

【重症度別集計】表

絞込条件

対象年月設定: 201404 ~ 201503
 病院の経営主体選択: 全て
 【詳細】病床数: 下限なし ~ 上限なし
 【詳細】手術の有無: 全て
 【詳細】ステージ集約選択: 有り
 【詳細】年代: 全て
 【詳細】性別: 全て
 【詳細】入院経路: 全て
 並べ替え条件: 退院年月・昇順

■ 死亡退院率

| | (単位:%) | 平均 | 退院年月 | | | | | | | |
|----------------------|--------|-------|---------|---------|---------|---------|---------|---------|---------|---------|
| | | | 2014/04 | 2014/05 | 2014/06 | 2014/07 | 2014/08 | 2014/09 | 2014/10 | 2014/11 |
| 胃の悪性新生物 | (人) | 696 | 72 | 63 | 46 | 58 | 70 | 62 | 60 | 54 |
| ステージ0 | | 6.3% | 33.3% | N/A | 0.0% | 0.0% | N/A | 0.0% | 0.0% | N/A |
| ステージI | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージII | | 4.1% | 10.0% | 0.0% | 0.0% | 0.0% | 0.0% | 9.1% | 14.3% | 0.0% |
| ステージIII | | 13.6% | 0.0% | 14.3% | 0.0% | 0.0% | 0.0% | 0.0% | 20.0% | 0.0% |
| ステージIV | | 19.3% | 29.6% | 8.0% | 27.3% | 14.3% | 13.8% | 34.8% | 9.5% | 25.0% |
| 不明 | | 13.3% | 28.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 12.5% |
| 計 | | 10.3% | 16.7% | 4.8% | 13.0% | 6.9% | 5.7% | 14.5% | 6.7% | 9.3% |
| 結腸の悪性新生物 | (人) | 866 | 83 | 75 | 83 | 76 | 80 | 70 | 70 | 70 |
| ステージ0 | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージI | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージII | | 1.5% | 0.0% | 0.0% | 0.0% | 9.1% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージIII | | 2.6% | 4.0% | 0.0% | 5.6% | 9.5% | 0.0% | 4.8% | 0.0% | 0.0% |
| ステージIV | | 9.7% | 3.4% | 8.3% | 10.8% | 8.3% | 10.5% | 15.0% | 12.5% | 12.5% |
| 不明 | | 10.0% | 42.9% | 0.0% | 12.5% | 0.0% | 0.0% | 16.7% | 0.0% | 16.7% |
| 計 | | 5.1% | 6.0% | 2.7% | 7.2% | 7.9% | 5.0% | 7.1% | 4.3% | 5.7% |
| 直腸の悪性新生物 | (人) | 453 | 34 | 29 | 29 | 32 | 49 | 36 | 39 | 44 |
| ステージ0 | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | N/A | 0.0% | N/A |
| ステージI | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージII | | 5.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 11.1% | 0.0% | 0.0% |
| ステージIII | | 1.4% | 0.0% | 0.0% | 0.0% | 9.1% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージIV | | 11.8% | 0.0% | 14.3% | 28.6% | 0.0% | 21.1% | 9.1% | 6.7% | 5.9% |
| 不明 | | 6.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | N/A | 33.3% | 0.0% |
| 計 | | 5.5% | 0.0% | 3.4% | 13.8% | 3.1% | 8.2% | 5.6% | 5.1% | 2.3% |
| 気管支および肺の悪性新生物 | (人) | 783 | 80 | 88 | 70 | 86 | 78 | 89 | 51 | 56 |
| ステージ0 | | 0.0% | 0.0% | 0.0% | 0.0% | N/A | N/A | N/A | N/A | N/A |
| ステージI | | 5.1% | 5.9% | 0.0% | 0.0% | 7.7% | 7.1% | 8.3% | 0.0% | 12.5% |
| ステージII | | 3.1% | 0.0% | 0.0% | 0.0% | 9.1% | 0.0% | 0.0% | 50.0% | 0.0% |
| ステージIII | | 7.6% | 11.8% | 0.0% | 13.3% | 15.0% | 12.5% | 4.5% | 0.0% | 8.3% |
| ステージIV | | 20.4% | 18.5% | 18.4% | 13.8% | 12.9% | 13.8% | 24.3% | 23.1% | 21.1% |
| 不明 | | 10.9% | 0.0% | 16.7% | 10.0% | 9.1% | 9.1% | 9.1% | 9.1% | 25.0% |
| 計 | | 12.3% | 10.0% | 11.4% | 10.0% | 11.6% | 10.3% | 13.5% | 9.8% | 16.1% |
| 急性心筋梗塞 | (人) | 266 | 14 | 19 | 20 | 20 | 28 | 20 | 29 | 25 |
| Killip I | | 1.8% | 0.0% | 20.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| Killip II | | 6.8% | 0.0% | 0.0% | 0.0% | 20.0% | 0.0% | 33.3% | 10.0% | 10.0% |
| Killip III | | 12.9% | N/A | 0.0% | 0.0% | 20.0% | 0.0% | 0.0% | 100.0% | 25.0% |
| Killip IV | | 40.0% | 50.0% | 25.0% | 50.0% | 50.0% | 33.3% | 0.0% | 60.0% | 50.0% |
| 不明 | | 60.0% | N/A | N/A | 0.0% | 100.0% | N/A | N/A | N/A | 50.0% |
| 計 | | 12.0% | 7.1% | 15.8% | 5.0% | 25.0% | 3.6% | 5.0% | 17.2% | 16.0% |
| 肺炎 | (人) | 2,889 | 270 | 283 | 272 | 241 | 229 | 183 | 190 | 175 |
| 軽症 | | 1.8% | 3.1% | 1.5% | 0.0% | 0.0% | 4.3% | 2.7% | 0.0% | 0.0% |
| 中等症 | | 6.6% | 0.0% | 12.5% | 15.8% | 0.0% | 11.1% | 10.0% | 9.5% | 0.0% |
| 重症 | | 14.1% | 17.2% | 14.6% | 15.7% | 10.9% | 10.2% | 16.7% | 10.4% | 20.4% |
| 不明 | | 2.1% | 4.3% | 0.9% | 0.9% | 3.0% | 1.3% | 0.0% | 6.6% | 1.7% |
| 計 | | 6.3% | 8.5% | 6.4% | 6.6% | 4.1% | 6.1% | 7.1% | 5.8% | 6.9% |
| 喘息 | (人) | 916 | 74 | 95 | 60 | 76 | 71 | 86 | 82 | 85 |
| 軽症間欠型 | | 0.4% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 軽症持続型 | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 中等症持続型 | | 0.5% | 0.0% | 0.0% | 0.0% | 5.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 重症持続型 | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 不明 | | 0.0% | N/A | 0.0% | N/A | 0.0% | N/A | 0.0% | 0.0% | N/A |
| 計 | | 0.2% | 0.0% | 0.0% | 0.0% | 1.3% | 0.0% | 0.0% | 0.0% | 0.0% |
| 脳梗塞 | (人) | 1,365 | 119 | 123 | 108 | 120 | 118 | 119 | 121 | 96 |
| JCS 0-1 | | 2.0% | 5.3% | 3.9% | 3.1% | 0.0% | 2.7% | 2.3% | 2.4% | 0.0% |
| JCS 2-3 | | 2.5% | 0.0% | 2.9% | 0.0% | 0.0% | 0.0% | 0.0% | 4.3% | 3.8% |
| JCS 10-100 | | 21.6% | 0.0% | 27.3% | 35.7% | 0.0% | 27.3% | 21.4% | 0.0% | 18.2% |

【重症度別集計】表

| | (単位:%) | 平均 | 退院年月 | | | |
|----------------------|--------|-------|---------|---------|---------|---------|
| | | | 2014/12 | 2015/01 | 2015/02 | 2015/03 |
| 胃の悪性新生物 | (人) | 696 | 62 | 54 | 45 | 50 |
| ステージ0 | | 6.3% | 0.0% | 0.0% | N/A | N/A |
| ステージI | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージII | | 4.1% | 14.3% | 0.0% | 0.0% | 0.0% |
| ステージIII | | 13.6% | 16.7% | 18.2% | 33.3% | 50.0% |
| ステージIV | | 19.3% | 13.6% | 17.6% | 23.1% | 18.8% |
| 不明 | | 13.3% | N/A | N/A | 16.7% | 50.0% |
| 計 | | 10.3% | 9.7% | 9.3% | 15.6% | 14.0% |
| 結腸の悪性新生物 | (人) | 866 | 67 | 56 | 81 | 55 |
| ステージ0 | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージI | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージII | | 1.5% | 0.0% | 5.9% | 0.0% | 0.0% |
| ステージIII | | 2.6% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージIV | | 9.7% | 9.5% | 6.7% | 9.1% | 9.1% |
| 不明 | | 10.0% | 0.0% | 0.0% | 16.7% | 0.0% |
| 計 | | 5.1% | 3.0% | 3.6% | 3.7% | 3.6% |
| 直腸の悪性新生物 | (人) | 453 | 44 | 46 | 32 | 39 |
| ステージ0 | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージI | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージII | | 5.3% | 0.0% | 0.0% | 0.0% | 23.1% |
| ステージIII | | 1.4% | 0.0% | 0.0% | 0.0% | 9.1% |
| ステージIV | | 11.8% | 8.3% | 12.5% | 15.4% | 16.7% |
| 不明 | | 6.7% | N/A | N/A | N/A | 0.0% |
| 計 | | 5.5% | 2.3% | 4.3% | 6.3% | 12.8% |
| 気管支および肺の悪性新生物 | (人) | 783 | 50 | 43 | 45 | 47 |
| ステージ0 | | 0.0% | N/A | N/A | N/A | N/A |
| ステージI | | 5.1% | 0.0% | 33.3% | 0.0% | 0.0% |
| ステージII | | 3.1% | 0.0% | N/A | 0.0% | 0.0% |
| ステージIII | | 7.6% | 0.0% | 0.0% | 11.1% | 6.7% |
| ステージIV | | 20.4% | 16.7% | 31.8% | 11.1% | 50.0% |
| 不明 | | 10.9% | 16.7% | 0.0% | 10.0% | 0.0% |
| 計 | | 12.3% | 10.0% | 18.6% | 8.9% | 21.3% |
| 急性心筋梗塞 | (人) | 266 | 34 | 21 | 22 | 14 |
| Killip I | | 1.8% | 0.0% | 0.0% | 0.0% | 0.0% |
| Killip II | | 6.8% | 12.5% | 0.0% | 0.0% | 0.0% |
| Killip III | | 12.9% | 0.0% | 0.0% | 0.0% | 100.0% |
| Killip IV | | 40.0% | 44.4% | 50.0% | 0.0% | 60.0% |
| 不明 | | 60.0% | 100.0% | N/A | N/A | N/A |
| 計 | | 12.0% | 17.6% | 4.8% | 0.0% | 28.6% |
| 肺炎 | (人) | 2,889 | 267 | 333 | 225 | 221 |
| 軽症 | | 1.8% | 4.3% | 1.4% | 1.7% | 2.0% |
| 中等症 | | 6.6% | 6.7% | 0.0% | 6.3% | 5.3% |
| 重症 | | 14.1% | 11.0% | 16.5% | 16.2% | 7.1% |
| 不明 | | 2.1% | 1.2% | 2.8% | 2.6% | 1.2% |
| 計 | | 6.3% | 5.6% | 7.5% | 7.1% | 3.6% |
| 喘息 | (人) | 916 | 66 | 63 | 60 | 98 |
| 軽症間欠型 | | 0.4% | 0.0% | 6.3% | 0.0% | 0.0% |
| 軽症持続型 | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 中等症持続型 | | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% |
| 重症持続型 | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 不明 | | 0.0% | N/A | N/A | N/A | 0.0% |
| 計 | | 0.2% | 0.0% | 1.6% | 0.0% | 0.0% |
| 脳梗塞 | (人) | 1,365 | 133 | 91 | 92 | 125 |
| JCS 0-1 | | 2.0% | 1.0% | 2.9% | 0.0% | 0.0% |
| JCS 2-3 | | 2.5% | 5.0% | 7.7% | 5.0% | 3.6% |
| JCS 10-100 | | 21.6% | 37.5% | 37.5% | 12.5% | 28.6% |

【重症度別集計】表

| | | 平均 | 退院年月 | | | |
|-----------|------------------------|-------|---------|---------|---------|---------|
| | | | 2014/12 | 2015/01 | 2015/02 | 2015/03 |
| 脳梗塞 | JCS 200-300 | 38.9% | 50.0% | 0.0% | 33.3% | 20.0% |
| | 不明 | N/A | N/A | N/A | N/A | N/A |
| | 計 | 4.8% | 5.3% | 6.6% | 3.3% | 4.8% |
| 脳出血 | (人) | 586 | 62 | 68 | 51 | 48 |
| | JCS 0-1 | 1.5% | 0.0% | 0.0% | 0.0% | 0.0% |
| | JCS 2-3 | 4.4% | 0.0% | 0.0% | 15.4% | 0.0% |
| | JCS 10-100 | 18.8% | 18.2% | 30.0% | 16.7% | 22.2% |
| | JCS 200-300 | 61.5% | 68.8% | 37.5% | 50.0% | 71.4% |
| | 不明 | N/A | N/A | N/A | N/A | N/A |
| | 計 | 14.7% | 21.0% | 8.8% | 13.7% | 14.6% |
| 糖尿病 | (人) | 585 | 54 | 40 | 55 | 39 |
| | 軽症 | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 中等症 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 重症 | 4.6% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 不明 | 0.0% | 0.0% | N/A | N/A | 0.0% |
| | 計 | 0.9% | 0.0% | 0.0% | 0.0% | 0.0% |
| 大腿骨頸部骨折 | (人) | 511 | 56 | 44 | 41 | 50 |
| | 重症度Ⅰ | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 重症度Ⅱ | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 重症度Ⅲ | 0.7% | 5.3% | 0.0% | 0.0% | 0.0% |
| | 重症度Ⅳ | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 不明 | 0.0% | 0.0% | 0.0% | 0.0% | N/A |
| | 計 | 0.2% | 1.8% | 0.0% | 0.0% | 0.0% |
| 胃潰瘍 | (人) | 202 | 19 | 18 | 16 | 15 |
| | ショックなし(収縮期血圧90mmHg以上) | 0.6% | 0.0% | 0.0% | 6.7% | 0.0% |
| | ショックあり(収縮期血圧90mmHg未満) | 0.0% | 0.0% | N/A | 0.0% | 0.0% |
| | 不明 | 0.0% | 0.0% | N/A | N/A | N/A |
| | 計 | 0.5% | 0.0% | 0.0% | 6.3% | 0.0% |
| 急性腸炎 | (人) | 701 | 72 | 53 | 48 | 49 |
| | 血便なし | 0.6% | 1.4% | 0.0% | 0.0% | 0.0% |
| | 血便あり | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 不明 | 0.0% | N/A | N/A | N/A | N/A |
| | 計 | 0.6% | 1.4% | 0.0% | 0.0% | 0.0% |
| 急性虫垂炎 | (人) | 522 | 41 | 33 | 31 | 43 |
| | 白血球15000/mm3未満 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 白血球15000/mm3以上 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 不明 | 0.0% | N/A | N/A | N/A | N/A |
| | 計 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 胆石症 | (人) | 1,108 | 107 | 74 | 83 | 88 |
| | 黄疸なし(T Bill 5mg/dl 未満) | 0.7% | 1.0% | 0.0% | 0.0% | 0.0% |
| | 黄疸あり(T Bill 5mg/dl 以上) | 2.6% | 0.0% | 0.0% | 0.0% | 16.7% |
| | 不明 | N/A | N/A | N/A | N/A | N/A |
| | 計 | 0.8% | 0.9% | 0.0% | 0.0% | 1.1% |
| 前立腺肥大症 | (人) | 95 | 10 | 7 | 6 | 4 |
| | 前立腺容積20ml未満 | 0.0% | 0.0% | 0.0% | 0.0% | N/A |
| | 前立腺容積50ml未満 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 前立腺容積50ml以上 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 不明 | 0.0% | 0.0% | N/A | N/A | N/A |
| | 計 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 白内障 | (人) | 679 | 44 | 40 | 48 | 56 |
| | 1眼(処置対象のもの) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 2眼(処置対象のもの) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 不明 | N/A | N/A | N/A | N/A | N/A |
| | 計 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 痔核 | (人) | 74 | 7 | 2 | 5 | 6 |
| | 第一度(出血のみ) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 第二度(還納可能) | 0.0% | 0.0% | 0.0% | N/A | N/A |
| | 第三度(脱肛) | 0.0% | 0.0% | N/A | 0.0% | 0.0% |
| | 不明 | 0.0% | N/A | N/A | N/A | N/A |
| | 計 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 子宮筋腫 | (人) | 142 | 10 | 8 | 13 | 11 |
| | 貧血なし(Hb10g/dl以上) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 貧血あり(Hb10g/dl未満) | 0.0% | N/A | 0.0% | 0.0% | 0.0% |
| | 不明 | 0.0% | N/A | N/A | N/A | N/A |
| | 計 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 狭心症 | (人) | 1,963 | 171 | 158 | 167 | 167 |
| | 重症度の指標なし | 0.2% | 0.6% | 1.3% | 0.0% | 0.0% |
| | 計 | 0.2% | 0.6% | 1.3% | 0.0% | 0.0% |
| 腎結石及び尿管結石 | (人) | 487 | 47 | 38 | 36 | 36 |
| | 水腎・水尿管なし | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 水腎・水尿管あり | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |

【重症度別集計】表

| | | 平均 | 退院年月 | | | |
|-----------|----------|-------|---------|---------|---------|---------|
| | | | 2014/12 | 2015/01 | 2015/02 | 2015/03 |
| 腎結石及び尿管結石 | 不明 | 0.0% | 0.0% | N/A | 0.0% | N/A |
| | 計 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 乳房の悪性新生物 | (人) | 493 | 51 | 37 | 50 | 41 |
| | ステージ0 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | ステージI | 0.6% | 0.0% | 0.0% | 0.0% | 0.0% |
| | ステージII | 2.8% | 4.3% | 0.0% | 6.7% | 0.0% |
| | ステージIII | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | ステージIV | 19.5% | 14.3% | 0.0% | 16.7% | 0.0% |
| | 不明 | 7.4% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 計 | 3.2% | 3.9% | 0.0% | 4.0% | 0.0% |
| 膝関節症 | (人) | 313 | 24 | 14 | 18 | 24 |
| | 骨移植なし | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 骨移植あり | N/A | N/A | N/A | N/A | N/A |
| | 不明 | N/A | N/A | N/A | N/A | N/A |
| | 計 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 鼠径ヘルニア | (人) | 727 | 63 | 47 | 60 | 58 |
| | 重症度の指標なし | 0.1% | 0.0% | 2.1% | 0.0% | 0.0% |
| | 計 | 0.1% | 0.0% | 2.1% | 0.0% | 0.0% |
| 全疾患合計 | | 3.7% | 3.9% | 4.3% | 3.4% | 3.6% |
| 集計病院数 | - | | 13 | 12 | 12 | 11 |