

【重症度別集計】表

絞込条件

対象年月設定: 202004 ~ 202103
 病院の経営主体選択: 全て
 【詳細】病床数: 下限なし ~ 上限なし
 【詳細】手術の有無: 全て
 【詳細】ステージ集約選択: 有り
 【詳細】年代: 全て
 【詳細】性別: 全て
 【詳細】入院経路: 全て
 並べ替え条件: 退院年月・昇順

■ 死亡退院率

| | (単位:%) | 平均 | 退院年月 | | | | | | | |
|----------------------|--------|-------|---------|---------|---------|---------|---------|---------|---------|---------|
| | | | 2020/04 | 2020/05 | 2020/06 | 2020/07 | 2020/08 | 2020/09 | 2020/10 | 2020/11 |
| 胃の悪性新生物 | (人) | - | - | - | - | - | - | - | - | - |
| ステージ0 | | 0.0% | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| ステージI | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージII | | 7.7% | 0.0% | 100.0% | N/A | 0.0% | N/A | 0.0% | N/A | N/A |
| ステージIII | | 18.2% | 100.0% | N/A | 0.0% | 100.0% | 0.0% | N/A | 0.0% | N/A |
| ステージIV | | 14.7% | 0.0% | 0.0% | 33.3% | 0.0% | 0.0% | 33.3% | 25.0% | 0.0% |
| 不明 | | 0.0% | N/A | N/A | 0.0% | N/A | N/A | 0.0% | 0.0% | N/A |
| 計 | | 6.8% | 8.3% | 16.7% | 10.0% | 14.3% | 0.0% | 7.7% | 6.3% | 0.0% |
| 結腸の悪性新生物 | (人) | - | - | - | - | - | - | - | - | - |
| ステージ0 | | 0.0% | 0.0% | N/A | 0.0% | 0.0% | 0.0% | N/A | 0.0% | N/A |
| ステージI | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージII | | 7.5% | 0.0% | 0.0% | 0.0% | 20.0% | 0.0% | 0.0% | 16.7% | 16.7% |
| ステージIII | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | N/A | 0.0% | 0.0% |
| ステージIV | | 6.3% | 11.1% | 20.0% | 0.0% | 0.0% | 0.0% | 20.0% | 0.0% | 0.0% |
| 不明 | | 25.0% | N/A | 0.0% | 0.0% | N/A | N/A | N/A | 0.0% | N/A |
| 計 | | 3.5% | 3.6% | 5.6% | 0.0% | 3.7% | 0.0% | 11.1% | 5.3% | 6.3% |
| 直腸の悪性新生物 | (人) | - | - | - | - | - | - | - | - | - |
| ステージ0 | | 0.0% | 0.0% | 0.0% | N/A | 0.0% | N/A | N/A | N/A | 0.0% |
| ステージI | | 3.2% | 0.0% | 0.0% | N/A | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージII | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | N/A | 0.0% | N/A | 0.0% |
| ステージIII | | 7.7% | 20.0% | 0.0% | 33.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージIV | | 5.1% | 0.0% | 0.0% | 0.0% | N/A | 0.0% | 22.2% | 0.0% | 0.0% |
| 不明 | | 0.0% | N/A | N/A | N/A | N/A | N/A | 0.0% | N/A | N/A |
| 計 | | 4.1% | 6.7% | 0.0% | 14.3% | 0.0% | 0.0% | 11.1% | 0.0% | 0.0% |
| 気管支および肺の悪性新生物 | (人) | - | - | - | - | - | - | - | - | - |
| ステージ0 | | 0.0% | 0.0% | 0.0% | 0.0% | N/A | N/A | N/A | N/A | N/A |
| ステージI | | 1.9% | 0.0% | N/A | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージII | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | N/A | N/A | N/A | 0.0% |
| ステージIII | | 3.8% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | N/A | 0.0% | 0.0% |
| ステージIV | | 8.2% | 0.0% | 0.0% | 25.0% | 0.0% | 0.0% | 25.0% | 0.0% | 0.0% |
| 不明 | | 15.6% | 0.0% | 50.0% | 0.0% | 25.0% | 20.0% | 25.0% | 0.0% | 0.0% |
| 計 | | 6.4% | 0.0% | 10.0% | 4.8% | 10.0% | 7.7% | 14.3% | 0.0% | 0.0% |
| 急性心筋梗塞 | (人) | - | - | - | - | - | - | - | - | - |
| Killip I | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| Killip II | | 0.0% | 0.0% | 0.0% | 0.0% | N/A | N/A | N/A | N/A | 0.0% |
| Killip III | | 16.7% | N/A | N/A | 0.0% | N/A | 0.0% | N/A | N/A | 100.0% |
| Killip IV | | 20.0% | 0.0% | 0.0% | 25.0% | 20.0% | 50.0% | 100.0% | 33.3% | 0.0% |
| 不明 | | 0.0% | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| 計 | | 6.9% | 0.0% | 0.0% | 7.7% | 8.3% | 16.7% | 20.0% | 12.5% | 12.5% |
| 肺炎 | (人) | - | - | - | - | - | - | - | - | - |
| 軽症 | | 9.5% | 23.1% | 14.3% | 13.3% | 9.1% | 0.0% | 0.0% | 0.0% | 16.7% |
| 中等症 | | 5.7% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 16.7% | 0.0% |
| 重症 | | 11.0% | 14.3% | 8.3% | 0.0% | 0.0% | 22.2% | 11.1% | 21.4% | 6.3% |
| 不明 | | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| 計 | | 9.8% | 16.7% | 9.5% | 7.1% | 4.0% | 9.1% | 3.8% | 10.3% | 10.0% |
| 喘息 | (人) | - | - | - | - | - | - | - | - | - |
| 軽症間欠型 | | 0.0% | 0.0% | N/A | N/A | N/A | N/A | 0.0% | N/A | 0.0% |
| 軽症持続型 | | 0.0% | 0.0% | N/A | N/A | N/A | N/A | 0.0% | N/A | 0.0% |
| 中等症持続型 | | 0.0% | 0.0% | N/A | N/A | N/A | 0.0% | 0.0% | 0.0% | 0.0% |
| 重症持続型 | | 40.0% | 100.0% | N/A | N/A | N/A | 0.0% | 0.0% | 100.0% | 0.0% |
| 不明 | | 0.0% | N/A | N/A | N/A | N/A | N/A | N/A | 0.0% | N/A |
| 計 | | 6.1% | 11.1% | N/A | N/A | N/A | 0.0% | 0.0% | 33.3% | 0.0% |
| 脳梗塞 | (人) | - | - | - | - | - | - | - | - | - |
| JCS 0-1 | | 1.5% | 2.6% | 0.0% | 0.0% | 0.0% | 0.0% | 1.8% | 0.0% | 2.9% |
| JCS 2-3 | | 2.2% | 0.0% | 0.0% | 3.3% | 4.8% | 0.0% | 15.0% | 0.0% | 0.0% |
| JCS 10-100 | | 19.6% | 0.0% | 33.3% | 0.0% | 50.0% | 0.0% | 25.0% | 0.0% | 16.7% |

【重症度別集計】表

| | (単位:%) | 平均 | 退院年月 | | | |
|---------------|--------|-------|---------|---------|---------|---------|
| | | | 2020/12 | 2021/01 | 2021/02 | 2021/03 |
| 胃の悪性新生物 | (人) | - | - | - | - | - |
| ステージ0 | | 0.0% | 0.0% | 0.0% | 0.0% | N/A |
| ステージI | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージII | | 7.7% | 0.0% | 0.0% | 0.0% | N/A |
| ステージIII | | 18.2% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージIV | | 14.7% | 40.0% | 0.0% | 0.0% | 0.0% |
| 不明 | | 0.0% | 0.0% | N/A | N/A | 0.0% |
| 計 | | 6.8% | 16.7% | 0.0% | 0.0% | 0.0% |
| 結腸の悪性新生物 | (人) | - | - | - | - | - |
| ステージ0 | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージI | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージII | | 7.5% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージIII | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| ステージIV | | 6.3% | 0.0% | 0.0% | 0.0% | 33.3% |
| 不明 | | 25.0% | N/A | 100.0% | N/A | N/A |
| 計 | | 3.5% | 0.0% | 4.8% | 0.0% | 10.0% |
| 直腸の悪性新生物 | (人) | - | - | - | - | - |
| ステージ0 | | 0.0% | 0.0% | N/A | 0.0% | N/A |
| ステージI | | 3.2% | 0.0% | 50.0% | N/A | 0.0% |
| ステージII | | 0.0% | 0.0% | 0.0% | 0.0% | N/A |
| ステージIII | | 7.7% | 0.0% | N/A | 0.0% | 0.0% |
| ステージIV | | 5.1% | 0.0% | 0.0% | 0.0% | 0.0% |
| 不明 | | 0.0% | N/A | N/A | N/A | N/A |
| 計 | | 4.1% | 0.0% | 20.0% | 0.0% | 0.0% |
| 気管支および肺の悪性新生物 | (人) | - | - | - | - | - |
| ステージ0 | | 0.0% | N/A | N/A | N/A | N/A |
| ステージI | | 1.9% | 16.7% | 0.0% | 0.0% | 0.0% |
| ステージII | | 0.0% | N/A | 0.0% | N/A | N/A |
| ステージIII | | 3.8% | 50.0% | 0.0% | 0.0% | 0.0% |
| ステージIV | | 8.2% | 0.0% | 16.7% | 33.3% | 0.0% |
| 不明 | | 15.6% | 0.0% | N/A | 0.0% | 0.0% |
| 計 | | 6.4% | 15.4% | 8.3% | 8.3% | 0.0% |
| 急性心筋梗塞 | (人) | - | - | - | - | - |
| Killip I | | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| Killip II | | 0.0% | 0.0% | N/A | 0.0% | N/A |
| Killip III | | 16.7% | 0.0% | 0.0% | N/A | N/A |
| Killip IV | | 20.0% | 33.3% | 0.0% | 0.0% | 0.0% |
| 不明 | | 0.0% | N/A | N/A | N/A | 0.0% |
| 計 | | 6.9% | 9.1% | 0.0% | 0.0% | 0.0% |
| 肺炎 | (人) | - | - | - | - | - |
| 軽症 | | 9.5% | 20.0% | 0.0% | 0.0% | 0.0% |
| 中等症 | | 5.7% | 0.0% | N/A | N/A | 25.0% |
| 重症 | | 11.0% | 16.7% | 25.0% | 0.0% | 0.0% |
| 不明 | | N/A | N/A | N/A | N/A | N/A |
| 計 | | 9.8% | 17.1% | 15.4% | 0.0% | 4.3% |
| 喘息 | (人) | - | - | - | - | - |
| 軽症間欠型 | | 0.0% | 0.0% | 0.0% | N/A | N/A |
| 軽症持続型 | | 0.0% | 0.0% | 0.0% | N/A | 0.0% |
| 中等症持続型 | | 0.0% | 0.0% | N/A | N/A | N/A |
| 重症持続型 | | 40.0% | N/A | N/A | N/A | N/A |
| 不明 | | 0.0% | N/A | N/A | N/A | N/A |
| 計 | | 6.1% | 0.0% | 0.0% | N/A | 0.0% |
| 脳梗塞 | (人) | - | - | - | - | - |
| JCS 0-1 | | 1.5% | 1.9% | 0.0% | 6.5% | 3.1% |
| JCS 2-3 | | 2.2% | 0.0% | 0.0% | 0.0% | 0.0% |
| JCS 10-100 | | 19.6% | 20.0% | 66.7% | 20.0% | 25.0% |

【重症度別集計】表

| (単位:%) | | 平均 | 退院年月 | | | | | | | |
|-----------|------------------------|-------|---------|---------|---------|---------|---------|---------|---------|---------|
| | | | 2020/04 | 2020/05 | 2020/06 | 2020/07 | 2020/08 | 2020/09 | 2020/10 | 2020/11 |
| 脳梗塞 | JCS 200-300 | 39.1% | N/A | 33.3% | 20.0% | 50.0% | 50.0% | N/A | N/A | 66.7% |
| | 不明 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 計 | 3.5% | 1.7% | 2.8% | 2.0% | 4.3% | 1.4% | 6.3% | 0.0% | 5.0% |
| 脳出血 | (人) | - | - | - | - | - | - | - | - | - |
| | JCS 0-1 | 1.6% | 0.0% | 0.0% | 5.6% | 0.0% | 0.0% | 0.0% | 0.0% | 6.7% |
| | JCS 2-3 | 2.2% | 0.0% | 0.0% | 11.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | JCS 10-100 | 11.9% | 0.0% | 0.0% | 14.3% | 20.0% | 0.0% | 0.0% | 50.0% | 0.0% |
| | JCS 200-300 | 54.3% | 57.1% | 0.0% | 57.1% | 100.0% | 25.0% | 66.7% | 37.5% | 50.0% |
| | 不明 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 計 | 12.3% | 14.8% | 0.0% | 22.9% | 10.7% | 4.3% | 13.8% | 13.8% | 13.3% |
| 糖尿病 | (人) | - | - | - | - | - | - | - | - | - |
| | 軽症 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | N/A | 0.0% | N/A | 0.0% |
| | 中等症 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 重症 | 0.0% | N/A | N/A | 0.0% | 0.0% | N/A | 0.0% | 0.0% | N/A |
| | 不明 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 計 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 大腿骨頸部骨折 | (人) | - | - | - | - | - | - | - | - | - |
| | 重症度Ⅰ | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 重症度Ⅱ | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 重症度Ⅲ | 0.0% | N/A | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 重症度Ⅳ | 14.3% | N/A | 0.0% | N/A | N/A | N/A | 0.0% | 0.0% | 50.0% |
| | 不明 | 0.0% | N/A | N/A | N/A | 0.0% | N/A | N/A | N/A | N/A |
| | 計 | 0.6% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 8.3% |
| 胃潰瘍 | (人) | - | - | - | - | - | - | - | - | - |
| | ショックなし(収縮期血圧90mmHg以上) | 5.9% | 0.0% | 0.0% | 25.0% | 0.0% | 0.0% | 0.0% | 33.3% | N/A |
| | ショックあり(収縮期血圧90mmHg未満) | 0.0% | N/A | 0.0% | N/A | 0.0% | N/A | 0.0% | N/A | 0.0% |
| | 不明 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 計 | 4.7% | 0.0% | 0.0% | 25.0% | 0.0% | 0.0% | 0.0% | 33.3% | 0.0% |
| 急性腸炎 | (人) | - | - | - | - | - | - | - | - | - |
| | 血便なし | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 血便あり | 0.0% | N/A | N/A | N/A | N/A | 0.0% | N/A | N/A | 0.0% |
| | 不明 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 計 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 急性虫垂炎 | (人) | - | - | - | - | - | - | - | - | - |
| | 白血球15000/mm3未満 | 0.0% | 0.0% | N/A | 0.0% | 0.0% | N/A | 0.0% | 0.0% | 0.0% |
| | 白血球15000/mm3以上 | 0.0% | N/A | N/A | 0.0% | 0.0% | N/A | 0.0% | N/A | 0.0% |
| | 不明 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 計 | 0.0% | 0.0% | N/A | 0.0% | 0.0% | N/A | 0.0% | 0.0% | 0.0% |
| 胆石症 | (人) | - | - | - | - | - | - | - | - | - |
| | 黄疸なし(T Bill 5mg/dl 未満) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 黄疸あり(T Bill 5mg/dl 以上) | 5.3% | 0.0% | 0.0% | N/A | 0.0% | N/A | 50.0% | 0.0% | 0.0% |
| | 不明 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 計 | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 4.0% | 0.0% | 0.0% |
| 前立腺肥大症 | (人) | - | - | - | - | - | - | - | - | - |
| | 前立腺容積20ml未満 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 前立腺容積50ml未満 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 前立腺容積50ml以上 | 0.0% | 0.0% | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 不明 | 0.0% | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 計 | 0.0% | 0.0% | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| 白内障 | (人) | - | - | - | - | - | - | - | - | - |
| | 1眼(処置対象のもの) | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 2眼(処置対象のもの) | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 不明 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 計 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| 痔核 | (人) | - | - | - | - | - | - | - | - | - |
| | 第一度(出血のみ) | 0.0% | N/A | N/A | 0.0% | N/A | N/A | N/A | N/A | N/A |
| | 第二度(還納可能) | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 第三度(脱肛) | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 不明 | 0.0% | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 計 | 0.0% | N/A | N/A | 0.0% | N/A | N/A | N/A | N/A | N/A |
| 子宮筋腫 | (人) | - | - | - | - | - | - | - | - | - |
| | 貧血なし(Hb10g/dl以上) | 0.0% | 0.0% | N/A | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 貧血あり(Hb10g/dl未満) | 0.0% | N/A | N/A | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 不明 | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A | N/A |
| | 計 | 0.0% | 0.0% | N/A | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 狭心症 | (人) | 434 | 20 | 17 | 45 | 52 | 33 | 40 | 44 | 44 |
| | 重症度の指標なし | 0.5% | 0.0% | 0.0% | 0.0% | 1.9% | 0.0% | 0.0% | 0.0% | 2.3% |
| | 計 | 0.5% | 0.0% | 0.0% | 0.0% | 1.9% | 0.0% | 0.0% | 0.0% | 2.3% |
| 腎結石及び尿管結石 | (人) | - | - | - | - | - | - | - | - | - |
| | 水腎・水尿管なし | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | N/A | N/A |
| | 水腎・水尿管あり | 0.0% | N/A | 0.0% | 0.0% | 0.0% | N/A | N/A | N/A | N/A |

【重症度別集計】表

| | | 平均 | 退院年月 | | | |
|-----------|------------------------|--------|---------|---------|---------|---------|
| | | | 2020/12 | 2021/01 | 2021/02 | 2021/03 |
| | | (単位:%) | | | | |
| 脳梗塞 | JCS 200-300 | 39.1% | 100.0% | 0.0% | 0.0% | 66.7% |
| | 不明 | N/A | N/A | N/A | N/A | N/A |
| | 計 | 3.5% | 4.1% | 2.5% | 6.5% | 6.8% |
| 脳出血 | | (人) | - | - | - | - |
| | JCS 0-1 | 1.6% | 0.0% | 0.0% | 0.0% | 6.3% |
| | JCS 2-3 | 2.2% | 0.0% | 7.7% | 0.0% | 0.0% |
| | JCS 10-100 | 11.9% | 40.0% | 0.0% | 0.0% | 0.0% |
| | JCS 200-300 | 54.3% | 40.0% | 55.6% | 100.0% | 50.0% |
| | 不明 | N/A | N/A | N/A | N/A | N/A |
| | 計 | 12.3% | 10.0% | 14.3% | 9.5% | 12.0% |
| 糖尿病 | | (人) | - | - | - | - |
| | 軽症 | 0.0% | 0.0% | N/A | 0.0% | 0.0% |
| | 中等症 | 0.0% | 0.0% | 0.0% | N/A | 0.0% |
| | 重症 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 不明 | N/A | N/A | N/A | N/A | N/A |
| | 計 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 大腿骨頸部骨折 | | (人) | - | - | - | - |
| | 重症度Ⅰ | 0.0% | 0.0% | 0.0% | 0.0% | N/A |
| | 重症度Ⅱ | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 重症度Ⅲ | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 重症度Ⅳ | 14.3% | N/A | 0.0% | N/A | N/A |
| | 不明 | 0.0% | N/A | N/A | N/A | N/A |
| | 計 | 0.6% | 0.0% | 0.0% | 0.0% | 0.0% |
| 胃潰瘍 | | (人) | - | - | - | - |
| | ショックなし(収縮期血圧90mmHg以上) | 5.9% | 0.0% | 0.0% | 0.0% | 0.0% |
| | ショックあり(収縮期血圧90mmHg未満) | 0.0% | N/A | 0.0% | 0.0% | N/A |
| | 不明 | N/A | N/A | N/A | N/A | N/A |
| | 計 | 4.7% | 0.0% | 0.0% | 0.0% | 0.0% |
| 急性腸炎 | | (人) | - | - | - | - |
| | 血便なし | 0.0% | 0.0% | 0.0% | 0.0% | N/A |
| | 血便あり | 0.0% | 0.0% | 0.0% | N/A | N/A |
| | 不明 | N/A | N/A | N/A | N/A | N/A |
| | 計 | 0.0% | 0.0% | 0.0% | 0.0% | N/A |
| 急性虫垂炎 | | (人) | - | - | - | - |
| | 白血球15000/mm3未満 | 0.0% | 0.0% | 0.0% | N/A | 0.0% |
| | 白血球15000/mm3以上 | 0.0% | 0.0% | N/A | N/A | N/A |
| | 不明 | N/A | N/A | N/A | N/A | N/A |
| | 計 | 0.0% | 0.0% | 0.0% | N/A | 0.0% |
| 胆石症 | | (人) | - | - | - | - |
| | 黄疸なし(T Bill 5mg/dl 未満) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 黄疸あり(T Bill 5mg/dl 以上) | 5.3% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 不明 | N/A | N/A | N/A | N/A | N/A |
| | 計 | 0.3% | 0.0% | 0.0% | 0.0% | 0.0% |
| 前立腺肥大症 | | (人) | - | - | - | - |
| | 前立腺容積20ml未満 | N/A | N/A | N/A | N/A | N/A |
| | 前立腺容積50ml未満 | N/A | N/A | N/A | N/A | N/A |
| | 前立腺容積50ml以上 | 0.0% | N/A | N/A | N/A | N/A |
| | 不明 | 0.0% | N/A | N/A | 0.0% | N/A |
| | 計 | 0.0% | N/A | N/A | 0.0% | N/A |
| 白内障 | | (人) | - | - | - | - |
| | 1眼(処置対象のもの) | N/A | N/A | N/A | N/A | N/A |
| | 2眼(処置対象のもの) | N/A | N/A | N/A | N/A | N/A |
| | 不明 | N/A | N/A | N/A | N/A | N/A |
| | 計 | N/A | N/A | N/A | N/A | N/A |
| 痔核 | | (人) | - | - | - | - |
| | 第一度(出血のみ) | 0.0% | N/A | N/A | 0.0% | N/A |
| | 第二度(還納可能) | N/A | N/A | N/A | N/A | N/A |
| | 第三度(脱肛) | N/A | N/A | N/A | N/A | N/A |
| | 不明 | 0.0% | 0.0% | N/A | N/A | N/A |
| | 計 | 0.0% | 0.0% | N/A | 0.0% | N/A |
| 子宮筋腫 | | (人) | - | - | - | - |
| | 貧血なし(Hb10g/dl以上) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 貧血あり(Hb10g/dl未満) | 0.0% | N/A | 0.0% | N/A | 0.0% |
| | 不明 | N/A | N/A | N/A | N/A | N/A |
| | 計 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 狭心症 | | (人) | 434 | 41 | 40 | 35 |
| | 重症度の指標なし | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 計 | 0.5% | 0.0% | 0.0% | 0.0% | 0.0% |
| 腎結石及び尿管結石 | | (人) | - | - | - | - |
| | 水腎・水尿管なし | 0.0% | N/A | N/A | 0.0% | N/A |
| | 水腎・水尿管あり | 0.0% | N/A | 0.0% | N/A | N/A |

【重症度別集計】表

| | | 平均 | 退院年月 | | | |
|-----------|----------|------|---------|---------|---------|---------|
| | | | 2020/12 | 2021/01 | 2021/02 | 2021/03 |
| 腎結石及び尿管結石 | 不明 | N/A | N/A | N/A | N/A | N/A |
| | 計 | 0.0% | N/A | 0.0% | 0.0% | N/A |
| 乳房の悪性新生物 | (人) | - | - | - | - | - |
| | ステージ0 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | ステージI | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | ステージII | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | ステージIII | 0.0% | 0.0% | 0.0% | N/A | 0.0% |
| | ステージIV | 0.0% | N/A | 0.0% | N/A | 0.0% |
| | 不明 | 0.0% | N/A | N/A | N/A | 0.0% |
| | 計 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 膝関節症 | (人) | - | - | - | - | - |
| | 骨移植なし | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 骨移植あり | N/A | N/A | N/A | N/A | N/A |
| | 不明 | N/A | N/A | N/A | N/A | N/A |
| | 計 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 鼠径ヘルニア | (人) | 76 | 4 | 3 | 8 | 5 |
| | 重症度の指標なし | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| | 計 | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| 全疾患合計 | | 4.1% | 5.3% | 4.2% | 3.1% | 3.6% |
| 集計病院数 | | - | 3 | 2 | 2 | 2 |