

【重症度別集計】表

絞込条件

対象年月設定: 201304 ~ 201403
 病院の経営主体選択: 全て
 【詳細】病床数: 下限なし ~ 上限なし
 【詳細】手術の有無: 全て
 【詳細】ステージ集約選択: 有り
 【詳細】年代: 全て
 【詳細】性別: 全て
 【詳細】入院経路: 全て
 並べ替え条件: 退院年月・昇順

■ 退院患者数

(単位: %)	平均	退院年月								
		2013/04	2013/05	2013/06	2013/07	2013/08	2013/09	2013/10	2013/11	
胃の悪性新生物	(人)	969	88	91	96	92	65	70	76	80
	ステージ0	1.2%	0.0%	2.2%	1.0%	2.2%	0.0%	0.0%	1.3%	2.5%
	ステージI	26.7%	26.1%	24.2%	28.1%	32.6%	30.8%	25.7%	23.7%	23.8%
	ステージII	10.6%	13.6%	9.9%	3.1%	7.6%	7.7%	21.4%	5.3%	12.5%
	ステージIII	13.9%	12.5%	18.7%	17.7%	17.4%	16.9%	14.3%	10.5%	10.0%
	ステージIV	36.0%	34.1%	33.0%	33.3%	27.2%	38.5%	25.7%	47.4%	41.3%
	不明	11.5%	13.6%	12.1%	16.7%	13.0%	6.2%	12.9%	11.8%	10.0%
結腸の悪性新生物	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	1,187	111	98	73	122	102	100	94	99
	ステージ0	6.3%	8.1%	9.2%	8.2%	4.1%	4.9%	8.0%	3.2%	6.1%
	ステージI	10.6%	8.1%	9.2%	11.0%	12.3%	8.8%	15.0%	5.3%	13.1%
	ステージII	17.6%	12.6%	16.3%	19.2%	18.0%	16.7%	14.0%	13.8%	18.2%
	ステージIII	22.0%	18.9%	15.3%	19.2%	18.0%	21.6%	26.0%	30.9%	18.2%
	ステージIV	30.0%	33.3%	37.8%	38.4%	28.7%	36.3%	27.0%	31.9%	25.3%
直腸の悪性新生物	不明	13.5%	18.9%	12.2%	4.1%	18.9%	11.8%	10.0%	14.9%	19.2%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	557	42	62	51	46	46	33	58	48
	ステージ0	4.8%	7.1%	1.6%	2.0%	2.2%	6.5%	0.0%	3.4%	8.3%
	ステージI	12.7%	7.1%	14.5%	11.8%	13.0%	8.7%	9.1%	8.6%	18.8%
	ステージII	16.3%	11.9%	17.7%	17.6%	19.6%	10.9%	18.2%	17.2%	20.8%
	ステージIII	32.1%	21.4%	29.0%	35.3%	45.7%	37.0%	33.3%	37.9%	29.2%
気管支および肺の悪性新生物	ステージIV	25.0%	38.1%	33.9%	25.5%	17.4%	26.1%	30.3%	22.4%	18.8%
	不明	9.0%	14.3%	3.2%	7.8%	2.2%	10.9%	9.1%	10.3%	4.2%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	1,312	142	119	106	146	113	106	113	99
	ステージ0	0.5%	0.0%	0.8%	0.0%	0.7%	0.9%	0.9%	0.0%	0.0%
	ステージI	16.5%	16.2%	16.0%	13.2%	15.1%	15.9%	15.1%	23.9%	16.2%
	ステージII	7.9%	6.3%	5.0%	6.6%	3.4%	8.0%	7.5%	14.2%	9.1%
急性心筋梗塞	ステージIII	19.4%	17.6%	18.5%	19.8%	19.2%	19.5%	17.0%	20.4%	21.2%
	ステージIV	38.8%	39.4%	40.3%	41.5%	45.9%	40.7%	41.5%	25.7%	41.4%
	不明	16.9%	20.4%	19.3%	18.9%	15.8%	15.0%	17.9%	15.9%	12.1%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	370	30	32	43	31	24	16	26	43
	Killip I	43.5%	36.7%	46.9%	53.5%	35.5%	29.2%	43.8%	42.3%	46.5%
	Killip II	32.7%	33.3%	37.5%	32.6%	41.9%	33.3%	43.8%	30.8%	39.5%
肺炎	Killip III	8.1%	16.7%	6.3%	4.7%	12.9%	12.5%	12.5%	7.7%	0.0%
	Killip IV	12.4%	13.3%	3.1%	9.3%	9.7%	16.7%	0.0%	15.4%	9.3%
	不明	3.2%	0.0%	6.3%	0.0%	0.0%	8.3%	0.0%	3.8%	4.7%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	3,253	295	361	297	293	222	198	198	196
	軽症	27.6%	31.2%	25.2%	21.5%	22.2%	25.2%	29.3%	40.4%	26.0%
	中等症	8.8%	7.1%	8.6%	10.4%	7.2%	9.0%	9.1%	6.1%	12.2%
喘息	重症	33.8%	32.2%	34.3%	29.6%	33.4%	33.8%	36.9%	36.9%	37.2%
	不明	29.9%	29.5%	31.9%	38.4%	37.2%	32.0%	24.7%	16.7%	24.5%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	1,141	94	119	106	113	96	118	100	105
	軽症間欠型	26.6%	30.9%	30.3%	31.1%	31.9%	22.9%	38.1%	22.0%	16.2%
	軽症持続型	36.6%	31.9%	39.5%	35.8%	35.4%	44.8%	33.1%	22.0%	35.2%
	中等症持続型	22.6%	20.2%	20.2%	23.6%	11.5%	17.7%	18.6%	33.0%	34.3%
脳梗塞	重症持続型	11.9%	12.8%	5.9%	9.4%	16.8%	13.5%	9.3%	19.0%	14.3%
	不明	2.2%	4.3%	4.2%	0.0%	4.4%	1.0%	0.8%	4.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	1,735	155	150	143	138	125	138	161	143
	JCS 0-1	70.5%	66.5%	70.0%	72.0%	68.8%	73.6%	78.3%	64.6%	67.1%
	JCS 2-3	18.2%	19.4%	18.0%	16.1%	15.2%	16.0%	13.8%	24.2%	21.7%
	JCS 10-100	8.8%	11.0%	8.7%	9.1%	13.8%	8.8%	5.8%	9.3%	8.4%

【重症度別集計】表

	(単位:%)	平均	退院年月			
			2013/12	2014/01	2014/02	2014/03
胃の悪性新生物	(人)	969	79	82	68	82
	ステージ0	1.2%	1.3%	2.4%	0.0%	1.2%
	ステージI	26.7%	24.1%	20.7%	33.8%	28.0%
	ステージII	10.6%	8.9%	13.4%	14.7%	12.2%
	ステージIII	13.9%	12.7%	13.4%	10.3%	11.0%
	ステージIV	36.0%	45.6%	35.4%	36.8%	36.6%
	不明	11.5%	7.6%	14.6%	4.4%	11.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%
結腸の悪性新生物	(人)	1,187	107	96	84	101
	ステージ0	6.3%	5.6%	7.3%	4.8%	6.9%
	ステージI	10.6%	10.3%	11.5%	10.7%	11.9%
	ステージII	17.6%	21.5%	18.8%	25.0%	18.8%
	ステージIII	22.0%	29.9%	20.8%	22.6%	22.8%
	ステージIV	30.0%	25.2%	29.2%	22.6%	25.7%
	不明	13.5%	7.5%	12.5%	14.3%	13.9%
計		100.0%	100.0%	100.0%	100.0%	100.0%
直腸の悪性新生物	(人)	557	47	42	40	42
	ステージ0	4.8%	4.3%	2.4%	7.5%	14.3%
	ステージI	12.7%	14.9%	26.2%	10.0%	9.5%
	ステージII	16.3%	23.4%	7.1%	22.5%	7.1%
	ステージIII	32.1%	23.4%	21.4%	32.5%	38.1%
	ステージIV	25.0%	14.9%	31.0%	20.0%	21.4%
	不明	9.0%	19.1%	11.9%	7.5%	9.5%
計		100.0%	100.0%	100.0%	100.0%	100.0%
気管支および肺の悪性新生物	(人)	1,312	117	88	77	86
	ステージ0	0.5%	0.9%	0.0%	2.6%	0.0%
	ステージI	16.5%	24.8%	13.6%	13.0%	11.6%
	ステージII	7.9%	6.8%	9.1%	15.6%	7.0%
	ステージIII	19.4%	21.4%	22.7%	15.6%	20.9%
	ステージIV	38.8%	32.5%	35.2%	33.8%	45.3%
	不明	16.9%	13.7%	19.3%	19.5%	15.1%
計		100.0%	100.0%	100.0%	100.0%	100.0%
急性心筋梗塞	(人)	370	33	30	26	36
	Killip I	43.5%	45.5%	46.7%	69.2%	25.0%
	Killip II	32.7%	21.2%	13.3%	15.4%	47.2%
	Killip III	8.1%	3.0%	10.0%	7.7%	11.1%
	Killip IV	12.4%	24.2%	20.0%	7.7%	16.7%
	不明	3.2%	6.1%	10.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
肺炎	(人)	3,253	274	294	330	295
	軽症	27.6%	28.5%	27.9%	29.1%	28.5%
	中等症	8.8%	9.5%	9.2%	9.4%	7.8%
	重症	33.8%	36.1%	31.3%	32.7%	33.9%
	不明	29.9%	25.9%	31.6%	28.8%	29.8%
計		100.0%	100.0%	100.0%	100.0%	100.0%
喘息	(人)	1,141	59	60	74	97
	軽症間欠型	26.6%	10.2%	18.3%	18.9%	34.0%
	軽症持続型	36.6%	40.7%	40.0%	48.6%	39.2%
	中等症持続型	22.6%	32.2%	31.7%	18.9%	17.5%
	重症持続型	11.9%	11.9%	10.0%	13.5%	7.2%
	不明	2.2%	5.1%	0.0%	0.0%	2.1%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
脳梗塞	(人)	1,735	139	152	135	156
	JCS 0-1	70.5%	75.5%	77.6%	68.9%	65.4%
	JCS 2-3	18.2%	12.9%	13.2%	21.5%	24.4%
	JCS 10-100	8.8%	7.2%	7.2%	8.1%	7.7%

【重症度別集計】表

(単位: %)		平均	退院年月							
			2013/04	2013/05	2013/06	2013/07	2013/08	2013/09	2013/10	2013/11
脳梗塞	JCS 200-300	2.5%	3.2%	3.3%	2.8%	2.2%	1.6%	2.2%	1.9%	2.8%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
脳出血	(人)	667	61	66	55	53	45	41	48	35
	JCS 0-1	45.3%	49.2%	51.5%	40.0%	56.6%	26.7%	53.7%	39.6%	34.3%
	JCS 2-3	17.2%	19.7%	15.2%	20.0%	11.3%	17.8%	4.9%	18.8%	14.3%
	JCS 10-100	18.7%	16.4%	16.7%	23.6%	24.5%	26.7%	17.1%	14.6%	22.9%
	JCS 200-300	18.7%	14.8%	16.7%	16.4%	7.5%	28.9%	24.4%	27.1%	28.6%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
糖尿病	(人)	855	98	84	88	85	64	58	62	66
	軽症	62.1%	58.2%	54.8%	51.1%	58.8%	70.3%	70.7%	67.7%	60.6%
	中等症	14.0%	8.2%	10.7%	15.9%	12.9%	9.4%	13.8%	21.0%	18.2%
	重症	10.9%	10.2%	4.8%	8.0%	5.9%	9.4%	13.8%	9.7%	13.6%
	不明	13.0%	23.5%	29.8%	25.0%	22.4%	10.9%	1.7%	1.6%	7.6%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
大腿骨頸部骨折	(人)	648	64	52	56	66	44	52	40	51
	重症度 I	41.7%	40.6%	34.6%	50.0%	42.4%	34.1%	59.6%	37.5%	41.2%
	重症度 II	16.5%	18.8%	5.8%	19.6%	13.6%	22.7%	13.5%	7.5%	11.8%
	重症度 III	24.7%	18.8%	23.1%	14.3%	21.2%	20.5%	17.3%	40.0%	29.4%
	重症度 IV	5.6%	1.6%	3.8%	5.4%	4.5%	15.9%	7.7%	7.5%	7.8%
	不明	11.6%	20.3%	32.7%	10.7%	18.2%	6.8%	1.9%	7.5%	9.8%
胃潰瘍	(人)	252	19	25	16	27	24	19	23	20
	ショックなし(収縮期血圧90mmHg以上)	83.3%	73.7%	76.0%	87.5%	77.8%	75.0%	78.9%	91.3%	80.0%
	ショックあり(収縮期血圧90mmHg未満)	10.3%	15.8%	8.0%	0.0%	3.7%	16.7%	21.1%	8.7%	20.0%
	不明	6.3%	10.5%	16.0%	12.5%	18.5%	8.3%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
急性腸炎	(人)	896	88	90	57	76	98	59	77	68
	血便なし	88.6%	92.0%	87.8%	80.7%	90.8%	91.8%	79.7%	96.1%	94.1%
	血便あり	8.0%	2.3%	8.9%	10.5%	3.9%	7.1%	18.6%	3.9%	5.9%
	不明	3.3%	5.7%	3.3%	8.8%	5.3%	1.0%	1.7%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
急性虫垂炎	(人)	703	59	63	73	58	60	66	73	49
	白血球15000/mm ³ 未満	68.4%	71.2%	66.7%	72.6%	67.2%	73.3%	81.8%	68.5%	59.2%
	白血球15000/mm ³ 以上	28.2%	18.6%	28.6%	20.5%	17.2%	26.7%	18.2%	30.1%	40.8%
	不明	3.4%	10.2%	4.8%	6.8%	15.5%	0.0%	0.0%	1.4%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
胆石症	(人)	1,413	129	117	127	111	108	124	132	136
	黄疸なし(T Bill 5mg/dl 未満)	92.0%	87.6%	90.6%	89.8%	87.4%	90.7%	95.2%	93.2%	92.6%
	黄疸あり(T Bill 5mg/dl 以上)	5.7%	4.7%	5.1%	3.9%	4.5%	9.3%	4.8%	6.8%	7.4%
	不明	2.3%	7.8%	4.3%	6.3%	8.1%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
前立腺肥大症	(人)	189	12	23	16	22	20	17	10	23
	前立腺容積20ml未満	25.4%	50.0%	17.4%	25.0%	18.2%	25.0%	23.5%	30.0%	26.1%
	前立腺容積50ml未満	25.9%	8.3%	21.7%	25.0%	27.3%	20.0%	17.6%	40.0%	43.5%
	前立腺容積50ml以上	16.4%	8.3%	13.0%	25.0%	9.1%	15.0%	11.8%	20.0%	26.1%
	不明	32.3%	33.3%	47.8%	25.0%	45.5%	40.0%	47.1%	10.0%	4.3%
白内障	(人)	1,194	116	103	132	135	97	75	106	101
	1眼(処置対象のもの)	58.7%	44.0%	41.7%	41.7%	47.4%	67.0%	62.7%	70.8%	75.2%
	2眼(処置対象のもの)	28.9%	26.7%	28.2%	30.3%	19.3%	33.0%	36.0%	29.2%	24.8%
	不明	12.4%	29.3%	30.1%	28.0%	33.3%	0.0%	1.3%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
痔核	(人)	140	14	13	13	10	13	4	8	15
	第一度(出血のみ)	27.1%	7.1%	30.8%	7.7%	30.0%	30.8%	50.0%	50.0%	40.0%
	第二度(還納可能)	17.9%	7.1%	15.4%	23.1%	20.0%	23.1%	25.0%	12.5%	20.0%
	第三度(脱肛)	52.1%	85.7%	38.5%	61.5%	50.0%	46.2%	25.0%	37.5%	40.0%
	不明	2.9%	0.0%	15.4%	7.7%	0.0%	0.0%	0.0%	0.0%	0.0%
子宮筋腫	(人)	476	41	37	46	51	51	36	37	37
	貧血なし(Hb10g/dl以上)	71.2%	61.0%	59.5%	58.7%	54.9%	70.6%	83.3%	78.4%	83.8%
	貧血あり(Hb10g/dl未満)	21.6%	19.5%	21.6%	21.7%	27.5%	29.4%	16.7%	21.6%	13.5%
	不明	7.1%	19.5%	18.9%	19.6%	17.6%	0.0%	0.0%	0.0%	2.7%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
狭心症	(人)	2,070	213	211	242	202	175	121	160	172
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
腎結石及び尿管結石	(人)	632	55	57	63	56	71	49	51	54
	水腎・水尿管なし	49.7%	54.5%	50.9%	46.0%	41.1%	59.2%	55.1%	58.8%	48.1%
	水腎・水尿管あり	44.9%	30.9%	43.9%	39.7%	46.4%	39.4%	44.9%	37.3%	50.0%

【重症度別集計】表

(単位:%)		平均	退院年月			
			2013/12	2014/01	2014/02	2014/03
脳梗塞	JCS 200-300	2.5%	4.3%	2.0%	1.5%	2.6%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
脳出血	(人)	667	48	72	62	81
	JCS 0-1	45.3%	43.8%	30.6%	46.8%	60.5%
	JCS 2-3	17.2%	18.8%	29.2%	22.6%	9.9%
	JCS 10-100	18.7%	14.6%	25.0%	11.3%	14.8%
	JCS 200-300	18.7%	22.9%	15.3%	19.4%	14.8%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
糖尿病	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	855	59	54	61	76
	軽症	62.1%	64.4%	66.7%	63.9%	68.4%
	中等症	14.0%	15.3%	16.7%	16.4%	14.5%
	重症	10.9%	18.6%	11.1%	14.8%	15.8%
	不明	13.0%	1.7%	5.6%	4.9%	1.3%
大腿骨頸部骨折	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	648	64	55	52	52
	重症度 I	41.7%	43.8%	30.9%	30.8%	51.9%
	重症度 II	16.5%	17.2%	25.5%	25.0%	15.4%
	重症度 III	24.7%	28.1%	29.1%	30.8%	28.8%
	重症度 IV	5.6%	6.3%	5.5%	1.9%	1.9%
腎潰瘍	不明	11.6%	4.7%	9.1%	11.5%	1.9%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	252	20	18	21	20
	ショックなし (収縮期血圧90mmHg以上)	83.3%	95.0%	88.9%	85.7%	95.0%
	ショックあり (収縮期血圧90mmHg未満)	10.3%	0.0%	11.1%	14.3%	5.0%
	不明	6.3%	5.0%	0.0%	0.0%	0.0%
急性腸炎	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	896	90	70	52	71
	血便なし	88.6%	77.8%	82.9%	94.2%	94.4%
	血便あり	8.0%	14.4%	12.9%	3.8%	5.6%
	不明	3.3%	7.8%	4.3%	1.9%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
急性虫垂炎	(人)	703	71	41	42	48
	白血球15000/mm ³ 未満	68.4%	53.5%	68.3%	61.9%	75.0%
	白血球15000/mm ³ 以上	28.2%	46.5%	31.7%	38.1%	25.0%
	不明	3.4%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	1,413	102	101	96	130
胆石症	黄疸なし(T Bill 5mg/dl 未満)	92.0%	94.1%	94.1%	93.8%	95.4%
	黄疸あり(T Bill 5mg/dl 以上)	5.7%	5.9%	5.9%	6.3%	4.6%
	不明	2.3%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	189	20	11	10	5
	前立腺容積20ml未満	25.4%	35.0%	18.2%	20.0%	20.0%
前立腺肥大症	前立腺容積50ml未満	25.9%	10.0%	45.5%	30.0%	40.0%
	前立腺容積50ml以上	16.4%	25.0%	18.2%	10.0%	0.0%
	不明	32.3%	30.0%	18.2%	40.0%	40.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	1,194	85	77	86	81
	1眼(処置対象のもの)	58.7%	65.9%	70.1%	75.6%	61.7%
白内障	2眼(処置対象のもの)	28.9%	34.1%	29.9%	24.4%	38.3%
	不明	12.4%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	140	11	10	13	16
	第一度(出血のみ)	27.1%	36.4%	30.0%	23.1%	18.8%
	第二度(還納可能)	17.9%	0.0%	40.0%	7.7%	25.0%
痔核	第三度(脱肛)	52.1%	63.6%	30.0%	69.2%	50.0%
	不明	2.9%	0.0%	0.0%	0.0%	6.3%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	476	46	21	34	39
	貧血なし(Hb10g/dl以上)	71.2%	71.7%	66.7%	85.3%	89.7%
	貧血あり(Hb10g/dl未満)	21.6%	28.3%	33.3%	14.7%	10.3%
子宮筋腫	不明	7.1%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	2,070	143	145	144	142
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%
狭心症	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	632	56	36	37	47
	水腎・水尿管なし	49.7%	42.9%	33.3%	48.6%	51.1%
	水腎・水尿管あり	44.9%	55.4%	63.9%	51.4%	46.8%
腎結石及び尿管結石	(人)					

【重症度別集計】表

(単位: %)		平均	退院年月							
			2013/04	2013/05	2013/06	2013/07	2013/08	2013/09	2013/10	2013/11
腎結石及び尿管結石	不明	5.4%	14.5%	5.3%	14.3%	12.5%	1.4%	0.0%	3.9%	1.9%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
乳房の悪性新生物	(人)	576	50	52	63	68	40	43	48	51
	ステージⅠ	4.9%	0.0%	1.9%	1.6%	1.5%	5.0%	7.0%	8.3%	3.9%
	ステージⅡ	33.7%	34.0%	23.1%	39.7%	32.4%	47.5%	27.9%	29.2%	33.3%
	ステージⅢ	32.1%	36.0%	38.5%	25.4%	33.8%	27.5%	41.9%	33.3%	21.6%
	ステージⅣ	10.4%	10.0%	7.7%	7.9%	13.2%	7.5%	9.3%	10.4%	11.8%
	不明	10.1%	8.0%	15.4%	12.7%	10.3%	7.5%	7.0%	6.3%	17.6%
	計	8.9%	12.0%	13.5%	12.7%	8.8%	5.0%	7.0%	12.5%	11.8%
膝関節症	(人)	386	37	46	37	44	35	33	30	29
	骨移植なし	90.7%	75.7%	76.1%	75.7%	88.6%	94.3%	100.0%	100.0%	100.0%
	骨移植あり	1.0%	2.7%	0.0%	2.7%	2.3%	2.9%	0.0%	0.0%	0.0%
	不明	8.3%	21.6%	23.9%	21.6%	9.1%	2.9%	0.0%	0.0%	0.0%
鼠径ヘルニア	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	912	81	78	85	92	104	57	69	63
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
全疾患合計		100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
集計病院数			-	17	17	17	16	16	16	16

【重症度別集計】表

(単位:%)		平均	退院年月			
			2013/12	2014/01	2014/02	2014/03
腎結石及び尿管結石	不明	5.4%	1.8%	2.8%	0.0%	2.1%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
乳房の悪性新生物	(人)	576	43	35	40	43
	ステージⅠ	4.9%	4.7%	11.4%	7.5%	11.6%
	ステージⅡ	33.7%	44.2%	31.4%	22.5%	39.5%
	ステージⅢ	32.1%	20.9%	25.7%	52.5%	30.2%
	ステージⅣ	10.4%	11.6%	17.1%	12.5%	7.0%
	不明	10.1%	14.0%	8.6%	2.5%	7.0%
	計	8.9%	4.7%	5.7%	2.5%	4.7%
膝関節症	(人)	386	23	22	28	22
	骨移植なし	90.7%	100.0%	100.0%	100.0%	100.0%
	骨移植あり	1.0%	0.0%	0.0%	0.0%	0.0%
	不明	8.3%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
鼠径ヘルニア	(人)	912	66	57	78	82
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
全疾患合計		100.0%	0.0%	0.0%	0.0%	0.0%
集計病院数	-	16	16	16	16	16