

# 【重症度別集計】表

## 絞込条件

対象年月設定: 201404 ~ 201503  
 病院の経営主体選択: 全て  
 【詳細】病床数: 下限なし ~ 上限なし  
 【詳細】手術の有無: 全て  
 【詳細】ステージ集約選択: 有り  
 【詳細】年代: 全て  
 【詳細】性別: 全て  
 【詳細】入院経路: 全て  
 並べ替え条件: 退院年月・昇順

## ■ 退院患者数

	(単位:%)	平均	退院年月							
			2014/04	2014/05	2014/06	2014/07	2014/08	2014/09	2014/10	2014/11
<b>胃の悪性新生物</b>	(人)	696	72	63	46	58	70	62	60	54
ステージ0		2.3%	4.2%	0.0%	2.2%	1.7%	0.0%	3.2%	8.3%	0.0%
ステージI		28.3%	27.8%	31.7%	23.9%	15.5%	28.6%	29.0%	31.7%	24.1%
ステージII		14.1%	13.9%	11.1%	13.0%	19.0%	17.1%	17.7%	11.7%	14.8%
ステージIII		11.6%	6.9%	11.1%	2.2%	8.6%	8.6%	11.3%	8.3%	16.7%
ステージIV		37.2%	37.5%	39.7%	47.8%	48.3%	41.4%	37.1%	35.0%	29.6%
不明		6.5%	9.7%	6.3%	10.9%	6.9%	4.3%	1.6%	5.0%	14.8%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
<b>結腸の悪性新生物</b>	(人)	866	83	75	83	76	80	70	70	70
ステージ0		3.7%	2.4%	5.3%	2.4%	1.3%	1.3%	7.1%	1.4%	5.7%
ステージI		8.5%	7.2%	12.0%	4.8%	3.9%	7.5%	5.7%	8.6%	10.0%
ステージII		22.7%	16.9%	25.3%	16.9%	28.9%	21.3%	20.0%	27.1%	21.4%
ステージIII		22.3%	30.1%	16.0%	21.7%	27.6%	16.3%	30.0%	12.9%	20.0%
ステージIV		34.6%	34.9%	32.0%	44.6%	31.6%	47.5%	28.6%	34.3%	34.3%
不明		8.1%	8.4%	9.3%	9.6%	6.6%	6.3%	8.6%	15.7%	8.6%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
<b>直腸の悪性新生物</b>	(人)	453	34	29	29	32	49	36	39	44
ステージ0		3.1%	2.9%	3.4%	3.4%	6.3%	2.0%	0.0%	5.1%	0.0%
ステージI		10.6%	17.6%	13.8%	6.9%	6.3%	8.2%	8.3%	2.6%	9.1%
ステージII		16.6%	26.5%	13.8%	6.9%	6.3%	18.4%	25.0%	7.7%	15.9%
ステージIII		32.7%	17.6%	37.9%	27.6%	34.4%	30.6%	36.1%	38.5%	34.1%
ステージIV		33.8%	29.4%	24.1%	48.3%	40.6%	38.8%	30.6%	38.5%	38.6%
不明		3.3%	5.9%	6.9%	6.9%	6.3%	2.0%	0.0%	7.7%	2.3%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
<b>気管支および肺の悪性新生物</b>	(人)	783	80	88	70	86	78	89	51	56
ステージ0		0.4%	1.3%	1.1%	1.4%	0.0%	0.0%	0.0%	0.0%	0.0%
ステージI		15.1%	21.3%	11.4%	11.4%	15.1%	17.9%	13.5%	25.5%	14.3%
ステージII		8.2%	16.3%	8.0%	10.0%	12.8%	10.3%	7.9%	3.9%	8.9%
ステージIII		21.7%	21.3%	15.9%	21.4%	23.3%	20.5%	24.7%	23.5%	21.4%
ステージIV		38.2%	33.8%	43.2%	41.4%	36.0%	37.2%	41.6%	25.5%	33.9%
不明		16.5%	6.3%	20.5%	14.3%	12.8%	14.1%	12.4%	21.6%	21.4%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
<b>急性心筋梗塞</b>	(人)	266	14	19	20	20	28	20	29	25
Killip I		41.7%	50.0%	52.6%	35.0%	25.0%	50.0%	55.0%	44.8%	28.0%
Killip II		27.8%	35.7%	21.1%	25.0%	25.0%	25.0%	15.0%	34.5%	40.0%
Killip III		11.7%	0.0%	5.3%	25.0%	25.0%	14.3%	15.0%	3.4%	16.0%
Killip IV		16.9%	14.3%	21.1%	10.0%	20.0%	10.7%	15.0%	17.2%	8.0%
不明		1.9%	0.0%	0.0%	5.0%	5.0%	0.0%	0.0%	0.0%	8.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
<b>肺炎</b>	(人)	2,889	270	283	272	241	229	183	190	175
軽症		23.6%	23.7%	23.0%	18.4%	25.3%	20.1%	20.2%	31.6%	28.0%
中等症		7.9%	5.6%	5.7%	7.0%	7.1%	7.9%	10.9%	11.1%	7.4%
重症		32.9%	36.7%	33.9%	32.7%	26.6%	38.4%	32.8%	25.3%	30.9%
不明		35.5%	34.1%	37.5%	41.9%	41.1%	33.6%	36.1%	32.1%	33.7%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
<b>喘息</b>	(人)	916	74	95	60	76	71	86	82	85
軽症間欠型		30.2%	36.5%	25.3%	21.7%	27.6%	36.6%	33.7%	31.7%	35.3%
軽症持続型		36.2%	35.1%	45.3%	36.7%	36.8%	39.4%	32.6%	34.1%	30.6%
中等症持続型		20.1%	18.9%	16.8%	28.3%	26.3%	16.9%	18.6%	17.1%	21.2%
重症持続型		12.2%	9.5%	11.6%	13.3%	7.9%	7.0%	7.0%	15.9%	12.9%
不明		1.2%	0.0%	1.1%	0.0%	1.3%	0.0%	8.1%	1.2%	0.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
<b>脳梗塞</b>	(人)	1,365	119	123	108	120	118	119	121	96
JCS 0-1		67.4%	63.9%	61.8%	60.2%	75.0%	63.6%	73.1%	68.6%	61.5%
JCS 2-3		20.8%	26.1%	27.6%	23.1%	16.7%	23.7%	13.4%	19.0%	27.1%
JCS 10-100		9.2%	5.9%	8.9%	13.0%	5.0%	9.3%	11.8%	10.7%	11.5%

# 【重症度別集計】表

	(単位:%)	平均	退院年月			
			2014/12	2015/01	2015/02	2015/03
<b>胃の悪性新生物</b>	(人)	696	62	54	45	50
ステージ0		2.3%	3.2%	3.7%	0.0%	0.0%
ステージⅠ		28.3%	30.6%	31.5%	26.7%	38.0%
ステージⅡ		14.1%	11.3%	13.0%	11.1%	14.0%
ステージⅢ		11.6%	19.4%	20.4%	20.0%	8.0%
ステージⅣ		37.2%	35.5%	31.5%	28.9%	32.0%
不明		6.5%	0.0%	0.0%	13.3%	8.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%
<b>結腸の悪性新生物</b>	(人)	866	67	56	81	55
ステージ0		3.7%	3.0%	7.1%	6.2%	1.8%
ステージⅠ		8.5%	14.9%	10.7%	7.4%	12.7%
ステージⅡ		22.7%	22.4%	30.4%	21.0%	25.5%
ステージⅢ		22.3%	22.4%	19.6%	30.9%	16.4%
ステージⅣ		34.6%	31.3%	26.8%	27.2%	40.0%
不明		8.1%	6.0%	5.4%	7.4%	3.6%
計		100.0%	100.0%	100.0%	100.0%	100.0%
<b>直腸の悪性新生物</b>	(人)	453	44	46	32	39
ステージ0		3.1%	2.3%	4.3%	3.1%	5.1%
ステージⅠ		10.6%	18.2%	13.0%	9.4%	12.8%
ステージⅡ		16.6%	13.6%	17.4%	9.4%	33.3%
ステージⅢ		32.7%	38.6%	30.4%	37.5%	28.2%
ステージⅣ		33.8%	27.3%	34.8%	40.6%	15.4%
不明		3.3%	0.0%	0.0%	0.0%	5.1%
計		100.0%	100.0%	100.0%	100.0%	100.0%
<b>気管支および肺の悪性新生物</b>	(人)	783	50	43	45	47
ステージ0		0.4%	0.0%	0.0%	0.0%	0.0%
ステージⅠ		15.1%	16.0%	7.0%	13.3%	12.8%
ステージⅡ		8.2%	2.0%	0.0%	4.4%	2.1%
ステージⅢ		21.7%	22.0%	16.3%	20.0%	31.9%
ステージⅣ		38.2%	36.0%	51.2%	40.0%	38.3%
不明		16.5%	24.0%	25.6%	22.2%	14.9%
計		100.0%	100.0%	100.0%	100.0%	100.0%
<b>急性心筋梗塞</b>	(人)	266	34	21	22	14
KillipⅠ		41.7%	38.2%	38.1%	45.5%	42.9%
KillipⅡ		27.8%	23.5%	42.9%	27.3%	14.3%
KillipⅢ		11.7%	8.8%	9.5%	9.1%	7.1%
KillipⅣ		16.9%	26.5%	9.5%	18.2%	35.7%
不明		1.9%	2.9%	0.0%	0.0%	0.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%
<b>肺炎</b>	(人)	2,889	267	333	225	221
軽症		23.6%	25.8%	21.9%	26.2%	22.6%
中等症		7.9%	11.2%	7.5%	7.1%	8.6%
重症		32.9%	30.7%	38.1%	32.9%	31.7%
不明		35.5%	32.2%	32.4%	33.8%	37.1%
計		100.0%	100.0%	100.0%	100.0%	100.0%
<b>喘息</b>	(人)	916	66	63	60	98
軽症間欠型		30.2%	33.3%	25.4%	38.3%	20.4%
軽症持続型		36.2%	31.8%	31.7%	30.0%	44.9%
中等症持続型		20.1%	19.7%	22.2%	18.3%	19.4%
重症持続型		12.2%	15.2%	20.6%	13.3%	14.3%
不明		1.2%	0.0%	0.0%	0.0%	1.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%
<b>脳梗塞</b>	(人)	1,365	133	91	92	125
JCS 0-1		67.4%	75.9%	75.8%	66.3%	62.4%
JCS 2-3		20.8%	15.0%	14.3%	21.7%	22.4%
JCS 10-100		9.2%	6.0%	8.8%	8.7%	11.2%

【重症度別集計】表

(単位:%)		平均	退院年月							
			2014/04	2014/05	2014/06	2014/07	2014/08	2014/09	2014/10	2014/11
脳梗塞	JCS 200-300	2.6%	4.2%	1.6%	3.7%	3.3%	3.4%	1.7%	1.7%	0.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
脳出血	(人)	586	58	58	42	40	40	33	39	47
	JCS 0-1	46.1%	43.1%	53.4%	52.4%	57.5%	55.0%	48.5%	56.4%	34.0%
	JCS 2-3	19.3%	15.5%	15.5%	11.9%	15.0%	17.5%	9.1%	12.8%	25.5%
	JCS 10-100	19.1%	17.2%	19.0%	21.4%	17.5%	15.0%	24.2%	20.5%	23.4%
	JCS 200-300	15.5%	24.1%	12.1%	14.3%	10.0%	12.5%	18.2%	10.3%	17.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
糖尿病	(人)	585	55	51	43	54	66	42	44	42
	軽症	65.3%	56.4%	72.5%	62.8%	64.8%	63.6%	81.0%	70.5%	71.4%
	中等症	17.9%	20.0%	17.6%	16.3%	11.1%	19.7%	11.9%	18.2%	7.1%
	重症	11.1%	9.1%	5.9%	7.0%	14.8%	9.1%	4.8%	6.8%	16.7%
	不明	5.6%	14.5%	3.9%	14.0%	9.3%	7.6%	2.4%	4.5%	4.8%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
大腿骨頸部骨折	(人)	511	38	41	46	37	47	40	45	26
	重症度Ⅰ	41.5%	28.9%	31.7%	34.8%	43.2%	34.0%	50.0%	35.6%	46.2%
	重症度Ⅱ	19.8%	28.9%	24.4%	19.6%	24.3%	17.0%	17.5%	26.7%	15.4%
	重症度Ⅲ	27.4%	34.2%	29.3%	23.9%	21.6%	31.9%	12.5%	24.4%	34.6%
	重症度Ⅳ	5.9%	0.0%	4.9%	13.0%	5.4%	4.3%	17.5%	6.7%	3.8%
	不明	5.5%	7.9%	9.8%	8.7%	5.4%	12.8%	2.5%	6.7%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
胃潰瘍	(人)	202	18	21	13	19	11	14	20	18
	ショックなし(収縮期血圧90mmHg以上)	87.6%	88.9%	85.7%	92.3%	84.2%	90.9%	85.7%	95.0%	77.8%
	ショックあり(収縮期血圧90mmHg未満)	11.9%	11.1%	14.3%	7.7%	15.8%	9.1%	14.3%	5.0%	22.2%
	不明	0.5%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
急性腸炎	(人)	701	66	76	59	57	56	52	49	64
	血便なし	92.3%	92.4%	89.5%	94.9%	93.0%	89.3%	92.3%	81.6%	93.8%
	血便あり	6.6%	6.1%	10.5%	1.7%	5.3%	3.6%	7.7%	18.4%	6.3%
	不明	1.1%	1.5%	0.0%	3.4%	1.8%	7.1%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
急性虫垂炎	(人)	522	46	43	40	54	49	49	58	35
	白血球15000/mm3未満	67.4%	67.4%	60.5%	70.0%	64.8%	61.2%	75.5%	74.1%	68.6%
	白血球15000/mm3以上	32.2%	32.6%	37.2%	30.0%	35.2%	36.7%	24.5%	25.9%	31.4%
	不明	0.4%	0.0%	2.3%	0.0%	0.0%	2.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
胆石症	(人)	1,108	100	94	75	88	87	106	102	104
	黄疸なし(T Bill 5mg/dl 未満)	93.1%	93.0%	94.7%	94.7%	87.5%	97.7%	97.2%	86.3%	95.2%
	黄疸あり(T Bill 5mg/dl 以上)	6.9%	7.0%	5.3%	5.3%	12.5%	2.3%	2.8%	13.7%	4.8%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
前立腺肥大症	(人)	95	6	4	4	11	9	7	10	17
	前立腺容積20ml未満	20.0%	16.7%	25.0%	50.0%	0.0%	22.2%	28.6%	0.0%	5.9%
	前立腺容積50ml未満	40.0%	33.3%	50.0%	25.0%	81.8%	44.4%	0.0%	50.0%	35.3%
	前立腺容積50ml以上	30.5%	16.7%	25.0%	0.0%	18.2%	22.2%	42.9%	40.0%	52.9%
	不明	9.5%	33.3%	0.0%	25.0%	0.0%	11.1%	28.6%	10.0%	5.9%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
白内障	(人)	679	53	60	68	69	61	45	83	52
	1眼(処置対象のもの)	91.2%	96.2%	100.0%	100.0%	100.0%	100.0%	100.0%	95.2%	100.0%
	2眼(処置対象のもの)	8.8%	3.8%	0.0%	0.0%	0.0%	0.0%	0.0%	4.8%	0.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
痔核	(人)	74	6	12	4	10	3	6	6	7
	第一度(出血のみ)	24.3%	16.7%	8.3%	0.0%	10.0%	33.3%	33.3%	16.7%	28.6%
	第二度(還納可能)	12.2%	0.0%	16.7%	0.0%	40.0%	0.0%	0.0%	0.0%	14.3%
	第三度(脱肛)	62.2%	83.3%	75.0%	100.0%	50.0%	33.3%	66.7%	83.3%	57.1%
	不明	1.4%	0.0%	0.0%	0.0%	0.0%	33.3%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
子宮筋腫	(人)	142	14	18	7	15	13	11	14	8
	貧血なし(Hb10g/dl以上)	65.5%	71.4%	50.0%	42.9%	73.3%	69.2%	63.6%	64.3%	75.0%
	貧血あり(Hb10g/dl未満)	33.8%	28.6%	50.0%	57.1%	26.7%	30.8%	36.4%	28.6%	25.0%
	不明	0.7%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	7.1%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
狭心症	(人)	1,963	135	112	125	176	182	196	216	158
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
腎結石及び尿管結石	(人)	487	26	34	43	33	58	46	47	43
	水腎・尿管なし	35.7%	65.4%	29.4%	34.9%	33.3%	29.3%	28.3%	40.4%	27.9%
	水腎・尿管あり	57.5%	30.8%	47.1%	58.1%	60.6%	65.5%	65.2%	44.7%	62.8%

# 【重症度別集計】表

		平均	退院年月			
			2014/12	2015/01	2015/02	2015/03
		(単位:%)				
脳梗塞	JCS 200-300	2.6%	3.0%	1.1%	3.3%	4.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
脳出血	(人)	586	62	68	51	48
	JCS 0-1	46.1%	43.5%	39.7%	39.2%	39.6%
	JCS 2-3	19.3%	12.9%	33.8%	25.5%	27.1%
	JCS 10-100	19.1%	17.7%	14.7%	23.5%	18.8%
	JCS 200-300	15.5%	25.8%	11.8%	11.8%	14.6%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
糖尿病	(人)	585	54	40	55	39
	軽症	65.3%	68.5%	52.5%	58.2%	64.1%
	中等症	17.9%	20.4%	20.0%	29.1%	20.5%
	重症	11.1%	9.3%	27.5%	12.7%	12.8%
	不明	5.6%	1.9%	0.0%	0.0%	2.6%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
大腿骨頸部骨折	(人)	511	56	44	41	50
	重症度Ⅰ	41.5%	42.9%	54.5%	39.0%	56.0%
	重症度Ⅱ	19.8%	14.3%	15.9%	14.6%	20.0%
	重症度Ⅲ	27.4%	33.9%	25.0%	39.0%	20.0%
	重症度Ⅳ	5.9%	3.6%	2.3%	4.9%	4.0%
	不明	5.5%	5.4%	2.3%	2.4%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
胃潰瘍	(人)	202	19	18	16	15
	ショックなし(収縮期血圧90mmHg以上)	87.6%	84.2%	100.0%	93.8%	73.3%
	ショックあり(収縮期血圧90mmHg未満)	11.9%	10.5%	0.0%	6.3%	26.7%
	不明	0.5%	5.3%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
急性腸炎	(人)	701	72	53	48	49
	血便なし	92.3%	98.6%	96.2%	93.8%	89.8%
	血便あり	6.6%	1.4%	3.8%	6.3%	10.2%
	不明	1.1%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
急性虫垂炎	(人)	522	41	33	31	43
	白血球15000/mm3未満	67.4%	70.7%	66.7%	67.7%	60.5%
	白血球15000/mm3以上	32.2%	29.3%	33.3%	32.3%	39.5%
	不明	0.4%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
胆石症	(人)	1,108	107	74	83	88
	黄疸なし(T Bill 5mg/dl 未満)	93.1%	92.5%	93.2%	91.6%	93.2%
	黄疸あり(T Bill 5mg/dl 以上)	6.9%	7.5%	6.8%	8.4%	6.8%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
前立腺肥大症	(人)	95	10	7	6	4
	前立腺容積20ml未満	20.0%	40.0%	57.1%	33.3%	0.0%
	前立腺容積50ml未満	40.0%	40.0%	28.6%	16.7%	50.0%
	前立腺容積50ml以上	30.5%	10.0%	14.3%	50.0%	50.0%
	不明	9.5%	10.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
白内障	(人)	679	44	40	48	56
	1眼(処置対象のもの)	91.2%	95.5%	65.0%	60.4%	66.1%
	2眼(処置対象のもの)	8.8%	4.5%	35.0%	39.6%	33.9%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
痔核	(人)	74	7	2	5	6
	第一度(出血のみ)	24.3%	42.9%	50.0%	20.0%	66.7%
	第二度(還納可能)	12.2%	14.3%	50.0%	0.0%	0.0%
	第三度(脱肛)	62.2%	42.9%	0.0%	80.0%	33.3%
	不明	1.4%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
子宮筋腫	(人)	142	10	8	13	11
	貧血なし(Hb10g/dl以上)	65.5%	100.0%	75.0%	61.5%	45.5%
	貧血あり(Hb10g/dl未満)	33.8%	0.0%	25.0%	38.5%	54.5%
	不明	0.7%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
狭心症	(人)	1,963	171	158	167	167
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
腎結石及び尿管結石	(人)	487	47	38	36	36
	水腎・水尿管なし	35.7%	38.3%	50.0%	25.0%	38.9%
	水腎・水尿管あり	57.5%	59.6%	50.0%	72.2%	61.1%



# 【重症度別集計】表

		平均	退院年月			
			2014/12	2015/01	2015/02	2015/03
腎結石及び尿管結石	不明	6.8%	2.1%	0.0%	2.8%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
乳房の悪性新生物	(人)	493	51	37	50	41
	ステージ0	6.5%	5.9%	5.4%	6.0%	4.9%
	ステージI	34.7%	27.5%	27.0%	34.0%	48.8%
	ステージII	36.1%	45.1%	37.8%	30.0%	24.4%
	ステージIII	8.9%	5.9%	16.2%	10.0%	4.9%
	ステージIV	8.3%	13.7%	2.7%	12.0%	7.3%
	不明	5.5%	2.0%	10.8%	8.0%	9.8%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
膝関節症	(人)	313	24	14	18	24
	骨移植なし	100.0%	100.0%	100.0%	100.0%	100.0%
	骨移植あり	N/A	0.0%	0.0%	0.0%	0.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
鼠径ヘルニア	(人)	727	63	47	60	58
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
全疾患合計		100.0%	0.0%	0.0%	0.0%	0.0%
集計病院数	-		13	12	12	11