

【重症度別集計】表

絞込条件

対象年月設定: 201504 ~ 201603
 病院の経営主体選択: 全て
 【詳細】病床数: 下限なし ~ 上限なし
 【詳細】手術の有無: 全て
 【詳細】ステージ集約選択: 有り
 【詳細】年代: 全て
 【詳細】性別: 全て
 【詳細】入院経路: 全て
 並べ替え条件: 退院年月・昇順

■ 退院患者数

(単位: %)	平均	退院年月								
		2015/04	2015/05	2015/06	2015/07	2015/08	2015/09	2015/10	2015/11	
胃の悪性新生物	(人)	586	44	53	49	46	55	52	31	46
	ステージ0	1.5%	2.3%	3.8%	4.1%	2.2%	0.0%	0.0%	0.0%	0.0%
	ステージI	27.3%	22.7%	20.8%	18.4%	30.4%	21.8%	32.7%	29.0%	34.8%
	ステージII	12.3%	13.6%	22.6%	16.3%	19.6%	18.2%	7.7%	6.5%	2.2%
	ステージIII	17.2%	13.6%	18.9%	10.2%	23.9%	21.8%	19.2%	22.6%	10.9%
	ステージIV	33.4%	29.5%	28.3%	42.9%	19.6%	32.7%	36.5%	32.3%	41.3%
	不明	8.2%	18.2%	5.7%	8.2%	4.3%	5.5%	3.8%	9.7%	10.9%
結腸の悪性新生物	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	957	74	62	74	86	77	79	88	79
	ステージ0	2.8%	5.4%	3.2%	2.7%	3.5%	1.3%	1.3%	2.3%	2.5%
	ステージI	7.7%	13.5%	3.2%	5.4%	11.6%	5.2%	2.5%	9.1%	6.3%
	ステージII	23.2%	14.9%	29.0%	24.3%	23.3%	31.2%	26.6%	22.7%	22.8%
	ステージIII	32.2%	27.0%	27.4%	32.4%	26.7%	27.3%	32.9%	31.8%	39.2%
	ステージIV	26.4%	27.0%	25.8%	25.7%	30.2%	24.7%	27.8%	26.1%	22.8%
直腸の悪性新生物	不明	7.6%	12.2%	11.3%	9.5%	4.7%	10.4%	8.9%	8.0%	6.3%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	515	42	30	35	46	41	43	55	44
	ステージ0	2.3%	0.0%	0.0%	0.0%	4.3%	7.3%	2.3%	1.8%	2.3%
	ステージI	9.9%	11.9%	0.0%	17.1%	4.3%	9.8%	7.0%	12.7%	13.6%
	ステージII	18.1%	23.8%	20.0%	14.3%	21.7%	14.6%	25.6%	21.8%	20.5%
	ステージIII	27.0%	21.4%	23.3%	34.3%	34.8%	17.1%	20.9%	29.1%	29.5%
気管支および肺の悪性新生物	ステージIV	38.1%	35.7%	50.0%	28.6%	32.6%	46.3%	34.9%	30.9%	31.8%
	不明	4.7%	7.1%	6.7%	5.7%	2.2%	4.9%	9.3%	3.6%	2.3%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	522	49	39	39	56	32	40	40	44
	ステージ0	0.6%	0.0%	0.0%	0.0%	0.0%	0.0%	2.5%	0.0%	0.0%
	ステージI	17.6%	18.4%	15.4%	10.3%	10.7%	21.9%	17.5%	17.5%	27.3%
	ステージII	5.7%	2.0%	5.1%	0.0%	7.1%	0.0%	2.5%	12.5%	13.6%
急性心筋梗塞	ステージIII	16.7%	30.6%	20.5%	20.5%	16.1%	15.6%	10.0%	15.0%	6.8%
	ステージIV	35.2%	26.5%	38.5%	25.6%	41.1%	34.4%	40.0%	32.5%	20.5%
	不明	24.1%	22.4%	20.5%	43.6%	25.0%	28.1%	27.5%	22.5%	31.8%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	316	31	24	31	14	31	28	28	24
	Killip I	42.4%	32.3%	54.2%	51.6%	50.0%	48.4%	39.3%	53.6%	16.7%
	Killip II	32.9%	45.2%	29.2%	35.5%	35.7%	22.6%	32.1%	25.0%	41.7%
肺炎	Killip III	7.6%	6.5%	4.2%	0.0%	7.1%	6.5%	14.3%	7.1%	4.2%
	Killip IV	11.4%	12.9%	12.5%	9.7%	0.0%	12.9%	7.1%	14.3%	20.8%
	不明	5.7%	3.2%	0.0%	3.2%	7.1%	9.7%	7.1%	0.0%	16.7%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	1,893	139	135	158	143	147	142	173	155
	軽症	32.8%	33.1%	29.6%	29.1%	32.2%	32.0%	40.1%	31.2%	26.5%
	中等症	9.1%	10.1%	13.3%	7.6%	2.8%	6.1%	6.3%	5.8%	7.1%
喘息	重症	38.5%	43.2%	37.8%	41.8%	43.4%	38.1%	35.2%	37.6%	38.7%
	不明	19.7%	13.7%	19.3%	21.5%	21.7%	23.8%	18.3%	25.4%	27.7%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	367	18	36	24	28	23	60	44	28
	軽症間欠型	31.6%	27.8%	27.8%	29.2%	25.0%	39.1%	41.7%	29.5%	32.1%
	軽症持続型	16.1%	5.6%	22.2%	16.7%	25.0%	13.0%	13.3%	9.1%	14.3%
	中等症持続型	37.6%	50.0%	41.7%	54.2%	28.6%	34.8%	31.7%	34.1%	32.1%
脳梗塞	重症持続型	13.9%	16.7%	5.6%	0.0%	17.9%	13.0%	13.3%	27.3%	21.4%
	不明	0.8%	0.0%	2.8%	0.0%	3.6%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	1,234	91	117	121	101	114	75	116	99
	JCS 0-1	60.6%	60.4%	61.5%	52.1%	55.4%	57.9%	61.3%	63.8%	62.6%
	JCS 2-3	29.7%	30.8%	33.3%	35.5%	28.7%	36.0%	30.7%	30.2%	25.3%
	JCS 10-100	7.7%	7.7%	4.3%	10.7%	11.9%	6.1%	5.3%	5.2%	10.1%

【重症度別集計】表

	(単位:%)	平均	退院年月			
			2015/12	2016/01	2016/02	2016/03
胃の悪性新生物	(人)	586	53	48	46	63
	ステージ0	1.5%	3.8%	0.0%	2.2%	0.0%
	ステージI	27.3%	20.8%	22.9%	32.6%	39.7%
	ステージII	12.3%	5.7%	12.5%	10.9%	9.5%
	ステージIII	17.2%	17.0%	18.8%	19.6%	12.7%
	ステージIV	33.4%	45.3%	33.3%	30.4%	28.6%
	不明	8.2%	7.5%	12.5%	4.3%	9.5%
計		100.0%	100.0%	100.0%	100.0%	100.0%
結腸の悪性新生物	(人)	957	84	77	79	98
	ステージ0	2.8%	3.6%	2.6%	3.8%	2.0%
	ステージI	7.7%	3.6%	9.1%	11.4%	10.2%
	ステージII	23.2%	21.4%	24.7%	16.5%	22.4%
	ステージIII	32.2%	34.5%	33.8%	38.0%	33.7%
	ステージIV	26.4%	27.4%	23.4%	27.8%	27.6%
	不明	7.6%	9.5%	6.5%	2.5%	4.1%
計		100.0%	100.0%	100.0%	100.0%	100.0%
直腸の悪性新生物	(人)	515	44	55	37	43
	ステージ0	2.3%	0.0%	7.3%	0.0%	0.0%
	ステージI	9.9%	9.1%	12.7%	16.2%	2.3%
	ステージII	18.1%	15.9%	12.7%	18.9%	7.0%
	ステージIII	27.0%	25.0%	29.1%	27.0%	30.2%
	ステージIV	38.1%	47.7%	32.7%	35.1%	55.8%
	不明	4.7%	2.3%	5.5%	2.7%	4.7%
計		100.0%	100.0%	100.0%	100.0%	100.0%
気管支および肺の悪性新生物	(人)	522	43	46	41	53
	ステージ0	0.6%	2.3%	0.0%	0.0%	1.9%
	ステージI	17.6%	23.3%	19.6%	17.1%	15.1%
	ステージII	5.7%	4.7%	4.3%	7.3%	7.5%
	ステージIII	16.7%	14.0%	13.0%	22.0%	15.1%
	ステージIV	35.2%	41.9%	43.5%	29.3%	45.3%
	不明	24.1%	14.0%	19.6%	24.4%	15.1%
計		100.0%	100.0%	100.0%	100.0%	100.0%
急性心筋梗塞	(人)	316	34	24	18	29
	Killip I	42.4%	38.2%	33.3%	27.8%	58.6%
	Killip II	32.9%	29.4%	37.5%	38.9%	27.6%
	Killip III	7.6%	17.6%	8.3%	11.1%	3.4%
	Killip IV	11.4%	8.8%	16.7%	11.1%	6.9%
	不明	5.7%	5.9%	4.2%	11.1%	3.4%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
肺炎	(人)	1,893	182	169	185	165
	軽症	32.8%	34.6%	34.3%	31.9%	38.2%
	中等症	9.1%	10.4%	14.8%	13.0%	10.3%
	重症	38.5%	33.0%	35.5%	36.2%	43.6%
	不明	19.7%	22.0%	15.4%	18.9%	7.9%
計		100.0%	100.0%	100.0%	100.0%	100.0%
喘息	(人)	367	35	28	17	26
	軽症間欠型	31.6%	34.3%	14.3%	29.4%	38.5%
	軽症持続型	16.1%	17.1%	25.0%	23.5%	11.5%
	中等症持続型	37.6%	31.4%	46.4%	41.2%	42.3%
	重症持続型	13.9%	17.1%	14.3%	5.9%	3.8%
	不明	0.8%	0.0%	0.0%	0.0%	3.8%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
脳梗塞	(人)	1,234	111	91	77	121
	JCS 0-1	60.6%	64.9%	51.6%	64.9%	70.2%
	JCS 2-3	29.7%	25.2%	38.5%	23.4%	18.2%
	JCS 10-100	7.7%	5.4%	5.5%	10.4%	9.9%

【重症度別集計】表

(単位: %)		平均	退院年月							
			2015/04	2015/05	2015/06	2015/07	2015/08	2015/09	2015/10	2015/11
脳梗塞	JCS 200-300	2.0%	1.1%	0.9%	1.7%	4.0%	0.0%	2.7%	0.9%	2.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
脳出血	(人)	525	48	40	48	45	38	25	43	37
	JCS 0-1	38.1%	31.3%	32.5%	43.8%	40.0%	26.3%	52.0%	44.2%	37.8%
	JCS 2-3	24.0%	25.0%	12.5%	25.0%	24.4%	31.6%	28.0%	30.2%	18.9%
	JCS 10-100	21.3%	16.7%	32.5%	22.9%	17.8%	28.9%	12.0%	14.0%	32.4%
	JCS 200-300	16.6%	27.1%	22.5%	8.3%	17.8%	13.2%	8.0%	11.6%	10.8%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
糖尿病	(人)	407	38	39	34	32	45	35	24	30
	軽症	61.4%	55.3%	71.8%	58.8%	50.0%	57.8%	65.7%	70.8%	66.7%
	中等症	22.4%	28.9%	23.1%	23.5%	25.0%	22.2%	11.4%	20.8%	20.0%
	重症	16.0%	15.8%	5.1%	17.6%	25.0%	17.8%	22.9%	8.3%	13.3%
	不明	0.2%	0.0%	0.0%	0.0%	0.0%	2.2%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
大腿骨頸部骨折	(人)	429	32	29	24	40	37	35	39	32
	重症度 I	43.8%	43.8%	48.3%	41.7%	45.0%	56.8%	51.4%	33.3%	43.8%
	重症度 II	22.4%	3.1%	13.8%	12.5%	20.0%	16.2%	8.6%	41.0%	25.0%
	重症度 III	28.7%	43.8%	37.9%	45.8%	25.0%	24.3%	37.1%	15.4%	25.0%
	重症度 IV	4.2%	9.4%	0.0%	0.0%	10.0%	0.0%	2.9%	10.3%	6.3%
	不明	0.9%	0.0%	0.0%	0.0%	0.0%	2.7%	0.0%	0.0%	0.0%
胃潰瘍	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	157	12	13	14	10	15	18	11	9
	ショックなし(収縮期血圧90mmHg以上)	91.7%	100.0%	84.6%	100.0%	80.0%	93.3%	88.9%	90.9%	66.7%
	ショックあり(収縮期血圧90mmHg未満)	8.3%	0.0%	15.4%	0.0%	20.0%	6.7%	11.1%	9.1%	33.3%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
急性腸炎	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	518	42	52	34	48	49	43	40	38
	血便なし	90.3%	90.5%	90.4%	91.2%	75.0%	93.9%	79.1%	92.5%	92.1%
	血便あり	9.3%	7.1%	9.6%	8.8%	22.9%	6.1%	20.9%	7.5%	7.9%
	不明	0.4%	2.4%	0.0%	0.0%	2.1%	0.0%	0.0%	0.0%	0.0%
急性虫垂炎	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	423	34	36	38	36	35	35	51	30
	白血球15000/mm ³ 未満	72.3%	64.7%	75.0%	63.2%	77.8%	74.3%	80.0%	78.4%	66.7%
	白血球15000/mm ³ 以上	27.0%	35.3%	25.0%	36.8%	22.2%	25.7%	20.0%	21.6%	33.3%
	不明	0.7%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
胆石症	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	923	57	83	83	78	76	81	98	71
	黄疸なし(T Bill 5mg/dl 未満)	92.8%	94.7%	92.8%	92.8%	96.2%	97.4%	90.1%	90.8%	94.4%
	黄疸あり(T Bill 5mg/dl 以上)	7.2%	5.3%	7.2%	7.2%	3.8%	2.6%	9.9%	9.2%	5.6%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
前立腺肥大症	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	43	6	3	2	5	3	1	3	2
	前立腺容積20ml未満	7.0%	16.7%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	前立腺容積50ml未満	27.9%	50.0%	33.3%	50.0%	60.0%	33.3%	0.0%	66.7%	0.0%
	前立腺容積50ml以上	46.5%	33.3%	66.7%	50.0%	40.0%	0.0%	0.0%	33.3%	100.0%
白内障	不明	18.6%	0.0%	0.0%	0.0%	0.0%	66.7%	100.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	647	58	48	56	65	43	43	60	63
	1眼(処置対象のもの)	96.6%	69.0%	100.0%	100.0%	96.9%	97.7%	100.0%	100.0%	98.4%
	2眼(処置対象のもの)	3.1%	31.0%	0.0%	0.0%	1.5%	2.3%	0.0%	0.0%	0.0%
痔核	不明	0.3%	0.0%	0.0%	0.0%	1.5%	0.0%	0.0%	0.0%	1.6%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	109	14	8	5	8	6	11	6	12
	第一度(出血のみ)	24.8%	21.4%	25.0%	40.0%	12.5%	0.0%	27.3%	50.0%	25.0%
	第二度(還納可能)	18.3%	7.1%	25.0%	60.0%	25.0%	16.7%	36.4%	16.7%	0.0%
子宮筋腫	第三度(脱肛)	52.3%	71.4%	37.5%	0.0%	62.5%	83.3%	36.4%	16.7%	66.7%
	不明	4.6%	0.0%	12.5%	0.0%	0.0%	0.0%	0.0%	16.7%	8.3%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	90	7	6	5	7	7	8	11	9
	貧血なし(Hb10g/dl以上)	80.0%	71.4%	50.0%	60.0%	85.7%	57.1%	87.5%	81.8%	100.0%
狭心症	貧血あり(Hb10g/dl未満)	20.0%	28.6%	50.0%	40.0%	14.3%	42.9%	12.5%	18.2%	0.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	1,688	161	157	153	144	122	127	126	143
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
腎結石及び尿管結石	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	363	21	34	43	39	44	25	36	22
	水腎・水尿管なし	38.0%	38.1%	32.4%	27.9%	23.1%	45.5%	48.0%	38.9%	36.4%
水腎・水尿管あり	水腎・水尿管あり	60.3%	57.1%	67.6%	69.8%	76.9%	52.			

【重症度別集計】表

(単位:%)		平均	退院年月			
			2015/12	2016/01	2016/02	2016/03
脳梗塞	JCS 200-300	2.0%	4.5%	4.4%	1.3%	1.7%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
脳出血	(人)	525	60	41	48	52
	JCS 0-1	38.1%	38.3%	36.6%	39.6%	38.5%
	JCS 2-3	24.0%	23.3%	19.5%	16.7%	32.7%
	JCS 10-100	21.3%	18.3%	24.4%	22.9%	15.4%
	JCS 200-300	16.6%	20.0%	19.5%	20.8%	13.5%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
糖尿病	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	407	28	30	34	38
	軽症	61.4%	53.6%	56.7%	67.6%	63.2%
	中等症	22.4%	32.1%	30.0%	17.6%	15.8%
	重症	16.0%	14.3%	13.3%	14.7%	21.1%
	不明	0.2%	0.0%	0.0%	0.0%	0.0%
大腿骨頸部骨折	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	429	47	35	38	41
	重症度 I	43.8%	46.8%	22.9%	55.3%	36.6%
	重症度 II	22.4%	36.2%	28.6%	18.4%	31.7%
	重症度 III	28.7%	17.0%	40.0%	23.7%	24.4%
	重症度 IV	4.2%	0.0%	8.6%	2.6%	0.0%
腎潰瘍	不明	0.9%	0.0%	0.0%	0.0%	7.3%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	157	10	15	16	14
	ショックなし (収縮期血圧90mmHg以上)	91.7%	100.0%	93.3%	100.0%	92.9%
	ショックあり (収縮期血圧90mmHg未満)	8.3%	0.0%	6.7%	0.0%	7.1%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
急性腸炎	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	518	50	49	32	41
	血便なし	90.3%	96.0%	98.0%	93.8%	92.7%
	血便あり	9.3%	4.0%	2.0%	6.3%	7.3%
	不明	0.4%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
急性虫垂炎	(人)	423	32	16	35	45
	白血球15000/mm ³ 未満	72.3%	71.9%	68.8%	68.6%	73.3%
	白血球15000/mm ³ 以上	27.0%	28.1%	31.3%	22.9%	26.7%
	不明	0.7%	0.0%	0.0%	8.6%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	923	77	67	71	81
胆石症	黄疸なし(T Bill 5mg/dl 未満)	92.8%	90.9%	88.1%	94.4%	92.6%
	黄疸あり(T Bill 5mg/dl 以上)	7.2%	9.1%	11.9%	5.6%	7.4%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	43	7	0	2	9
	前立腺容積20ml未満	7.0%	14.3%	0.0%	0.0%	11.1%
前立腺肥大症	前立腺容積50ml未満	27.9%	0.0%	0.0%	0.0%	11.1%
	前立腺容積50ml以上	46.5%	42.9%	0.0%	100.0%	55.6%
	不明	18.6%	42.9%	0.0%	0.0%	22.2%
	計	100.0%	100.0%	0.0%	100.0%	100.0%
	(人)	647	40	43	59	69
	1眼(処置対象のもの)	96.6%	100.0%	100.0%	100.0%	100.0%
白内障	2眼(処置対象のもの)	3.1%	0.0%	0.0%	0.0%	0.0%
	不明	0.3%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	109	11	3	9	16
	第一度(出血のみ)	24.8%	18.2%	33.3%	22.2%	31.3%
	第二度(還納可能)	18.3%	9.1%	0.0%	11.1%	25.0%
痔核	第三度(脱肛)	52.3%	72.7%	33.3%	55.6%	43.8%
	不明	4.6%	0.0%	33.3%	11.1%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	90	6	6	5	13
	貧血なし(Hb10g/dl以上)	80.0%	83.3%	100.0%	80.0%	84.6%
	貧血あり(Hb10g/dl未満)	20.0%	16.7%	0.0%	20.0%	15.4%
子宮筋腫	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	1,688	143	120	136	156
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%
狭窄症	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	363	21	18	30	30
	水腎・水尿管なし	38.0%	28.6%	66.7%	53.3%	33.3%
	水腎・水尿管あり	60.3%	66.7%	33.3%	46.7%	60.0%
腎結石及び尿管結石	(人)					

【重症度別集計】表

(単位: %)		平均	退院年月							
			2015/04	2015/05	2015/06	2015/07	2015/08	2015/09	2015/10	2015/11
腎結石及び尿管結石	不明	1.7%	4.8%	0.0%	2.3%	0.0%	2.3%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
乳房の悪性新生物	(人)	469	40	26	46	36	38	42	41	42
	ステージⅠ	4.5%	5.0%	3.8%	2.2%	0.0%	2.6%	0.0%	4.9%	11.9%
	ステージⅡ	29.9%	27.5%	53.8%	32.6%	30.6%	13.2%	21.4%	26.8%	16.7%
	ステージⅢ	36.9%	37.5%	26.9%	37.0%	41.7%	47.4%	40.5%	41.5%	42.9%
	ステージⅣ	11.9%	2.5%	0.0%	13.0%	11.1%	21.1%	21.4%	14.6%	11.9%
	不明	8.3%	12.5%	15.4%	10.9%	13.9%	5.3%	9.5%	4.9%	9.5%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
膝関節症	(人)	311	32	25	32	41	23	30	28	23
	骨移植なし	98.7%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	骨移植あり	1.3%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
鼠径ヘルニア	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	574	44	49	58	40	42	54	58	50
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
全疾患合計		100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
集計病院数			-	11	11	11	11	11	11	11

【重症度別集計】表

(単位:%)		平均	退院年月			
			2015/12	2016/01	2016/02	2016/03
腎結石及び尿管結石	不明	1.7%	4.8%	0.0%	0.0%	6.7%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
乳房の悪性新生物	(人)	469	40	34	47	37
	ステージⅠ	4.5%	5.0%	11.8%	2.1%	5.4%
	ステージⅡ	29.9%	25.0%	44.1%	40.4%	35.1%
	ステージⅢ	36.9%	30.0%	23.5%	29.8%	40.5%
	ステージⅣ	11.9%	15.0%	8.8%	12.8%	5.4%
	不明	8.3%	20.0%	11.8%	4.3%	8.1%
	計	8.5%	5.0%	0.0%	10.6%	5.4%
膝関節症	(人)	311	23	16	20	18
	骨移植なし	98.7%	100.0%	93.8%	85.0%	100.0%
	骨移植あり	1.3%	0.0%	6.3%	15.0%	0.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
鼠径ヘルニア	(人)	574	47	40	43	49
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
全疾患合計		100.0%	0.0%	0.0%	0.0%	0.0%
集計病院数	-	11	11	11	11	11