

【重症度別集計】表

絞込条件

対象年月設定: 201604 ~ 201703
 病院の経営主体選択: 全て
 【詳細】病床数: 下限なし ~ 上限なし
 【詳細】手術の有無: 全て
 【詳細】ステージ集約選択: 有り
 【詳細】年代: 全て
 【詳細】性別: 全て
 【詳細】入院経路: 全て
 並べ替え条件: 退院年月・昇順

■ 退院患者数

	(単位:%)	平均	退院年月							
			2016/04	2016/05	2016/06	2016/07	2016/08	2016/09	2016/10	2016/11
胃の悪性新生物	(人)	578	39	51	54	55	43	52	39	45
ステージ0		3.6%	2.6%	2.0%	0.0%	3.6%	4.7%	1.9%	5.1%	2.2%
ステージI		30.8%	25.6%	23.5%	29.6%	29.1%	18.6%	28.8%	30.8%	31.1%
ステージII		11.6%	5.1%	9.8%	14.8%	14.5%	23.3%	11.5%	10.3%	17.8%
ステージIII		14.5%	20.5%	19.6%	13.0%	20.0%	18.6%	23.1%	15.4%	6.7%
ステージIV		29.2%	38.5%	33.3%	31.5%	23.6%	30.2%	26.9%	25.6%	33.3%
不明		10.2%	7.7%	11.8%	11.1%	9.1%	4.7%	7.7%	12.8%	8.9%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
結腸の悪性新生物	(人)	946	54	77	80	75	61	79	93	83
ステージ0		2.3%	0.0%	3.9%	1.3%	4.0%	3.3%	0.0%	3.2%	3.6%
ステージI		9.7%	11.1%	6.5%	15.0%	4.0%	4.9%	10.1%	6.5%	7.2%
ステージII		16.8%	13.0%	13.0%	18.8%	21.3%	23.0%	19.0%	23.7%	15.7%
ステージIII		33.4%	25.9%	36.4%	35.0%	30.7%	26.2%	36.7%	33.3%	36.1%
ステージIV		31.6%	42.6%	33.8%	26.3%	37.3%	41.0%	27.8%	31.2%	28.9%
不明		6.1%	7.4%	6.5%	3.8%	2.7%	1.6%	6.3%	2.2%	8.4%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
直腸の悪性新生物	(人)	523	29	45	39	47	49	45	52	42
ステージ0		3.8%	0.0%	6.7%	0.0%	2.1%	6.1%	0.0%	3.8%	7.1%
ステージI		9.0%	13.8%	15.6%	10.3%	6.4%	10.2%	8.9%	3.8%	16.7%
ステージII		16.3%	17.2%	20.0%	10.3%	14.9%	22.4%	13.3%	21.2%	9.5%
ステージIII		22.8%	17.2%	22.2%	28.2%	21.3%	12.2%	22.2%	23.1%	23.8%
ステージIV		43.6%	44.8%	31.1%	43.6%	53.2%	49.0%	51.1%	40.4%	40.5%
不明		4.6%	6.9%	4.4%	7.7%	2.1%	0.0%	4.4%	7.7%	2.4%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
気管支および肺の悪性新生物	(人)	518	48	41	38	36	35	43	37	51
ステージ0		N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
ステージI		18.7%	31.3%	14.6%	13.2%	13.9%	14.3%	23.3%	18.9%	13.7%
ステージII		4.1%	6.3%	4.9%	2.6%	2.8%	8.6%	0.0%	2.7%	5.9%
ステージIII		21.6%	27.1%	29.3%	18.4%	5.6%	17.1%	20.9%	21.6%	19.6%
ステージIV		35.7%	29.2%	34.1%	39.5%	47.2%	42.9%	30.2%	40.5%	35.3%
不明		19.9%	6.3%	17.1%	26.3%	30.6%	17.1%	25.6%	16.2%	25.5%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
急性心筋梗塞	(人)	289	17	21	25	21	23	20	34	28
Killip I		53.6%	64.7%	52.4%	56.0%	52.4%	43.5%	65.0%	61.8%	39.3%
Killip II		22.1%	23.5%	28.6%	32.0%	9.5%	26.1%	10.0%	11.8%	17.9%
Killip III		8.0%	5.9%	4.8%	0.0%	14.3%	8.7%	10.0%	2.9%	14.3%
Killip IV		14.5%	0.0%	9.5%	4.0%	23.8%	21.7%	15.0%	23.5%	25.0%
不明		1.7%	5.9%	4.8%	8.0%	0.0%	0.0%	0.0%	0.0%	3.6%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
肺炎	(人)	1,858	120	126	137	124	135	124	158	170
軽症		30.1%	29.2%	25.4%	32.8%	33.9%	28.1%	29.0%	28.5%	26.5%
中等症		9.1%	9.2%	5.6%	11.7%	5.6%	11.9%	4.8%	9.5%	9.4%
重症		44.0%	50.8%	45.2%	38.0%	39.5%	41.5%	42.7%	41.8%	42.4%
不明		16.8%	10.8%	23.8%	17.5%	21.0%	18.5%	23.4%	20.3%	21.8%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
喘息	(人)	274	22	15	15	21	21	26	40	28
軽症間欠型		32.1%	31.8%	40.0%	26.7%	14.3%	19.0%	46.2%	37.5%	21.4%
軽症持続型		24.1%	18.2%	26.7%	33.3%	19.0%	28.6%	15.4%	27.5%	21.4%
中等症持続型		31.0%	45.5%	33.3%	20.0%	47.6%	33.3%	26.9%	25.0%	35.7%
重症持続型		12.4%	4.5%	0.0%	20.0%	19.0%	19.0%	11.5%	10.0%	21.4%
不明		0.4%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
脳梗塞	(人)	1,184	60	96	87	78	104	81	117	104
JCS 0-1		62.3%	63.3%	54.2%	60.9%	57.7%	64.4%	71.6%	61.5%	57.7%
JCS 2-3		27.4%	25.0%	32.3%	32.2%	35.9%	30.8%	19.8%	23.1%	30.8%
JCS 10-100		8.0%	10.0%	12.5%	3.4%	5.1%	4.8%	6.2%	12.8%	9.6%

【重症度別集計】表

	(単位:%)	平均	退院年月			
			2016/12	2017/01	2017/02	2017/03
胃の悪性新生物	(人)	578	59	43	44	54
ステージ0		3.6%	3.4%	2.3%	13.6%	3.7%
ステージI		30.8%	42.4%	39.5%	34.1%	33.3%
ステージII		11.6%	5.1%	9.3%	6.8%	11.1%
ステージIII		14.5%	8.5%	4.7%	11.4%	13.0%
ステージIV		29.2%	30.5%	25.6%	20.5%	31.5%
不明		10.2%	10.2%	18.6%	13.6%	7.4%
計		100.0%	100.0%	100.0%	100.0%	100.0%
結腸の悪性新生物	(人)	946	96	74	79	95
ステージ0		2.3%	3.1%	1.4%	1.3%	2.1%
ステージI		9.7%	12.5%	9.5%	16.5%	11.6%
ステージII		16.8%	12.5%	14.9%	13.9%	13.7%
ステージIII		33.4%	32.3%	35.1%	27.8%	40.0%
ステージIV		31.6%	32.3%	31.1%	30.4%	24.2%
不明		6.1%	7.3%	8.1%	10.1%	8.4%
計		100.0%	100.0%	100.0%	100.0%	100.0%
直腸の悪性新生物	(人)	523	53	51	35	36
ステージ0		3.8%	3.8%	3.9%	2.9%	8.3%
ステージI		9.0%	1.9%	9.8%	5.7%	8.3%
ステージII		16.3%	17.0%	15.7%	14.3%	16.7%
ステージIII		22.8%	24.5%	17.6%	34.3%	30.6%
ステージIV		43.6%	49.1%	45.1%	42.9%	27.8%
不明		4.6%	3.8%	7.8%	0.0%	8.3%
計		100.0%	100.0%	100.0%	100.0%	100.0%
気管支および肺の悪性新生物	(人)	518	66	38	41	44
ステージ0		N/A	0.0%	0.0%	0.0%	0.0%
ステージI		18.7%	19.7%	13.2%	22.0%	22.7%
ステージII		4.1%	4.5%	7.9%	2.4%	0.0%
ステージIII		21.6%	19.7%	23.7%	24.4%	29.5%
ステージIV		35.7%	33.3%	39.5%	29.3%	34.1%
不明		19.9%	22.7%	15.8%	22.0%	13.6%
計		100.0%	100.0%	100.0%	100.0%	100.0%
急性心筋梗塞	(人)	289	28	23	21	28
Killip I		53.6%	50.0%	47.8%	47.6%	64.3%
Killip II		22.1%	35.7%	17.4%	42.9%	14.3%
Killip III		8.0%	7.1%	13.0%	9.5%	7.1%
Killip IV		14.5%	7.1%	21.7%	0.0%	14.3%
不明		1.7%	0.0%	0.0%	0.0%	0.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%
肺炎	(人)	1,858	205	202	188	169
軽症		30.1%	34.1%	28.7%	30.3%	33.7%
中等症		9.1%	8.3%	12.4%	8.0%	10.7%
重症		44.0%	42.4%	51.0%	48.9%	40.8%
不明		16.8%	15.1%	7.9%	12.8%	14.8%
計		100.0%	100.0%	100.0%	100.0%	100.0%
喘息	(人)	274	29	21	16	20
軽症間欠型		32.1%	55.2%	38.1%	12.5%	25.0%
軽症持続型		24.1%	6.9%	23.8%	56.3%	30.0%
中等症持続型		31.0%	24.1%	28.6%	31.3%	25.0%
重症持続型		12.4%	13.8%	9.5%	0.0%	15.0%
不明		0.4%	0.0%	0.0%	0.0%	5.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%
脳梗塞	(人)	1,184	139	106	113	99
JCS 0-1		62.3%	67.6%	62.3%	55.8%	70.7%
JCS 2-3		27.4%	22.3%	25.5%	31.0%	23.2%
JCS 10-100		8.0%	7.2%	9.4%	9.7%	4.0%

【重症度別集計】表

(単位:%)		平均	退院年月							
			2016/04	2016/05	2016/06	2016/07	2016/08	2016/09	2016/10	2016/11
脳梗塞	JCS 200-300	2.2%	1.7%	1.0%	3.4%	1.3%	0.0%	2.5%	2.6%	1.9%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
脳出血	(人)	480	36	38	38	34	33	31	33	51
	JCS 0-1	40.8%	44.4%	44.7%	42.1%	35.3%	36.4%	41.9%	39.4%	43.1%
	JCS 2-3	21.5%	27.8%	28.9%	13.2%	35.3%	21.2%	9.7%	15.2%	25.5%
	JCS 10-100	20.8%	19.4%	15.8%	18.4%	11.8%	21.2%	22.6%	15.2%	21.6%
	JCS 200-300	16.9%	8.3%	10.5%	26.3%	17.6%	21.2%	25.8%	30.3%	9.8%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
糖尿病	(人)	348	33	21	41	27	24	39	37	27
	軽症	53.4%	60.6%	61.9%	51.2%	40.7%	58.3%	59.0%	51.4%	63.0%
	中等症	31.6%	24.2%	23.8%	36.6%	44.4%	29.2%	28.2%	35.1%	25.9%
	重症	14.9%	15.2%	14.3%	12.2%	14.8%	12.5%	12.8%	13.5%	11.1%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
大腿骨頸部骨折	(人)	393	35	30	33	25	25	28	35	28
	重症度Ⅰ	46.3%	45.7%	63.3%	60.6%	44.0%	60.0%	32.1%	45.7%	39.3%
	重症度Ⅱ	26.5%	31.4%	10.0%	18.2%	28.0%	20.0%	21.4%	22.9%	39.3%
	重症度Ⅲ	20.4%	14.3%	23.3%	15.2%	20.0%	16.0%	32.1%	22.9%	21.4%
	重症度Ⅳ	6.4%	8.6%	3.3%	6.1%	8.0%	4.0%	14.3%	8.6%	0.0%
	不明	0.5%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
胃潰瘍	(人)	138	8	12	11	19	6	9	14	10
	ショックなし(収縮期血圧90mmHg以上)	89.9%	87.5%	91.7%	90.9%	94.7%	100.0%	88.9%	85.7%	100.0%
	ショックあり(収縮期血圧90mmHg未満)	10.1%	12.5%	8.3%	9.1%	5.3%	0.0%	11.1%	14.3%	0.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
急性腸炎	(人)	559	47	52	42	54	45	41	41	53
	血便なし	90.5%	91.5%	92.3%	90.5%	85.2%	86.7%	87.8%	90.2%	92.5%
	血便あり	9.3%	8.5%	7.7%	9.5%	14.8%	11.1%	12.2%	9.8%	7.5%
	不明	0.2%	0.0%	0.0%	0.0%	0.0%	2.2%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
急性虫垂炎	(人)	377	28	47	39	30	38	27	40	26
	白血球15000/mm ³ 未満	73.5%	71.4%	63.8%	71.8%	70.0%	71.1%	77.8%	77.5%	76.9%
	白血球15000/mm ³ 以上	26.3%	28.6%	34.0%	28.2%	30.0%	28.9%	22.2%	22.5%	23.1%
	不明	0.3%	0.0%	2.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
胆石症	(人)	955	75	73	75	81	81	85	93	92
	黄疸なし(T Bill 5mg/dl 未満)	91.6%	89.3%	94.5%	93.3%	92.6%	90.1%	91.8%	92.5%	88.0%
	黄疸あり(T Bill 5mg/dl 以上)	8.4%	10.7%	5.5%	6.7%	7.4%	9.9%	8.2%	7.5%	12.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
前立腺肥大症	(人)	54	4	5	8	10	0	6	2	5
	前立腺容積20ml未満	1.9%	0.0%	0.0%	0.0%	10.0%	0.0%	0.0%	0.0%	0.0%
	前立腺容積50ml未満	31.5%	50.0%	20.0%	25.0%	30.0%	0.0%	50.0%	0.0%	20.0%
	前立腺容積50ml以上	50.0%	50.0%	60.0%	62.5%	50.0%	0.0%	33.3%	50.0%	60.0%
	不明	16.7%	0.0%	20.0%	12.5%	10.0%	0.0%	16.7%	50.0%	20.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	
白内障	(人)	588	39	40	57	43	40	44	64	40
	1眼(処置対象のもの)	99.8%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	98.4%	100.0%
	2眼(処置対象のもの)	0.2%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.6%	0.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
痔核	(人)	109	6	8	8	10	6	6	10	11
	第一度(出血のみ)	25.7%	33.3%	37.5%	25.0%	30.0%	16.7%	33.3%	20.0%	36.4%
	第二度(還納可能)	21.1%	0.0%	12.5%	25.0%	0.0%	0.0%	16.7%	60.0%	18.2%
	第三度(脱肛)	50.5%	66.7%	37.5%	50.0%	70.0%	83.3%	50.0%	20.0%	36.4%
	不明	2.8%	0.0%	12.5%	0.0%	0.0%	0.0%	0.0%	0.0%	9.1%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
子宮筋腫	(人)	115	6	10	6	7	11	5	14	9
	貧血なし(Hb10g/dl以上)	81.7%	100.0%	50.0%	100.0%	100.0%	100.0%	80.0%	78.6%	88.9%
	貧血あり(Hb10g/dl未満)	18.3%	0.0%	50.0%	0.0%	0.0%	0.0%	20.0%	21.4%	11.1%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
狭心症	(人)	1,649	120	120	160	134	122	126	135	142
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
腎結石及び尿管結石	(人)	352	27	25	22	30	40	37	29	31
	水腎・水尿管なし	51.1%	51.9%	44.0%	40.9%	56.7%	55.0%	48.6%	62.1%	51.6%
	水腎・水尿管あり	46.9%	48.1%	56.0%	59.1%	43.3%	40.0%	45.9%	34.5%	48.4%

【重症度別集計】表

(単位:%)		平均	退院年月			
			2016/12	2017/01	2017/02	2017/03
脳梗塞	JCS 200-300	2.2%	2.9%	2.8%	3.5%	2.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
脳出血	(人)	480	52	36	45	53
	JCS 0-1	40.8%	50.0%	36.1%	31.1%	41.5%
	JCS 2-3	21.5%	19.2%	22.2%	28.9%	11.3%
	JCS 10-100	20.8%	19.2%	25.0%	24.4%	30.2%
	JCS 200-300	16.9%	11.5%	16.7%	15.6%	17.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
糖尿病	(人)	348	28	19	19	33
	軽症	53.4%	39.3%	36.8%	68.4%	51.5%
	中等症	31.6%	32.1%	42.1%	26.3%	30.3%
	重症	14.9%	28.6%	21.1%	5.3%	18.2%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
大腿骨頸部骨折	(人)	393	47	44	31	32
	重症度Ⅰ	46.3%	46.8%	36.4%	45.2%	40.6%
	重症度Ⅱ	26.5%	23.4%	36.4%	35.5%	28.1%
	重症度Ⅲ	20.4%	25.5%	18.2%	19.4%	15.6%
	重症度Ⅳ	6.4%	4.3%	6.8%	0.0%	12.5%
	不明	0.5%	0.0%	2.3%	0.0%	3.1%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
胃潰瘍	(人)	138	20	7	10	12
	ショックなし(収縮期血圧90mmHg以上)	89.9%	90.0%	71.4%	80.0%	91.7%
	ショックあり(収縮期血圧90mmHg未満)	10.1%	10.0%	28.6%	20.0%	8.3%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
急性腸炎	(人)	559	58	50	34	42
	血便なし	90.5%	98.3%	92.0%	82.4%	92.9%
	血便あり	9.3%	1.7%	8.0%	17.6%	7.1%
	不明	0.2%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
急性虫垂炎	(人)	377	29	24	27	22
	白血球15000/mm3未満	73.5%	79.3%	83.3%	88.9%	54.5%
	白血球15000/mm3以上	26.3%	20.7%	16.7%	11.1%	45.5%
	不明	0.3%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
胆石症	(人)	955	83	76	66	75
	黄疸なし(T Bill 5mg/dl 未満)	91.6%	90.4%	89.5%	93.9%	94.7%
	黄疸あり(T Bill 5mg/dl 以上)	8.4%	9.6%	10.5%	6.1%	5.3%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
前立腺肥大症	(人)	54	6	4	3	1
	前立腺容積20ml未満	1.9%	0.0%	0.0%	0.0%	0.0%
	前立腺容積50ml未満	31.5%	0.0%	75.0%	33.3%	100.0%
	前立腺容積50ml以上	50.0%	66.7%	0.0%	66.7%	0.0%
	不明	16.7%	33.3%	25.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
白内障	(人)	588	47	51	61	62
	1眼(処置対象のもの)	99.8%	100.0%	100.0%	100.0%	100.0%
	2眼(処置対象のもの)	0.2%	0.0%	0.0%	0.0%	0.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
痔核	(人)	109	8	4	11	21
	第一度(出血のみ)	25.7%	25.0%	0.0%	9.1%	28.6%
	第二度(還納可能)	21.1%	25.0%	0.0%	9.1%	38.1%
	第三度(脱肛)	50.5%	50.0%	75.0%	81.8%	33.3%
	不明	2.8%	0.0%	25.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
子宮筋腫	(人)	115	10	12	12	13
	貧血なし(Hb10g/dl以上)	81.7%	60.0%	91.7%	91.7%	61.5%
	貧血あり(Hb10g/dl未満)	18.3%	40.0%	8.3%	8.3%	38.5%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
狭心症	(人)	1,649	147	127	144	172
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
腎結石及び尿管結石	(人)	352	32	19	29	31
	水腎・水尿管なし	51.1%	50.0%	73.7%	34.5%	48.4%
	水腎・水尿管あり	46.9%	43.8%	26.3%	65.5%	51.6%

【重症度別集計】表

(単位:%)		平均	退院年月							
			2016/04	2016/05	2016/06	2016/07	2016/08	2016/09	2016/10	2016/11
腎結石及び尿管結石	不明	2.0%	0.0%	0.0%	0.0%	0.0%	5.0%	5.4%	3.4%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
乳房の悪性新生物	(人)	436	22	37	23	48	37	33	41	36
	ステージ0	6.7%	9.1%	2.7%	13.0%	2.1%	2.7%	9.1%	2.4%	11.1%
	ステージI	34.6%	40.9%	27.0%	30.4%	39.6%	32.4%	30.3%	41.5%	30.6%
	ステージII	29.8%	13.6%	35.1%	34.8%	31.3%	40.5%	30.3%	31.7%	27.8%
	ステージIII	12.6%	13.6%	13.5%	8.7%	10.4%	8.1%	12.1%	4.9%	11.1%
	ステージIV	7.6%	13.6%	13.5%	4.3%	8.3%	8.1%	12.1%	12.2%	8.3%
	不明	8.7%	9.1%	8.1%	8.7%	8.3%	8.1%	6.1%	7.3%	11.1%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
膝関節症	(人)	77	24	18	3	2	1	7	6	6
	骨移植なし	89.6%	100.0%	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	66.7%
	骨移植あり	10.4%	0.0%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	33.3%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
鼠径ヘルニア	(人)	535	31	41	49	51	51	46	39	53
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
全疾患合計		100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
集計病院数	-		9	10	10	10	10	10	11	11

【重症度別集計】表

		平均	退院年月			
			2016/12	2017/01	2017/02	2017/03
腎結石及び尿管結石	不明	2.0%	6.3%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
乳房の悪性新生物	(人)	436	37	44	34	44
	ステージ0	6.7%	2.7%	6.8%	2.9%	18.2%
	ステージI	34.6%	56.8%	25.0%	29.4%	31.8%
	ステージII	29.8%	16.2%	38.6%	32.4%	20.5%
	ステージIII	12.6%	10.8%	11.4%	23.5%	22.7%
	ステージIV	7.6%	0.0%	6.8%	2.9%	2.3%
	不明	8.7%	13.5%	11.4%	8.8%	4.5%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
膝関節症	(人)	77	2	1	3	4
	骨移植なし	89.6%	0.0%	100.0%	66.7%	75.0%
	骨移植あり	10.4%	100.0%	0.0%	33.3%	25.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
鼠径ヘルニア	(人)	535	55	33	42	44
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
全疾患合計		100.0%	0.0%	0.0%	0.0%	0.0%
集計病院数		-	11	11	10	10