

# 【重症度別集計】表

## 絞込条件

対象年月設定: 201704 ~ 201803  
 病院の経営主体選択: 全て  
 【詳細】病床数: 下限なし ~ 上限なし  
 【詳細】手術の有無: 全て  
 【詳細】ステージ集約選択: 有り  
 【詳細】年代: 全て  
 【詳細】性別: 全て  
 【詳細】入院経路: 全て  
 並べ替え条件: 退院年月・昇順

## ■ 退院患者数

(単位: %)	平均	退院年月								
		2017/04	2017/05	2017/06	2017/07	2017/08	2017/09	2017/10	2017/11	
胃の悪性新生物	(人)	491	36	31	38	47	46	43	49	37
	ステージ0	0.2%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	2.7%
	ステージI	32.6%	33.3%	29.0%	36.8%	25.5%	37.0%	30.2%	36.7%	35.1%
	ステージII	11.2%	16.7%	6.5%	10.5%	6.4%	13.0%	9.3%	12.2%	13.5%
	ステージIII	13.0%	5.6%	12.9%	10.5%	12.8%	8.7%	14.0%	6.1%	13.5%
	ステージIV	36.3%	41.7%	38.7%	31.6%	44.7%	30.4%	34.9%	38.8%	29.7%
	不明	6.7%	2.8%	12.9%	10.5%	10.6%	10.9%	11.6%	6.1%	5.4%
結腸の悪性新生物	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	755	67	71	73	75	61	65	60	68
	ステージ0	1.6%	1.5%	1.4%	0.0%	1.3%	0.0%	3.1%	1.7%	0.0%
	ステージI	7.2%	13.4%	12.7%	6.8%	2.7%	4.9%	4.6%	11.7%	14.7%
	ステージII	19.3%	20.9%	18.3%	27.4%	25.3%	16.4%	13.8%	16.7%	16.2%
	ステージIII	34.6%	34.3%	35.2%	32.9%	32.0%	41.0%	32.3%	35.0%	33.8%
	ステージIV	28.9%	23.9%	22.5%	23.3%	32.0%	27.9%	33.8%	31.7%	25.0%
直腸の悪性新生物	不明	8.5%	6.0%	9.9%	9.6%	6.7%	9.8%	12.3%	3.3%	10.3%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	424	36	36	35	28	32	41	34	33
	ステージ0	2.4%	5.6%	5.6%	5.7%	0.0%	3.1%	0.0%	5.9%	0.0%
	ステージI	12.5%	11.1%	5.6%	22.9%	10.7%	15.6%	14.6%	11.8%	15.2%
	ステージII	15.3%	13.9%	22.2%	8.6%	7.1%	25.0%	29.3%	23.5%	15.2%
	ステージIII	26.2%	41.7%	22.2%	34.3%	46.4%	18.8%	14.6%	20.6%	12.1%
気管支および肺の悪性新生物	ステージIV	39.4%	25.0%	33.3%	25.7%	35.7%	31.3%	39.0%	35.3%	51.5%
	不明	4.2%	2.8%	11.1%	2.9%	0.0%	6.3%	2.4%	2.9%	6.1%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	515	33	29	46	46	41	54	31	34
	ステージ0	1.2%	0.0%	6.9%	2.2%	0.0%	2.4%	0.0%	6.5%	0.0%
	ステージI	18.6%	24.2%	13.8%	19.6%	17.4%	19.5%	20.4%	16.1%	17.6%
	ステージII	8.0%	6.1%	6.9%	6.5%	2.2%	4.9%	9.3%	16.1%	8.8%
急性心筋梗塞	ステージIII	18.8%	18.2%	13.8%	10.9%	17.4%	22.0%	24.1%	16.1%	14.7%
	ステージIV	38.1%	33.3%	48.3%	37.0%	41.3%	39.0%	22.2%	38.7%	35.3%
	不明	15.3%	18.2%	10.3%	23.9%	21.7%	12.2%	24.1%	6.5%	23.5%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	306	30	28	22	31	19	28	16	29
	Killip I	52.0%	43.3%	50.0%	77.3%	54.8%	63.2%	64.3%	50.0%	41.4%
	Killip II	28.8%	26.7%	32.1%	13.6%	25.8%	21.1%	21.4%	25.0%	34.5%
肺炎	Killip III	6.2%	13.3%	0.0%	4.5%	3.2%	5.3%	10.7%	6.3%	10.3%
	Killip IV	10.1%	6.7%	17.9%	0.0%	16.1%	5.3%	3.6%	12.5%	10.3%
	不明	2.9%	10.0%	0.0%	4.5%	0.0%	5.3%	0.0%	6.3%	3.4%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	1,687	158	183	141	135	143	108	107	119
	軽症	29.9%	34.8%	37.2%	27.7%	33.3%	33.6%	37.0%	24.3%	25.2%
	中等症	9.3%	8.2%	5.5%	12.1%	9.6%	11.9%	8.3%	11.2%	11.8%
喘息	重症	48.0%	46.8%	45.9%	46.1%	46.7%	47.6%	43.5%	43.9%	44.5%
	不明	12.8%	10.1%	11.5%	14.2%	10.4%	7.0%	11.1%	20.6%	18.5%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	257	20	31	22	26	15	34	25	21
	軽症間欠型	30.7%	15.0%	19.4%	45.5%	34.6%	40.0%	47.1%	32.0%	33.3%
	軽症持続型	22.6%	20.0%	32.3%	31.8%	15.4%	13.3%	17.6%	8.0%	14.3%
	中等症持続型	31.5%	40.0%	35.5%	13.6%	19.2%	40.0%	23.5%	44.0%	33.3%
脳梗塞	重症持続型	12.5%	20.0%	12.9%	9.1%	23.1%	6.7%	11.8%	12.0%	19.0%
	不明	2.7%	5.0%	0.0%	0.0%	7.7%	0.0%	0.0%	4.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	1,218	111	108	111	104	118	106	107	90
	JCS 0-1	65.6%	64.0%	72.2%	65.8%	60.6%	78.0%	58.5%	65.4%	60.0%
	JCS 2-3	24.1%	26.1%	20.4%	20.7%	26.0%	14.4%	31.1%	25.2%	28.9%
	JCS 10-100	8.5%	8.1%	6.5%	10.8%	8.7%	5.9%	7.5%	7.5%	10.0%

# 【重症度別集計】表

(単位:%)		平均	退院年月			
			2017/12	2018/01	2018/02	2018/03
胃の悪性新生物	(人)	491	49	45	40	30
	ステージ0	0.2%	0.0%	0.0%	0.0%	0.0%
	ステージI	32.6%	32.7%	33.3%	37.5%	20.0%
	ステージII	11.2%	14.3%	8.9%	12.5%	10.0%
	ステージIII	13.0%	16.3%	15.6%	22.5%	20.0%
	ステージIV	36.3%	34.7%	40.0%	25.0%	46.7%
	不明	6.7%	2.0%	2.2%	2.5%	3.3%
計		100.0%	100.0%	100.0%	100.0%	100.0%
結腸の悪性新生物	(人)	755	56	48	56	55
	ステージ0	1.6%	1.8%	2.1%	3.6%	3.6%
	ステージI	7.2%	1.8%	4.2%	1.8%	3.6%
	ステージII	19.3%	19.6%	16.7%	17.9%	20.0%
	ステージIII	34.6%	28.6%	39.6%	42.9%	29.1%
	ステージIV	28.9%	37.5%	33.3%	25.0%	34.5%
	不明	8.5%	10.7%	4.2%	8.9%	9.1%
計		100.0%	100.0%	100.0%	100.0%	100.0%
直腸の悪性新生物	(人)	424	43	37	45	24
	ステージ0	2.4%	0.0%	0.0%	2.2%	0.0%
	ステージI	12.5%	11.6%	5.4%	13.3%	12.5%
	ステージII	15.3%	11.6%	8.1%	11.1%	4.2%
	ステージIII	26.2%	23.3%	32.4%	26.7%	25.0%
	ステージIV	39.4%	48.8%	51.4%	40.0%	58.3%
	不明	4.2%	4.7%	2.7%	6.7%	0.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%
気管支および肺の悪性新生物	(人)	515	48	46	48	59
	ステージ0	1.2%	0.0%	0.0%	0.0%	0.0%
	ステージI	18.6%	22.9%	15.2%	20.8%	15.3%
	ステージII	8.0%	12.5%	6.5%	8.3%	8.5%
	ステージIII	18.8%	20.8%	13.0%	22.9%	25.4%
	ステージIV	38.1%	35.4%	45.7%	43.8%	40.7%
	不明	15.3%	8.3%	19.6%	4.2%	10.2%
計		100.0%	100.0%	100.0%	100.0%	100.0%
急性心筋梗塞	(人)	306	31	17	28	27
	Killip I	52.0%	38.7%	64.7%	46.4%	44.4%
	Killip II	28.8%	41.9%	23.5%	28.6%	40.7%
	Killip III	6.2%	9.7%	0.0%	3.6%	3.7%
	Killip IV	10.1%	9.7%	5.9%	17.9%	11.1%
	不明	2.9%	0.0%	5.9%	3.6%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
肺炎	(人)	1,687	152	156	164	121
	軽症	29.9%	25.0%	26.3%	28.7%	23.1%
	中等症	9.3%	7.2%	12.2%	6.7%	9.1%
	重症	48.0%	52.0%	48.1%	51.2%	57.9%
	不明	12.8%	15.8%	13.5%	13.4%	9.9%
計		100.0%	100.0%	100.0%	100.0%	100.0%
喘息	(人)	257	19	19	14	11
	軽症間欠型	30.7%	26.3%	10.5%	28.6%	27.3%
	軽症持続型	22.6%	26.3%	42.1%	21.4%	36.4%
	中等症持続型	31.5%	31.6%	31.6%	42.9%	36.4%
	重症持続型	12.5%	10.5%	5.3%	7.1%	0.0%
	不明	2.7%	5.3%	10.5%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
脳梗塞	(人)	1,218	113	93	81	76
	JCS 0-1	65.6%	65.5%	60.2%	59.3%	76.3%
	JCS 2-3	24.1%	23.9%	29.0%	32.1%	13.2%
	JCS 10-100	8.5%	10.6%	8.6%	8.6%	9.2%

## 【重症度別集計】表

(単位: %)		平均	退院年月							
			2017/04	2017/05	2017/06	2017/07	2017/08	2017/09	2017/10	2017/11
脳梗塞	JCS 200-300	1.8%	1.8%	0.9%	2.7%	4.8%	1.7%	2.8%	1.9%	1.1%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
脳出血	(人)	432	43	48	41	43	34	36	29	23
	JCS 0-1	37.7%	41.9%	35.4%	34.1%	32.6%	23.5%	38.9%	44.8%	43.5%
	JCS 2-3	20.8%	25.6%	20.8%	26.8%	25.6%	11.8%	22.2%	20.7%	21.7%
	JCS 10-100	24.1%	16.3%	31.3%	22.0%	27.9%	35.3%	16.7%	13.8%	17.4%
	JCS 200-300	17.4%	16.3%	12.5%	17.1%	14.0%	29.4%	22.2%	20.7%	17.4%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
糖尿病	(人)	345	40	33	23	27	37	32	25	24
	軽症	60.9%	70.0%	63.6%	69.6%	55.6%	45.9%	75.0%	76.0%	62.5%
	中等症	23.8%	17.5%	18.2%	26.1%	29.6%	35.1%	9.4%	16.0%	25.0%
	重症	13.3%	12.5%	15.2%	4.3%	11.1%	16.2%	9.4%	8.0%	8.3%
	不明	2.0%	0.0%	3.0%	0.0%	3.7%	2.7%	6.3%	0.0%	4.2%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
大腿骨頸部骨折	(人)	379	31	35	38	31	34	27	24	40
	重症度 I	41.7%	32.3%	31.4%	44.7%	41.9%	55.9%	59.3%	37.5%	25.0%
	重症度 II	27.7%	29.0%	22.9%	42.1%	22.6%	14.7%	18.5%	25.0%	42.5%
	重症度 III	22.2%	29.0%	31.4%	7.9%	29.0%	26.5%	11.1%	29.2%	20.0%
	重症度 IV	6.9%	3.2%	11.4%	5.3%	6.5%	2.9%	11.1%	4.2%	10.0%
	不明	1.6%	6.5%	2.9%	0.0%	0.0%	0.0%	0.0%	4.2%	2.5%
胃潰瘍	(人)	133	10	8	10	8	12	14	11	20
	ショックなし(収縮期血圧90mmHg以上)	92.5%	100.0%	100.0%	100.0%	100.0%	91.7%	78.6%	90.9%	90.0%
	ショックあり(収縮期血圧90mmHg未満)	7.5%	0.0%	0.0%	0.0%	0.0%	8.3%	21.4%	9.1%	10.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
急性腸炎	(人)	487	54	48	60	45	38	47	32	30
	血便なし	91.2%	98.1%	87.5%	96.7%	84.4%	81.6%	93.6%	90.6%	96.7%
	血便あり	8.8%	1.9%	12.5%	3.3%	15.6%	18.4%	6.4%	9.4%	3.3%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
急性虫垂炎	(人)	368	29	46	25	29	25	41	50	31
	白血球15000/mm <sup>3</sup> 未満	71.7%	44.8%	76.1%	80.0%	93.1%	64.0%	63.4%	66.0%	74.2%
	白血球15000/mm <sup>3</sup> 以上	28.3%	55.2%	23.9%	20.0%	6.9%	36.0%	36.6%	34.0%	25.8%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
胆石症	(人)	910	83	76	82	84	83	88	73	96
	黄疸なし(T Bill 5mg/dl 未満)	91.4%	88.0%	89.5%	95.1%	90.5%	90.4%	94.3%	91.8%	89.6%
	黄疸あり(T Bill 5mg/dl 以上)	8.6%	12.0%	10.5%	4.9%	9.5%	9.6%	5.7%	8.2%	10.4%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
前立腺肥大症	(人)	43	3	2	1	5	5	5	3	6
	前立腺容積20ml未満	11.6%	0.0%	0.0%	0.0%	0.0%	0.0%	60.0%	33.3%	0.0%
	前立腺容積50ml未満	32.6%	0.0%	50.0%	0.0%	20.0%	60.0%	0.0%	33.3%	16.7%
	前立腺容積50ml以上	41.9%	100.0%	50.0%	0.0%	60.0%	20.0%	40.0%	33.3%	50.0%
	不明	14.0%	0.0%	0.0%	100.0%	20.0%	20.0%	0.0%	0.0%	33.3%
白内障	(人)	530	41	39	62	46	49	53	48	58
	1眼(処置対象のもの)	99.8%	100.0%	100.0%	100.0%	97.8%	100.0%	100.0%	100.0%	100.0%
	2眼(処置対象のもの)	0.2%	0.0%	0.0%	0.0%	2.2%	0.0%	0.0%	0.0%	0.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
痔核	(人)	105	10	7	15	8	7	10	10	4
	第一度(出血のみ)	20.0%	20.0%	28.6%	26.7%	12.5%	0.0%	40.0%	20.0%	0.0%
	第二度(還納可能)	37.1%	40.0%	28.6%	40.0%	50.0%	57.1%	10.0%	30.0%	75.0%
	第三度(脱肛)	41.9%	30.0%	42.9%	33.3%	37.5%	42.9%	50.0%	50.0%	25.0%
	不明	1.0%	10.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
子宮筋腫	(人)	118	14	8	9	13	17	13	10	5
	貧血なし(Hb10g/dl以上)	58.5%	57.1%	62.5%	88.9%	30.8%	52.9%	53.8%	80.0%	60.0%
	貧血あり(Hb10g/dl未満)	41.5%	42.9%	37.5%	11.1%	69.2%	47.1%	46.2%	20.0%	40.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
狭心症	(人)	1,534	135	132	144	149	123	137	122	117
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
腎結石及び尿管結石	(人)	306	20	18	30	40	36	25	27	23
	水腎・水尿管なし	47.7%	50.0%	61.1%	50.0%	45.0%	50.0%	36.0%	48.1%	47.8%
	水腎・水尿管あり	52.3%	50.0%	38.9%	50.0%	55.0%	50.0%	64.0%	51.9%	52.2%

## 【重症度別集計】表

(単位:%)		平均	退院年月			
			2017/12	2018/01	2018/02	2018/03
脳梗塞	JCS 200-300	1.8%	0.0%	2.2%	0.0%	1.3%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
脳出血	(人)	432	29	32	40	34
	JCS 0-1	37.7%	41.4%	46.9%	30.0%	47.1%
	JCS 2-3	20.8%	20.7%	12.5%	22.5%	14.7%
	JCS 10-100	24.1%	24.1%	28.1%	32.5%	17.6%
	JCS 200-300	17.4%	13.8%	12.5%	15.0%	20.6%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
糖尿病	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	345	29	21	30	24
	軽症	60.9%	55.2%	33.3%	60.0%	58.3%
	中等症	23.8%	24.1%	38.1%	26.7%	25.0%
	重症	13.3%	17.2%	28.6%	13.3%	16.7%
	不明	2.0%	3.4%	0.0%	0.0%	0.0%
大腿骨頸部骨折	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	379	37	38	22	22
	重症度 I	41.7%	40.5%	44.7%	50.0%	45.5%
	重症度 II	27.7%	18.9%	39.5%	18.2%	27.3%
	重症度 III	22.2%	24.3%	13.2%	22.7%	27.3%
	重症度 IV	6.9%	13.5%	2.6%	9.1%	0.0%
臓潰瘍	不明	1.6%	2.7%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	133	14	7	10	9
	ショックなし(収縮期血圧90mmHg以上)	92.5%	92.9%	85.7%	90.0%	100.0%
	ショックあり(収縮期血圧90mmHg未満)	7.5%	7.1%	14.3%	10.0%	0.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
急性腸炎	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	487	36	29	35	33
	血便なし	91.2%	91.7%	93.1%	85.7%	90.9%
	血便あり	8.8%	8.3%	6.9%	14.3%	9.1%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
急性虫垂炎	(人)	368	32	26	19	15
	白血球15000/mm <sup>3</sup> 未満	71.7%	84.4%	76.9%	63.2%	80.0%
	白血球15000/mm <sup>3</sup> 以上	28.3%	15.6%	23.1%	36.8%	20.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	910	80	44	55	66
胆石症	黄疸なし(T Bill 5mg/dl未満)	91.4%	92.5%	93.2%	90.9%	92.4%
	黄疸あり(T Bill 5mg/dl以上)	8.6%	7.5%	6.8%	9.1%	7.6%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	43	1	3	5	4
	前立腺容積20ml未満	11.6%	0.0%	33.3%	0.0%	0.0%
前立腺肥大症	前立腺容積50ml未満	32.6%	0.0%	33.3%	40.0%	100.0%
	前立腺容積50ml以上	41.9%	0.0%	33.3%	60.0%	0.0%
	不明	14.0%	100.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	530	31	38	48	17
	1眼(処置対象のもの)	99.8%	100.0%	100.0%	100.0%	100.0%
白内障	2眼(処置対象のもの)	0.2%	0.0%	0.0%	0.0%	0.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	105	12	8	6	8
	第一度(出血のみ)	20.0%	25.0%	12.5%	16.7%	12.5%
	第二度(還納可能)	37.1%	58.3%	25.0%	33.3%	12.5%
痔核	第三度(脱肛)	41.9%	16.7%	62.5%	50.0%	75.0%
	不明	1.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	118	7	13	7	2
	貧血なし(Hb10g/dl以上)	58.5%	42.9%	61.5%	71.4%	50.0%
	貧血あり(Hb10g/dl未満)	41.5%	57.1%	38.5%	28.6%	50.0%
子宮筋腫	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	1,534	124	114	119	118
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%
狭心症	計	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	306	31	19	25	12
	水腎・水尿管なし	47.7%	38.7%	42.1%	56.0%	58.3%
	水腎・水尿管あり	52.3%	61.3%	57.9%	44.0%	41.7%
腎結石及び尿管結石	(人)					

## 【重症度別集計】表

(単位: %)		平均	退院年月							
			2017/04	2017/05	2017/06	2017/07	2017/08	2017/09	2017/10	2017/11
腎結石及び尿管結石	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
乳房の悪性新生物	(人)	389	43	29	34	29	38	42	32	33
	ステージⅠ	8.7%	7.0%	17.2%	8.8%	3.4%	7.9%	2.4%	0.0%	6.1%
	ステージⅡ	33.4%	23.3%	31.0%	32.4%	31.0%	26.3%	26.2%	62.5%	24.2%
	ステージⅢ	33.4%	37.2%	31.0%	38.2%	37.9%	42.1%	40.5%	21.9%	42.4%
	ステージⅣ	13.4%	18.6%	10.3%	14.7%	17.2%	13.2%	16.7%	0.0%	15.2%
	不明	6.7%	9.3%	10.3%	2.9%	3.4%	0.0%	7.1%	3.1%	9.1%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
膝関節症	(人)	104	2	0	3	3	4	4	2	2
	骨移植なし	89.4%	100.0%	0.0%	33.3%	33.3%	75.0%	50.0%	100.0%	50.0%
	骨移植あり	10.6%	0.0%	0.0%	66.7%	66.7%	25.0%	50.0%	0.0%	50.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
鼠径ヘルニア	計	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	(人)	481	36	48	51	53	41	48	42	26
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
全疾患合計		100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
集計病院数			-	10	10	10	10	10	9	9

## 【重症度別集計】表

(単位:%)		平均	退院年月			
			2017/12	2018/01	2018/02	2018/03
腎結石及び尿管結石	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
乳房の悪性新生物	(人)	389	32	26	28	23
	ステージⅠ	8.7%	15.6%	15.4%	17.9%	8.7%
	ステージⅡ	33.4%	40.6%	23.1%	46.4%	43.5%
	ステージⅢ	33.4%	21.9%	30.8%	21.4%	26.1%
	ステージⅣ	13.4%	15.6%	15.4%	7.1%	13.0%
	不明	4.4%	0.0%	11.5%	7.1%	4.3%
	計	6.7%	6.3%	3.8%	0.0%	4.3%
膝関節症	(人)	104	31	14	18	21
	骨移植なし	89.4%	96.8%	92.9%	94.4%	100.0%
	骨移植あり	10.6%	3.2%	7.1%	5.6%	0.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
鼠径ヘルニア	(人)	481	41	31	36	28
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
全疾患合計		100.0%	0.0%	0.0%	0.0%	0.0%
集計病院数	-		9	7	7	6