

【重症度別集計】表

絞込条件

対象年月設定: 201804 ~ 201903
 病院の経営主体選択: 全て
 【詳細】病床数: 下限なし ~ 上限なし
 【詳細】手術の有無: 全て
 【詳細】ステージ集約選択: 有り
 【詳細】年代: 全て
 【詳細】性別: 全て
 【詳細】入院経路: 全て
 並べ替え条件: 退院年月・昇順

■ 退院患者数

	(単位:%)	平均	退院年月							
			2018/04	2018/05	2018/06	2018/07	2018/08	2018/09	2018/10	2018/11
胃の悪性新生物	(人)	-	-	-	-	-	-	-	-	-
ステージ0		0.3%	0.0%	0.0%	0.0%	2.2%	0.0%	0.0%	0.0%	0.0%
ステージI		31.3%	47.8%	31.0%	26.3%	21.7%	29.0%	35.5%	29.7%	34.9%
ステージII		10.0%	4.3%	10.3%	13.2%	6.5%	6.5%	9.7%	8.1%	9.3%
ステージIII		14.5%	21.7%	10.3%	21.1%	17.4%	19.4%	12.9%	21.6%	11.6%
ステージIV		38.6%	26.1%	41.4%	34.2%	45.7%	38.7%	32.3%	29.7%	41.9%
不明		5.3%	0.0%	6.9%	5.3%	6.5%	6.5%	9.7%	10.8%	2.3%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
結腸の悪性新生物	(人)	-	-	-	-	-	-	-	53	-
ステージ0		4.3%	0.0%	0.0%	2.2%	3.6%	0.0%	0.0%	9.4%	10.0%
ステージI		8.5%	5.5%	5.1%	6.5%	10.9%	10.6%	2.6%	11.3%	17.5%
ステージII		15.4%	20.0%	15.4%	21.7%	14.5%	23.4%	21.1%	9.4%	12.5%
ステージIII		36.5%	27.3%	30.8%	34.8%	30.9%	27.7%	42.1%	34.0%	40.0%
ステージIV		30.1%	43.6%	41.0%	30.4%	36.4%	27.7%	28.9%	28.3%	17.5%
不明		5.2%	3.6%	7.7%	4.3%	3.6%	10.6%	5.3%	7.5%	2.5%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
直腸の悪性新生物	(人)	-	-	-	-	-	-	-	-	-
ステージ0		1.8%	0.0%	3.4%	0.0%	3.8%	0.0%	4.8%	0.0%	3.2%
ステージI		8.5%	8.0%	10.3%	12.9%	11.5%	22.6%	4.8%	10.0%	6.5%
ステージII		16.7%	16.0%	13.8%	9.7%	11.5%	9.7%	23.8%	25.0%	22.6%
ステージIII		30.3%	36.0%	20.7%	29.0%	19.2%	29.0%	28.6%	40.0%	45.2%
ステージIV		38.8%	36.0%	51.7%	41.9%	50.0%	35.5%	38.1%	20.0%	22.6%
不明		3.9%	4.0%	0.0%	6.5%	3.8%	3.2%	0.0%	5.0%	0.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
気管支および肺の悪性新生物	(人)	-	-	-	-	-	-	-	-	-
ステージ0		1.0%	0.0%	0.0%	1.8%	0.0%	2.4%	2.6%	2.6%	2.0%
ステージI		17.3%	17.1%	17.9%	14.3%	13.8%	19.5%	15.8%	17.9%	15.7%
ステージII		6.4%	2.4%	7.7%	10.7%	1.7%	12.2%	10.5%	7.7%	2.0%
ステージIII		19.6%	36.6%	12.8%	14.3%	12.1%	17.1%	21.1%	17.9%	25.5%
ステージIV		39.4%	31.7%	51.3%	48.2%	46.6%	39.0%	34.2%	30.8%	39.2%
不明		16.3%	12.2%	10.3%	10.7%	25.9%	9.8%	15.8%	23.1%	15.7%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
急性心筋梗塞	(人)	-	-	-	-	-	-	-	-	-
Killip I		52.5%	46.7%	50.0%	47.4%	44.4%	45.0%	69.6%	40.0%	40.0%
Killip II		23.2%	26.7%	30.0%	31.6%	33.3%	35.0%	13.0%	26.7%	20.0%
Killip III		7.2%	6.7%	5.0%	5.3%	0.0%	5.0%	4.3%	6.7%	16.0%
Killip IV		14.1%	13.3%	15.0%	10.5%	16.7%	15.0%	13.0%	13.3%	20.0%
不明		3.0%	6.7%	0.0%	5.3%	5.6%	0.0%	0.0%	13.3%	4.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
肺炎	(人)	-	85	82	76	87	-	-	83	-
軽症		30.7%	25.9%	29.3%	39.5%	34.5%	35.4%	34.8%	34.9%	35.0%
中等症		9.8%	9.4%	9.8%	6.6%	11.5%	7.6%	8.7%	9.6%	7.5%
重症		56.4%	56.5%	57.3%	50.0%	46.0%	57.0%	56.5%	51.8%	56.3%
不明		3.1%	8.2%	3.7%	3.9%	8.0%	0.0%	0.0%	3.6%	1.3%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
喘息	(人)	-	-	-	-	-	-	-	-	-
軽症間欠型		19.8%	14.3%	40.0%	27.3%	30.0%	15.4%	4.5%	10.0%	41.2%
軽症持続型		23.4%	28.6%	20.0%	27.3%	30.0%	15.4%	22.7%	20.0%	23.5%
中等症持続型		43.7%	28.6%	10.0%	36.4%	40.0%	53.8%	72.7%	55.0%	23.5%
重症持続型		11.4%	21.4%	20.0%	9.1%	0.0%	15.4%	0.0%	10.0%	11.8%
不明		1.8%	7.1%	10.0%	0.0%	0.0%	0.0%	0.0%	5.0%	0.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
脳梗塞	(人)	-	-	-	-	-	-	-	-	-
JCS 0-1		66.8%	70.3%	73.8%	65.8%	72.8%	72.2%	65.9%	61.8%	67.7%
JCS 2-3		24.7%	21.9%	19.0%	28.9%	22.2%	20.8%	22.0%	28.9%	25.3%
JCS 10-100		6.5%	7.8%	4.8%	3.9%	4.9%	4.2%	9.8%	5.3%	4.0%

【重症度別集計】表

	(単位:%)	平均	退院年月			
			2018/12	2019/01	2019/02	2019/03
胃の悪性新生物	(人)	-	-	-	-	-
ステージ0		0.3%	0.0%	0.0%	0.0%	0.0%
ステージI		31.3%	37.9%	25.8%	37.0%	29.4%
ステージII		10.0%	10.3%	3.2%	29.6%	11.8%
ステージIII		14.5%	3.4%	9.7%	11.1%	11.8%
ステージIV		38.6%	44.8%	54.8%	22.2%	44.1%
不明		5.3%	3.4%	6.5%	0.0%	2.9%
計		100.0%	100.0%	100.0%	100.0%	100.0%
結腸の悪性新生物	(人)	-	-	-	-	-
ステージ0		4.3%	5.3%	8.1%	9.3%	4.2%
ステージI		8.5%	10.5%	10.8%	4.7%	6.3%
ステージII		15.4%	7.9%	10.8%	7.0%	18.8%
ステージIII		36.5%	50.0%	45.9%	44.2%	39.6%
ステージIV		30.1%	23.7%	18.9%	32.6%	25.0%
不明		5.2%	2.6%	5.4%	2.3%	6.3%
計		100.0%	100.0%	100.0%	100.0%	100.0%
直腸の悪性新生物	(人)	-	-	-	-	-
ステージ0		1.8%	0.0%	3.6%	0.0%	3.6%
ステージI		8.5%	0.0%	3.6%	6.5%	3.6%
ステージII		16.7%	17.2%	25.0%	16.1%	14.3%
ステージIII		30.3%	37.9%	28.6%	32.3%	17.9%
ステージIV		38.8%	37.9%	39.3%	38.7%	50.0%
不明		3.9%	6.9%	0.0%	6.5%	10.7%
計		100.0%	100.0%	100.0%	100.0%	100.0%
気管支および肺の悪性新生物	(人)	-	-	-	-	-
ステージ0		1.0%	0.0%	0.0%	0.0%	0.0%
ステージI		17.3%	17.1%	17.6%	22.0%	22.2%
ステージII		6.4%	9.8%	5.9%	4.9%	2.8%
ステージIII		19.6%	14.6%	26.5%	19.5%	22.2%
ステージIV		39.4%	36.6%	35.3%	36.6%	36.1%
不明		16.3%	22.0%	14.7%	17.1%	16.7%
計		100.0%	100.0%	100.0%	100.0%	100.0%
急性心筋梗塞	(人)	-	-	-	-	-
Killip I		52.5%	64.7%	52.9%	54.5%	58.3%
Killip II		23.2%	17.6%	35.3%	15.2%	12.5%
Killip III		7.2%	8.8%	5.9%	12.1%	4.2%
Killip IV		14.1%	5.9%	5.9%	18.2%	20.8%
不明		3.0%	2.9%	0.0%	0.0%	4.2%
計		100.0%	100.0%	100.0%	100.0%	100.0%
肺炎	(人)	-	73	160	-	-
軽症		30.7%	28.8%	25.6%	26.5%	27.1%
中等症		9.8%	9.6%	10.6%	11.8%	11.8%
重症		56.4%	57.5%	60.6%	60.3%	61.2%
不明		3.1%	4.1%	3.1%	1.5%	0.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%
喘息	(人)	-	-	-	-	-
軽症間欠型		19.8%	16.7%	0.0%	35.3%	50.0%
軽症持続型		23.4%	8.3%	31.6%	29.4%	0.0%
中等症持続型		43.7%	58.3%	52.6%	23.5%	50.0%
重症持続型		11.4%	16.7%	15.8%	11.8%	0.0%
不明		1.8%	0.0%	0.0%	0.0%	0.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%
脳梗塞	(人)	-	-	-	-	-
JCS 0-1		66.8%	61.3%	64.9%	54.7%	68.1%
JCS 2-3		24.7%	28.0%	28.4%	32.8%	20.9%
JCS 10-100		6.5%	10.7%	6.8%	9.4%	7.7%

【重症度別集計】表

(単位:%)		平均	退院年月							
			2018/04	2018/05	2018/06	2018/07	2018/08	2018/09	2018/10	2018/11
脳梗塞	JCS 200-300	1.9%	0.0%	2.4%	1.3%	0.0%	2.8%	2.4%	3.9%	3.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
脳出血	(人)	-	-	-	-	-	-	-	-	-
	JCS 0-1	38.9%	41.4%	52.2%	39.3%	32.3%	36.4%	51.6%	29.6%	33.3%
	JCS 2-3	23.4%	24.1%	26.1%	17.9%	12.9%	27.3%	22.6%	37.0%	19.4%
	JCS 10-100	19.0%	24.1%	17.4%	21.4%	32.3%	13.6%	16.1%	7.4%	25.0%
	JCS 200-300	18.7%	10.3%	4.3%	21.4%	22.6%	22.7%	9.7%	25.9%	22.2%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
糖尿病	(人)	-	-	-	-	-	-	-	-	-
	軽症	58.7%	56.5%	46.7%	89.5%	47.4%	55.0%	70.0%	59.1%	50.0%
	中等症	22.2%	26.1%	13.3%	10.5%	10.5%	30.0%	20.0%	22.7%	25.0%
	重症	18.7%	17.4%	40.0%	0.0%	42.1%	15.0%	10.0%	13.6%	25.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
大腿骨頸部骨折	(人)	-	-	-	-	-	-	-	-	-
	重症度Ⅰ	50.7%	45.0%	52.2%	42.1%	60.9%	75.0%	54.5%	30.0%	40.7%
	重症度Ⅱ	14.6%	15.0%	21.7%	10.5%	8.7%	14.3%	27.3%	10.0%	3.7%
	重症度Ⅲ	25.5%	25.0%	8.7%	42.1%	30.4%	7.1%	13.6%	60.0%	37.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
胃潰瘍	(人)	-	-	-	-	-	-	-	-	-
	ショックなし(収縮期血圧90mmHg以上)	92.1%	100.0%	100.0%	100.0%	100.0%	83.3%	100.0%	100.0%	80.0%
	ショックあり(収縮期血圧90mmHg未満)	7.9%	0.0%	0.0%	0.0%	0.0%	16.7%	0.0%	0.0%	20.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
急性腸炎	(人)	-	-	-	-	-	-	-	-	-
	血便なし	84.9%	63.6%	87.5%	90.0%	73.7%	71.4%	100.0%	85.7%	87.5%
	血便あり	15.1%	36.4%	12.5%	10.0%	26.3%	28.6%	0.0%	14.3%	12.5%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
急性虫垂炎	(人)	-	-	-	-	-	-	-	-	-
	白血球15000/mm3未満	68.0%	50.0%	50.0%	100.0%	52.6%	81.8%	50.0%	71.4%	50.0%
	白血球15000/mm3以上	32.0%	50.0%	50.0%	0.0%	47.4%	18.2%	50.0%	28.6%	50.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
胆石症	(人)	-	-	-	-	-	-	-	-	-
	黄疸なし(T Bill 5mg/dl 未満)	94.2%	95.4%	90.9%	93.3%	97.2%	96.7%	96.2%	92.6%	90.8%
	黄疸あり(T Bill 5mg/dl 以上)	5.8%	4.6%	9.1%	6.7%	2.8%	3.3%	3.8%	7.4%	9.2%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
前立腺肥大症	(人)	-	-	-	-	-	-	-	-	-
	前立腺容積20ml未満	5.7%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	50.0%
	前立腺容積50ml未満	25.7%	0.0%	0.0%	20.0%	20.0%	100.0%	0.0%	50.0%	25.0%
	前立腺容積50ml以上	54.3%	75.0%	50.0%	80.0%	80.0%	0.0%	100.0%	50.0%	25.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
白内障	(人)	-	-	-	-	-	-	-	-	-
	1眼(処置対象のもの)	92.2%	100.0%	97.4%	100.0%	86.2%	91.3%	85.7%	94.4%	82.8%
	2眼(処置対象のもの)	7.8%	0.0%	2.6%	0.0%	13.8%	8.7%	14.3%	5.6%	17.2%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
痔核	(人)	-	-	-	-	-	-	-	-	-
	第一度(出血のみ)	20.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	50.0%
	第二度(還納可能)	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	第三度(脱肛)	73.3%	0.0%	100.0%	0.0%	100.0%	100.0%	0.0%	0.0%	50.0%
	計	100.0%	0.0%	100.0%	0.0%	100.0%	100.0%	0.0%	100.0%	100.0%
子宮筋腫	(人)	-	-	-	-	-	-	-	-	-
	貧血なし(Hb10g/dl以上)	78.2%	50.0%	71.4%	83.3%	90.0%	55.6%	83.3%	77.8%	90.9%
	貧血あり(Hb10g/dl未満)	21.8%	50.0%	28.6%	16.7%	10.0%	44.4%	16.7%	22.2%	9.1%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
狭心症	(人)	1,328	100	117	118	136	124	100	103	132
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
腎結石及び尿管結石	(人)	-	-	-	-	-	-	-	-	-
	水腎・水尿管なし	55.6%	30.3%	66.7%	52.0%	58.7%	58.5%	59.3%	40.6%	54.5%
	水腎・水尿管あり	44.4%	69.7%	33.3%	48.0%	41.3%	41.5%	40.7%	59.4%	45.5%

【重症度別集計】表

(単位:%)		平均	退院年月			
			2018/12	2019/01	2019/02	2019/03
脳梗塞	JCS 200-300	1.9%	0.0%	0.0%	3.1%	3.3%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
脳出血	(人)	-	-	-	-	-
	JCS 0-1	38.9%	47.8%	44.0%	39.5%	24.1%
	JCS 2-3	23.4%	13.0%	36.0%	18.4%	31.0%
	JCS 10-100	19.0%	17.4%	8.0%	21.1%	17.2%
	JCS 200-300	18.7%	21.7%	12.0%	21.1%	27.6%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
糖尿病	(人)	-	-	-	-	-
	軽症	58.7%	68.8%	41.2%	73.3%	47.4%
	中等症	22.2%	6.3%	47.1%	13.3%	36.8%
	重症	18.7%	25.0%	11.8%	13.3%	15.8%
	不明	0.4%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
大腿骨頸部骨折	(人)	-	-	-	-	-
	重症度Ⅰ	50.7%	40.0%	53.3%	62.5%	40.0%
	重症度Ⅱ	14.6%	20.0%	20.0%	12.5%	15.0%
	重症度Ⅲ	25.5%	28.0%	20.0%	18.8%	25.0%
	重症度Ⅳ	8.0%	8.0%	6.7%	6.3%	20.0%
	不明	1.1%	4.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
胃潰瘍	(人)	-	-	-	-	-
	ショックなし(収縮期血圧90mmHg以上)	92.1%	100.0%	85.7%	50.0%	87.5%
	ショックあり(収縮期血圧90mmHg未満)	7.9%	0.0%	14.3%	50.0%	12.5%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
急性腸炎	(人)	-	-	-	-	-
	血便なし	84.9%	95.8%	83.3%	92.9%	90.9%
	血便あり	15.1%	4.2%	16.7%	7.1%	9.1%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
急性虫垂炎	(人)	-	-	-	-	-
	白血球15000/mm3未満	68.0%	100.0%	66.7%	66.7%	75.0%
	白血球15000/mm3以上	32.0%	0.0%	33.3%	33.3%	25.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
胆石症	(人)	-	-	-	-	-
	黄疸なし(T Bill 5mg/dl 未満)	94.2%	97.4%	93.1%	92.7%	93.0%
	黄疸あり(T Bill 5mg/dl 以上)	5.8%	2.6%	6.9%	7.3%	7.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
前立腺肥大症	(人)	-	-	-	-	-
	前立腺容積20ml未満	5.7%	0.0%	0.0%	0.0%	0.0%
	前立腺容積50ml未満	25.7%	50.0%	0.0%	0.0%	66.7%
	前立腺容積50ml以上	54.3%	25.0%	33.3%	100.0%	33.3%
	不明	14.3%	25.0%	66.7%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
白内障	(人)	-	-	-	-	-
	1眼(処置対象のもの)	92.2%	70.0%	100.0%	95.7%	91.7%
	2眼(処置対象のもの)	7.8%	30.0%	0.0%	4.3%	8.3%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
痔核	(人)	-	-	-	-	-
	第一度(出血のみ)	20.0%	0.0%	0.0%	33.3%	50.0%
	第二度(還納可能)	N/A	0.0%	0.0%	0.0%	0.0%
	第三度(脱肛)	73.3%	0.0%	100.0%	66.7%	50.0%
	不明	6.7%	0.0%	0.0%	0.0%	0.0%
計	100.0%	0.0%	100.0%	100.0%	100.0%	
子宮筋腫	(人)	-	-	-	-	-
	貧血なし(Hb10g/dl以上)	78.2%	87.5%	77.8%	87.5%	75.0%
	貧血あり(Hb10g/dl未満)	21.8%	12.5%	22.2%	12.5%	25.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
狭心症	(人)	1,328	98	108	100	92
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
腎結石及び尿管結石	(人)	-	-	-	-	-
	水腎・水尿管なし	55.6%	57.1%	64.1%	57.7%	71.4%
	水腎・水尿管あり	44.4%	42.9%	35.9%	42.3%	28.6%

【重症度別集計】表

(単位:%)		平均	退院年月							
			2018/04	2018/05	2018/06	2018/07	2018/08	2018/09	2018/10	2018/11
腎結石及び尿管結石	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
乳房の悪性新生物	(人)	-	-	-	-	-	-	-	-	-
	ステージ0	8.8%	9.1%	10.0%	5.0%	7.4%	0.0%	13.3%	18.5%	9.1%
	ステージI	43.1%	40.9%	45.0%	35.0%	29.6%	44.0%	40.0%	44.4%	40.9%
	ステージII	28.8%	40.9%	25.0%	30.0%	33.3%	32.0%	26.7%	25.9%	31.8%
	ステージIII	13.1%	4.5%	10.0%	20.0%	22.2%	16.0%	6.7%	3.7%	13.6%
	ステージIV	4.6%	4.5%	10.0%	0.0%	7.4%	4.0%	13.3%	3.7%	4.5%
	不明	1.5%	0.0%	0.0%	10.0%	0.0%	4.0%	0.0%	3.7%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
膝関節症	(人)	-	-	-	-	-	-	-	-	-
	骨移植なし	97.3%	100.0%	90.3%	100.0%	96.6%	94.7%	94.1%	95.8%	100.0%
	骨移植あり	2.7%	0.0%	9.7%	0.0%	3.4%	5.3%	5.9%	4.2%	0.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
鼠径ヘルニア	(人)	423	33	32	34	38	36	41	39	36
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
全疾患合計		100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
集計病院数		-	6	6	6	6	6	5	6	6

【重症度別集計】表

		平均	退院年月			
			2018/12	2019/01	2019/02	2019/03
腎結石及び尿管結石	不明	N/A	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%
乳房の悪性新生物	(人)	-	-	-	-	-
	ステージ0	8.8%	5.3%	11.8%	0.0%	14.3%
	ステージI	43.1%	52.6%	58.8%	47.1%	50.0%
	ステージII	28.8%	36.8%	11.8%	29.4%	14.3%
	ステージIII	13.1%	5.3%	17.6%	23.5%	21.4%
	ステージIV	4.6%	0.0%	0.0%	0.0%	0.0%
	不明	1.5%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
膝関節症	(人)	-	-	-	-	-
	骨移植なし	97.3%	100.0%	100.0%	100.0%	100.0%
	骨移植あり	2.7%	0.0%	0.0%	0.0%	0.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
鼠径ヘルニア	(人)	423	35	33	30	36
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	
全疾患合計		100.0%	0.0%	0.0%	0.0%	0.0%
集計病院数		-	6	6	6	6