

# 【重症度別集計】表

## 絞込条件

対象年月設定: 201904 ~ 202003  
 病院の経営主体選択: 全て  
 【詳細】病床数: 下限なし ~ 上限なし  
 【詳細】手術の有無: 全て  
 【詳細】ステージ集約選択: 有り  
 【詳細】年代: 全て  
 【詳細】性別: 全て  
 【詳細】入院経路: 全て  
 並べ替え条件: 退院年月・昇順

## ■ 退院患者数

| (単位: %)       | 平均         | 退院年月    |         |         |         |         |         |         |         |
|---------------|------------|---------|---------|---------|---------|---------|---------|---------|---------|
|               |            | 2019/04 | 2019/05 | 2019/06 | 2019/07 | 2019/08 | 2019/09 | 2019/10 | 2019/11 |
| 胃の悪性新生物       | (人)        | -       | -       | -       | -       | -       | -       | -       | -       |
|               | ステージ0      | 1.1%    | 0.0%    | 0.0%    | 3.7%    | 3.2%    | 3.7%    | 0.0%    | 0.0%    |
|               | ステージI      | 35.5%   | 29.7%   | 34.3%   | 25.9%   | 35.5%   | 22.2%   | 41.4%   | 34.6%   |
|               | ステージII     | 12.6%   | 8.1%    | 20.0%   | 18.5%   | 16.1%   | 14.8%   | 13.8%   | 7.7%    |
|               | ステージIII    | 12.8%   | 13.5%   | 5.7%    | 3.7%    | 12.9%   | 7.4%    | 13.8%   | 19.2%   |
|               | ステージIV     | 31.8%   | 45.9%   | 34.3%   | 40.7%   | 29.0%   | 40.7%   | 24.1%   | 30.8%   |
|               | 不明         | 6.1%    | 2.7%    | 5.7%    | 7.4%    | 3.2%    | 11.1%   | 6.9%    | 7.7%    |
| 結腸の悪性新生物      | 計          | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
|               | (人)        | -       | -       | -       | -       | -       | -       | 53      | 46      |
|               | ステージ0      | 3.9%    | 2.0%    | 1.9%    | 1.7%    | 4.3%    | 0.0%    | 5.7%    | 10.9%   |
|               | ステージI      | 11.0%   | 8.2%    | 13.0%   | 19.0%   | 6.4%    | 10.3%   | 7.5%    | 13.0%   |
|               | ステージII     | 15.6%   | 12.2%   | 16.7%   | 15.5%   | 19.1%   | 10.3%   | 11.3%   | 8.7%    |
|               | ステージIII    | 30.0%   | 40.8%   | 38.9%   | 29.3%   | 23.4%   | 24.1%   | 39.6%   | 30.4%   |
|               | ステージIV     | 34.0%   | 34.7%   | 25.9%   | 27.6%   | 40.4%   | 41.4%   | 26.4%   | 30.4%   |
| 直腸の悪性新生物      | 不明         | 5.5%    | 2.0%    | 3.7%    | 6.9%    | 6.4%    | 13.8%   | 9.4%    | 6.5%    |
|               | 計          | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
|               | (人)        | -       | -       | -       | -       | -       | -       | -       | -       |
|               | ステージ0      | 4.1%    | 3.1%    | 0.0%    | 5.9%    | 2.6%    | 6.9%    | 0.0%    | 0.0%    |
|               | ステージI      | 11.0%   | 0.0%    | 6.5%    | 17.6%   | 10.5%   | 13.8%   | 23.5%   | 6.9%    |
|               | ステージII     | 19.9%   | 28.1%   | 16.1%   | 17.6%   | 23.7%   | 20.7%   | 32.4%   | 17.2%   |
|               | ステージIII    | 42.3%   | 50.0%   | 51.6%   | 38.2%   | 42.1%   | 48.3%   | 20.6%   | 41.4%   |
| 気管支および肺の悪性新生物 | ステージIV     | 17.4%   | 15.6%   | 16.1%   | 14.7%   | 21.1%   | 10.3%   | 14.7%   | 24.1%   |
|               | 不明         | 5.2%    | 3.1%    | 9.7%    | 5.9%    | 0.0%    | 0.0%    | 8.8%    | 10.3%   |
|               | 計          | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
|               | (人)        | -       | -       | -       | -       | -       | -       | -       | -       |
|               | ステージ0      | 0.9%    | 2.9%    | 0.0%    | 0.0%    | 1.9%    | 0.0%    | 0.0%    | 0.0%    |
|               | ステージI      | 21.3%   | 8.8%    | 6.3%    | 16.7%   | 25.0%   | 20.0%   | 23.0%   | 22.9%   |
|               | ステージII     | 5.9%    | 2.9%    | 6.3%    | 5.6%    | 1.9%    | 7.5%    | 4.9%    | 2.9%    |
| 急性心筋梗塞        | ステージIII    | 19.3%   | 26.5%   | 21.9%   | 27.8%   | 13.5%   | 17.5%   | 16.4%   | 11.4%   |
|               | ステージIV     | 33.8%   | 47.1%   | 37.5%   | 25.0%   | 44.2%   | 40.0%   | 32.8%   | 31.4%   |
|               | 不明         | 18.7%   | 11.8%   | 28.1%   | 25.0%   | 13.5%   | 15.0%   | 23.0%   | 31.4%   |
|               | 計          | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
|               | (人)        | -       | -       | -       | -       | -       | -       | -       | -       |
|               | Killip I   | 49.6%   | 50.0%   | 43.8%   | 53.8%   | 50.0%   | 23.1%   | 41.9%   | 63.3%   |
|               | Killip II  | 25.4%   | 21.4%   | 25.0%   | 23.1%   | 15.4%   | 53.8%   | 29.0%   | 13.3%   |
| 肺炎            | Killip III | 3.6%    | 7.1%    | 9.4%    | 7.7%    | 3.8%    | 0.0%    | 0.0%    | 0.0%    |
|               | Killip IV  | 18.5%   | 21.4%   | 21.9%   | 7.7%    | 26.9%   | 23.1%   | 25.8%   | 16.7%   |
|               | 不明         | 2.8%    | 0.0%    | 0.0%    | 7.7%    | 3.8%    | 0.0%    | 3.2%    | 6.7%    |
|               | 計          | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
|               | (人)        | -       | 90      | 121     | 91      | 92      | 83      | -       | 109     |
|               | 軽症         | 31.9%   | 32.2%   | 28.9%   | 35.2%   | 22.8%   | 22.9%   | 45.0%   | 37.6%   |
|               | 中等症        | 9.9%    | 5.6%    | 5.8%    | 7.7%    | 12.0%   | 8.4%    | 9.0%    | 18.3%   |
| 喘息            | 重症         | 53.5%   | 52.2%   | 59.5%   | 52.7%   | 58.7%   | 61.4%   | 45.0%   | 40.4%   |
|               | 不明         | 4.7%    | 10.0%   | 5.8%    | 4.4%    | 6.5%    | 7.2%    | 1.0%    | 3.7%    |
|               | 計          | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
|               | (人)        | -       | -       | -       | -       | -       | -       | -       | -       |
|               | 軽症間欠型      | 16.6%   | 12.5%   | 10.5%   | 35.3%   | 16.7%   | 30.0%   | 16.7%   | 20.0%   |
|               | 軽症持続型      | 18.6%   | 18.8%   | 10.5%   | 29.4%   | 25.0%   | 40.0%   | 16.7%   | 13.3%   |
|               | 中等症持続型     | 40.0%   | 56.3%   | 47.4%   | 23.5%   | 50.0%   | 20.0%   | 50.0%   | 33.3%   |
| 脳梗塞           | 重症持続型      | 23.4%   | 12.5%   | 31.6%   | 11.8%   | 8.3%    | 10.0%   | 8.3%    | 26.7%   |
|               | 不明         | 1.4%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 8.3%    | 6.7%    |
|               | 計          | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
|               | (人)        | -       | -       | -       | -       | -       | -       | -       | -       |
|               | JCS 0-1    | 65.3%   | 69.0%   | 65.5%   | 73.4%   | 60.2%   | 59.4%   | 68.9%   | 66.2%   |
|               | JCS 2-3    | 28.0%   | 27.4%   | 25.9%   | 25.0%   | 31.2%   | 26.1%   | 27.2%   | 26.5%   |
|               | JCS 10-100 | 5.0%    | 2.4%    | 5.2%    | 1.6%    | 7.5%    | 13.0%   | 3.9%    | 4.4%    |

## 【重症度別集計】表

| (単位:%)        |            | 平均     | 退院年月    |         |         |         |
|---------------|------------|--------|---------|---------|---------|---------|
|               |            |        | 2019/12 | 2020/01 | 2020/02 | 2020/03 |
| 胃の悪性新生物       | (人)        | -      | -       | -       | -       | -       |
|               | ステージ0      | 1.1%   | 2.5%    | 0.0%    | 0.0%    | 0.0%    |
|               | ステージI      | 35.5%  | 35.0%   | 26.9%   | 57.7%   | 31.0%   |
|               | ステージII     | 12.6%  | 10.0%   | 11.5%   | 3.8%    | 13.8%   |
|               | ステージIII    | 12.8%  | 10.0%   | 19.2%   | 11.5%   | 20.7%   |
|               | ステージIV     | 31.8%  | 30.0%   | 34.6%   | 23.1%   | 34.5%   |
|               | 不明         | 6.1%   | 12.5%   | 7.7%    | 3.8%    | 0.0%    |
| 計             |            | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 結腸の悪性新生物      | (人)        | -      | -       | -       | -       | -       |
|               | ステージ0      | 3.9%   | 4.0%    | 2.4%    | 0.0%    | 12.8%   |
|               | ステージI      | 11.0%  | 10.0%   | 0.0%    | 11.8%   | 15.4%   |
|               | ステージII     | 15.6%  | 22.0%   | 16.7%   | 19.6%   | 12.8%   |
|               | ステージIII    | 30.0%  | 20.0%   | 26.2%   | 33.3%   | 25.6%   |
|               | ステージIV     | 34.0%  | 38.0%   | 50.0%   | 29.4%   | 33.3%   |
|               | 不明         | 5.5%   | 6.0%    | 4.8%    | 5.9%    | 0.0%    |
| 計             |            | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 直腸の悪性新生物      | (人)        | -      | -       | -       | -       | -       |
|               | ステージ0      | 4.1%   | 5.7%    | 4.3%    | 6.1%    | 14.3%   |
|               | ステージI      | 11.0%  | 14.3%   | 4.3%    | 9.1%    | 19.0%   |
|               | ステージII     | 19.9%  | 14.3%   | 21.7%   | 9.1%    | 14.3%   |
|               | ステージIII    | 42.3%  | 42.9%   | 47.8%   | 48.5%   | 33.3%   |
|               | ステージIV     | 17.4%  | 20.0%   | 13.0%   | 24.2%   | 19.0%   |
|               | 不明         | 5.2%   | 2.9%    | 8.7%    | 3.0%    | 0.0%    |
| 計             |            | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 気管支および肺の悪性新生物 | (人)        | -      | -       | -       | -       | -       |
|               | ステージ0      | 0.9%   | 0.0%    | 0.0%    | 0.0%    | 3.2%    |
|               | ステージI      | 21.3%  | 31.7%   | 33.3%   | 20.7%   | 29.0%   |
|               | ステージII     | 5.9%   | 9.8%    | 6.1%    | 13.8%   | 6.5%    |
|               | ステージIII    | 19.3%  | 24.4%   | 15.2%   | 13.8%   | 22.6%   |
|               | ステージIV     | 33.8%  | 26.8%   | 27.3%   | 37.9%   | 22.6%   |
|               | 不明         | 18.7%  | 7.3%    | 18.2%   | 13.8%   | 16.1%   |
| 計             |            | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 急性心筋梗塞        | (人)        | -      | -       | -       | -       | -       |
|               | Killip I   | 49.6%  | 61.9%   | 45.0%   | 50.0%   | 50.0%   |
|               | Killip II  | 25.4%  | 23.8%   | 35.0%   | 22.2%   | 22.2%   |
|               | Killip III | 3.6%   | 0.0%    | 5.0%    | 5.6%    | 5.6%    |
|               | Killip IV  | 18.5%  | 9.5%    | 10.0%   | 22.2%   | 22.2%   |
|               | 不明         | 2.8%   | 4.8%    | 5.0%    | 0.0%    | 0.0%    |
|               | 計          | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 肺炎            | (人)        | -      | 77      | 79      | -       | -       |
|               | 軽症         | 31.9%  | 28.6%   | 34.2%   | 37.5%   | 28.9%   |
|               | 中等症        | 9.9%   | 16.9%   | 7.6%    | 12.5%   | 5.3%    |
|               | 重症         | 53.5%  | 49.4%   | 54.4%   | 48.6%   | 63.2%   |
|               | 不明         | 4.7%   | 5.2%    | 3.8%    | 1.4%    | 2.6%    |
| 計             |            | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 喘息            | (人)        | -      | -       | -       | -       | -       |
|               | 軽症間欠型      | 16.6%  | 0.0%    | 16.7%   | 15.4%   | 16.7%   |
|               | 軽症持続型      | 18.6%  | 9.1%    | 33.3%   | 7.7%    | 16.7%   |
|               | 中等症持続型     | 40.0%  | 45.5%   | 16.7%   | 30.8%   | 50.0%   |
|               | 重症持続型      | 23.4%  | 45.5%   | 33.3%   | 46.2%   | 16.7%   |
|               | 不明         | 1.4%   | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
|               | 計          | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 脳梗塞           | (人)        | -      | -       | -       | -       | -       |
|               | JCS 0-1    | 65.3%  | 70.3%   | 67.1%   | 64.4%   | 61.4%   |
|               | JCS 2-3    | 28.0%  | 26.7%   | 25.0%   | 27.7%   | 30.0%   |
|               | JCS 10-100 | 5.0%   | 1.0%    | 6.6%    | 5.0%    | 8.6%    |

# 【重症度別集計】表

| (単位: %)   |                        | 平均     | 退院年月    |         |         |         |         |         |         |         |
|-----------|------------------------|--------|---------|---------|---------|---------|---------|---------|---------|---------|
|           |                        |        | 2019/04 | 2019/05 | 2019/06 | 2019/07 | 2019/08 | 2019/09 | 2019/10 | 2019/11 |
| 脳梗塞       | JCS 200-300            | 1.6%   | 1.2%    | 3.4%    | 0.0%    | 1.1%    | 1.4%    | 0.0%    | 2.9%    | 2.2%    |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
|           | 計                      | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 脳出血       | (人)                    | -      | -       | -       | -       | -       | -       | -       | -       | -       |
|           | JCS 0-1                | 39.7%  | 34.4%   | 35.3%   | 37.1%   | 27.6%   | 41.2%   | 48.5%   | 57.6%   | 38.2%   |
|           | JCS 2-3                | 24.9%  | 21.9%   | 20.6%   | 25.7%   | 27.6%   | 35.3%   | 24.2%   | 12.1%   | 26.5%   |
|           | JCS 10-100             | 19.2%  | 28.1%   | 26.5%   | 25.7%   | 27.6%   | 5.9%    | 6.1%    | 9.1%    | 17.6%   |
|           | JCS 200-300            | 16.1%  | 15.6%   | 17.6%   | 11.4%   | 17.2%   | 17.6%   | 21.2%   | 21.2%   | 17.6%   |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
| 糖尿病       | (人)                    | -      | -       | -       | -       | -       | -       | -       | -       | -       |
|           | 軽症                     | 54.0%  | 58.3%   | 63.6%   | 58.8%   | 54.5%   | 50.0%   | 52.6%   | 38.5%   | 52.6%   |
|           | 中等症                    | 28.4%  | 16.7%   | 24.2%   | 23.5%   | 36.4%   | 31.3%   | 31.6%   | 46.2%   | 36.8%   |
|           | 重症                     | 17.6%  | 25.0%   | 12.1%   | 17.6%   | 9.1%    | 18.8%   | 15.8%   | 15.4%   | 10.5%   |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
|           | 計                      | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 大腿骨頸部骨折   | (人)                    | -      | -       | -       | -       | -       | -       | -       | -       | -       |
|           | 重症度 I                  | 54.7%  | 45.5%   | 51.9%   | 45.0%   | 44.4%   | 65.0%   | 60.7%   | 66.7%   | 46.7%   |
|           | 重症度 II                 | 17.1%  | 13.6%   | 11.1%   | 20.0%   | 27.8%   | 15.0%   | 14.3%   | 4.2%    | 36.7%   |
|           | 重症度 III                | 22.1%  | 40.9%   | 29.6%   | 30.0%   | 16.7%   | 20.0%   | 21.4%   | 20.8%   | 13.3%   |
|           | 重症度 IV                 | 6.0%   | 0.0%    | 7.4%    | 5.0%    | 11.1%   | 0.0%    | 3.6%    | 8.3%    | 3.3%    |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
| 胃潰瘍       | (人)                    | -      | -       | -       | -       | -       | -       | -       | -       | -       |
|           | ショックなし (収縮期血圧90mmHg以上) | 89.3%  | 100.0%  | 100.0%  | 100.0%  | 85.7%   | 71.4%   | 100.0%  | 100.0%  | 60.0%   |
|           | ショックあり (収縮期血圧90mmHg未満) | 10.7%  | 0.0%    | 0.0%    | 0.0%    | 14.3%   | 28.6%   | 0.0%    | 0.0%    | 40.0%   |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
|           | 計                      | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
|           | 急性腸炎                   | -      | -       | -       | -       | -       | -       | -       | -       | -       |
| 急性虫垂炎     | (人)                    | -      | -       | -       | -       | -       | -       | -       | -       | -       |
|           | 血便なし                   | 85.9%  | 75.0%   | 90.0%   | 100.0%  | 85.0%   | 81.8%   | 83.3%   | 80.0%   | 81.8%   |
|           | 血便あり                   | 14.1%  | 25.0%   | 10.0%   | 0.0%    | 15.0%   | 18.2%   | 16.7%   | 20.0%   | 18.2%   |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
| 胆石症       | (人)                    | -      | -       | -       | -       | -       | -       | -       | -       | -       |
|           | 黄疸なし(T Bill 5mg/dl 未満) | 93.7%  | 87.3%   | 93.2%   | 92.7%   | 95.5%   | 95.9%   | 90.4%   | 94.8%   | 94.4%   |
|           | 黄疸あり(T Bill 5mg/dl 以上) | 6.3%   | 12.7%   | 6.8%    | 7.3%    | 4.5%    | 4.1%    | 9.6%    | 5.2%    | 5.6%    |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
| 前立腺肥大症    | (人)                    | -      | -       | -       | -       | -       | -       | -       | -       | -       |
|           | 前立腺容積20ml未満            | 11.6%  | 0.0%    | 0.0%    | 0.0%    | 33.3%   | 0.0%    | 0.0%    | 0.0%    | 33.3%   |
|           | 前立腺容積50ml未満            | 37.2%  | 0.0%    | 33.3%   | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 33.3%   |
|           | 前立腺容積50ml以上            | 46.5%  | 100.0%  | 66.7%   | 0.0%    | 33.3%   | 100.0%  | 0.0%    | 0.0%    | 33.3%   |
|           | 不明                     | 4.7%   | 0.0%    | 0.0%    | 0.0%    | 33.3%   | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
| 白内障       | (人)                    | -      | -       | -       | -       | -       | -       | -       | -       | -       |
|           | 1眼(処置対象のもの)            | 76.2%  | 77.8%   | 86.2%   | 88.9%   | 71.4%   | 46.2%   | 88.9%   | 92.0%   | 75.0%   |
|           | 2眼(処置対象のもの)            | 23.8%  | 22.2%   | 13.8%   | 11.1%   | 28.6%   | 53.8%   | 11.1%   | 8.0%    | 25.0%   |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
| 痔核        | (人)                    | -      | -       | -       | -       | -       | -       | -       | -       | -       |
|           | 第一度(出血のみ)              | 33.3%  | 0.0%    | 100.0%  | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 100.0%  | 0.0%    |
|           | 第二度(還納可能)              | 16.7%  | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 100.0%  |
|           | 第三度(脱肛)                | 50.0%  | 100.0%  | 0.0%    | 0.0%    | 0.0%    | 100.0%  | 100.0%  | 0.0%    | 0.0%    |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
| 子宮筋腫      | (人)                    | -      | -       | -       | -       | -       | -       | -       | -       | -       |
|           | 貧血なし(Hb10g/dl以上)       | 77.3%  | 85.7%   | 90.9%   | 75.0%   | 50.0%   | 64.3%   | 77.8%   | 85.7%   | 57.1%   |
|           | 貧血あり(Hb10g/dl未満)       | 22.7%  | 14.3%   | 9.1%    | 25.0%   | 50.0%   | 35.7%   | 22.2%   | 14.3%   | 42.9%   |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
| 狭心症       | (人)                    | 1,107  | 82      | 101     | 137     | 101     | 111     | 97      | 100     | 79      |
|           | 重症度の指標なし               | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
|           | 計                      | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 腎結石及び尿管結石 | (人)                    | -      | -       | -       | -       | -       | -       | -       | -       | -       |
|           | 水腎・水尿管なし               | 47.2%  | 48.3%   | 52.5%   | 54.3%   | 71.1%   | 37.1%   | 52.6%   | 41.7%   | 44.4%   |
|           | 水腎・水尿管あり               | 52.5%  | 51.7%   | 47.5%   | 45.7%   | 28.9%   | 62.9%   | 47.4%   | 58.3%   | 55.6%   |

## 【重症度別集計】表

| (単位:%)    |                        | 平均     | 退院年月    |         |         |         |
|-----------|------------------------|--------|---------|---------|---------|---------|
|           |                        |        | 2019/12 | 2020/01 | 2020/02 | 2020/03 |
| 脳梗塞       | JCS 200-300            | 1.6%   | 2.0%    | 1.3%    | 3.0%    | 0.0%    |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
|           | 計                      | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 脳出血       | (人)                    | -      | -       | -       | -       | -       |
|           | JCS 0-1                | 39.7%  | 43.2%   | 43.3%   | 20.6%   | 48.6%   |
|           | JCS 2-3                | 24.9%  | 24.3%   | 20.0%   | 38.2%   | 27.0%   |
|           | JCS 10-100             | 19.2%  | 18.9%   | 26.7%   | 23.5%   | 10.8%   |
|           | JCS 200-300            | 16.1%  | 13.5%   | 10.0%   | 17.6%   | 13.5%   |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
| 糖尿病       | (人)                    | -      | -       | -       | -       | -       |
|           | 軽症                     | 54.0%  | 44.8%   | 50.0%   | 56.7%   | 61.5%   |
|           | 中等症                    | 28.4%  | 34.5%   | 26.9%   | 23.3%   | 15.4%   |
|           | 重症                     | 17.6%  | 20.7%   | 23.1%   | 20.0%   | 23.1%   |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
|           | 計                      | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 大腿骨頸部骨折   | (人)                    | -      | -       | -       | -       | -       |
|           | 重症度 I                  | 54.7%  | 70.8%   | 60.9%   | 46.3%   | 57.1%   |
|           | 重症度 II                 | 17.1%  | 12.5%   | 13.0%   | 12.2%   | 28.6%   |
|           | 重症度 III                | 22.1%  | 8.3%    | 17.4%   | 29.3%   | 14.3%   |
|           | 重症度 IV                 | 6.0%   | 8.3%    | 8.7%    | 12.2%   | 0.0%    |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
| 胃潰瘍       | (人)                    | -      | -       | -       | -       | -       |
|           | ショックなし (収縮期血圧90mmHg以上) | 89.3%  | 77.8%   | 87.5%   | 100.0%  | 90.9%   |
|           | ショックあり (収縮期血圧90mmHg未満) | 10.7%  | 22.2%   | 12.5%   | 0.0%    | 9.1%    |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
|           | 計                      | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
|           | 急性腸炎                   | -      | -       | -       | -       | -       |
| 急性虫垂炎     | (人)                    | -      | -       | -       | -       | -       |
|           | 血便なし                   | 85.9%  | 88.2%   | 80.0%   | 93.3%   | 100.0%  |
|           | 血便あり                   | 14.1%  | 11.8%   | 20.0%   | 6.7%    | 0.0%    |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
| 胆石症       | (人)                    | -      | -       | -       | -       | -       |
|           | 黄疸なし(T Bill 5mg/dl 未満) | 93.7%  | 94.5%   | 93.0%   | 94.7%   | 98.0%   |
|           | 黄疸あり(T Bill 5mg/dl 以上) | 6.3%   | 5.5%    | 7.0%    | 5.3%    | 2.0%    |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
|           | 計                      | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
|           | 前立腺肥大症                 | -      | -       | -       | -       | -       |
| 白内障       | (人)                    | -      | -       | -       | -       | -       |
|           | 前立腺容積20ml未満            | 11.6%  | 14.3%   | 33.3%   | 14.3%   | 0.0%    |
|           | 前立腺容積50ml未満            | 37.2%  | 28.6%   | 33.3%   | 71.4%   | 60.0%   |
|           | 前立腺容積50ml以上            | 46.5%  | 42.9%   | 33.3%   | 14.3%   | 40.0%   |
|           | 不明                     | 4.7%   | 14.3%   | 0.0%    | 0.0%    | 0.0%    |
| 痔核        | (人)                    | -      | -       | -       | -       | -       |
|           | 第一度(出血のみ)              | 33.3%  | 66.7%   | 0.0%    | 0.0%    | 0.0%    |
|           | 第二度(還納可能)              | 16.7%  | 0.0%    | 50.0%   | 0.0%    | 0.0%    |
|           | 第三度(脱肛)                | 50.0%  | 33.3%   | 50.0%   | 100.0%  | 0.0%    |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
| 子宮筋腫      | (人)                    | -      | -       | -       | -       | -       |
|           | 貧血なし(Hb10g/dl以上)       | 77.3%  | 66.7%   | 88.9%   | 85.7%   | 100.0%  |
|           | 貧血あり(Hb10g/dl未満)       | 22.7%  | 33.3%   | 11.1%   | 14.3%   | 0.0%    |
|           | 不明                     | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
|           | 計                      | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 狭心症       | (人)                    | 1,107  | 76      | 77      | 87      | 59      |
|           | 重症度の指標なし               | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
|           | 計                      | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 腎結石及び尿管結石 | (人)                    | -      | -       | -       | -       | -       |
|           | 水腎・水尿管なし               | 47.2%  | 42.9%   | 26.5%   | 38.7%   | 55.6%   |
|           | 水腎・水尿管あり               | 52.5%  | 57.1%   | 73.5%   | 58.1%   | 44.4%   |

## 【重症度別集計】表

| (単位: %)   |          | 平均     | 退院年月    |         |         |         |         |         |         |         |
|-----------|----------|--------|---------|---------|---------|---------|---------|---------|---------|---------|
|           |          |        | 2019/04 | 2019/05 | 2019/06 | 2019/07 | 2019/08 | 2019/09 | 2019/10 | 2019/11 |
| 腎結石及び尿管結石 | 不明       | 0.3%   | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
|           | 計        | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 乳房の悪性新生物  | (人)      | -      | -       | -       | -       | -       | -       | -       | -       | -       |
|           | ステージⅠ    | 9.8%   | 8.7%    | 9.1%    | 0.0%    | 8.7%    | 4.5%    | 17.4%   | 12.5%   | 7.7%    |
|           | ステージⅡ    | 38.1%  | 34.8%   | 31.8%   | 35.7%   | 39.1%   | 36.4%   | 34.8%   | 33.3%   | 23.1%   |
|           | ステージⅢ    | 30.8%  | 30.4%   | 27.3%   | 32.1%   | 43.5%   | 27.3%   | 26.1%   | 37.5%   | 34.6%   |
|           | ステージⅣ    | 7.9%   | 8.7%    | 13.6%   | 10.7%   | 4.3%    | 4.5%    | 13.0%   | 4.2%    | 15.4%   |
|           | 不明       | 12.4%  | 17.4%   | 18.2%   | 17.9%   | 4.3%    | 27.3%   | 4.3%    | 12.5%   | 15.4%   |
|           | 計        | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 膝関節症      | (人)      | -      | -       | -       | -       | -       | -       | -       | -       | -       |
|           | 骨移植なし    | 90.8%  | 88.2%   | 91.7%   | 92.6%   | 100.0%  | 90.9%   | 87.5%   | 94.4%   | 100.0%  |
|           | 骨移植あり    | 9.2%   | 11.8%   | 8.3%    | 7.4%    | 0.0%    | 9.1%    | 12.5%   | 5.6%    | 0.0%    |
|           | 不明       | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
| 鼠径ヘルニア    | 計        | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
|           | (人)      | 427    | 35      | 34      | 46      | 31      | 50      | 42      | 38      | 30      |
|           | 重症度の指標なし | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 全疾患合計     |          | 100.0% | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
| 集計病院数     |          |        | -       | 6       | 6       | 6       | 6       | 6       | 6       | 5       |

## 【重症度別集計】表

| (単位:%)    |          | 平均     | 退院年月    |         |         |         |
|-----------|----------|--------|---------|---------|---------|---------|
|           |          |        | 2019/12 | 2020/01 | 2020/02 | 2020/03 |
| 腎結石及び尿管結石 | 不明       | 0.3%   | 0.0%    | 0.0%    | 3.2%    | 0.0%    |
|           | 計        | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 乳房の悪性新生物  | (人)      | -      | -       | -       | -       | -       |
|           | ステージⅠ    | 9.8%   | 17.9%   | 10.0%   | 12.2%   | 8.0%    |
|           | ステージⅡ    | 38.1%  | 46.4%   | 46.7%   | 39.0%   | 52.0%   |
|           | ステージⅢ    | 30.8%  | 21.4%   | 23.3%   | 39.0%   | 24.0%   |
|           | ステージⅣ    | 7.9%   | 7.1%    | 10.0%   | 2.4%    | 4.0%    |
|           | 不明       | 12.4%  | 7.1%    | 10.0%   | 7.3%    | 12.0%   |
|           | 計        | 1.0%   | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
| 膝関節症      | (人)      | -      | -       | -       | -       | -       |
|           | 骨移植なし    | 90.8%  | 90.5%   | 85.7%   | 85.2%   | 84.6%   |
|           | 骨移植あり    | 9.2%   | 9.5%    | 14.3%   | 14.8%   | 15.4%   |
|           | 不明       | N/A    | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
|           | 計        | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 鼠径ヘルニア    | (人)      | 427    | 36      | 39      | 30      | 16      |
|           | 重症度の指標なし | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
|           | 計        | 100.0% | 100.0%  | 100.0%  | 100.0%  | 100.0%  |
| 全疾患合計     |          | 100.0% | 0.0%    | 0.0%    | 0.0%    | 0.0%    |
| 集計病院数     |          | -      | 5       | 5       | 5       | 4       |