

【重症度別集計】表

絞込条件

対象年月設定: 201304 ~ 201403
 病院の経営主体選択: 全て
 【詳細】病床数: 下限なし ~ 上限なし
 【詳細】手術の有無: 全て
 【詳細】ステージ集約選択: 有り
 【詳細】年代: 全て
 【詳細】性別: 全て
 【詳細】入院経路: 全て
 並べ替え条件: 病院コード・昇順

■ 退院患者数

	(単位:%)	平均	病院コード							
			0003	0202	0203	0235	1001	1002	1003	1004
胃の悪性新生物	(人)	969	66	0	7	77	72	43	165	8
ステージ0		1.2%	0.0%	0.0%	0.0%	0.0%	6.9%	0.0%	0.0%	12.5%
ステージI		26.7%	22.7%	0.0%	0.0%	36.4%	19.4%	9.3%	38.2%	0.0%
ステージII		10.6%	9.1%	0.0%	0.0%	10.4%	12.5%	16.3%	9.1%	0.0%
ステージIII		13.9%	15.2%	0.0%	0.0%	9.1%	19.4%	9.3%	21.8%	0.0%
ステージIV		36.0%	37.9%	0.0%	0.0%	39.0%	38.9%	53.5%	28.5%	0.0%
不明		11.5%	15.2%	0.0%	100.0%	5.2%	2.8%	11.6%	2.4%	87.5%
計		100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
結腸の悪性新生物	(人)	1,187	90	0	7	116	322	22	125	1
ステージ0		6.3%	4.4%	0.0%	0.0%	3.4%	4.0%	13.6%	2.4%	0.0%
ステージI		10.6%	13.3%	0.0%	0.0%	14.7%	2.2%	4.5%	14.4%	0.0%
ステージII		17.6%	17.8%	0.0%	0.0%	12.1%	20.2%	18.2%	22.4%	0.0%
ステージIII		22.0%	20.0%	0.0%	0.0%	31.0%	28.6%	22.7%	31.2%	0.0%
ステージIV		30.0%	12.2%	0.0%	0.0%	36.2%	38.5%	22.7%	26.4%	0.0%
不明		13.5%	32.2%	0.0%	100.0%	2.6%	6.5%	18.2%	3.2%	100.0%
計		100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
直腸の悪性新生物	(人)	557	46	0	1	51	97	13	90	1
ステージ0		4.8%	8.7%	0.0%	0.0%	3.9%	3.1%	23.1%	3.3%	0.0%
ステージI		12.7%	19.6%	0.0%	0.0%	9.8%	5.2%	23.1%	14.4%	0.0%
ステージII		16.3%	6.5%	0.0%	0.0%	29.4%	15.5%	7.7%	24.4%	0.0%
ステージIII		32.1%	30.4%	0.0%	0.0%	21.6%	47.4%	7.7%	34.4%	0.0%
ステージIV		25.0%	23.9%	0.0%	100.0%	31.4%	27.8%	38.5%	20.0%	0.0%
不明		9.0%	10.9%	0.0%	0.0%	3.9%	1.0%	0.0%	3.3%	100.0%
計		100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
気管支および肺の悪性新生物	(人)	1,312	19	2	9	172	34	50	180	4
ステージ0		0.5%	0.0%	0.0%	0.0%	0.6%	2.9%	0.0%	1.1%	0.0%
ステージI		16.5%	10.5%	0.0%	0.0%	29.7%	23.5%	32.0%	15.0%	0.0%
ステージII		7.9%	5.3%	0.0%	0.0%	7.6%	14.7%	14.0%	5.6%	0.0%
ステージIII		19.4%	31.6%	0.0%	11.1%	20.9%	11.8%	4.0%	18.3%	0.0%
ステージIV		38.8%	47.4%	50.0%	33.3%	18.0%	44.1%	28.0%	38.3%	0.0%
不明		16.9%	5.3%	50.0%	55.6%	23.3%	2.9%	22.0%	21.7%	100.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
急性心筋梗塞	(人)	370	14	6	0	80	47	13	46	1
Killip I		43.5%	42.9%	33.3%	0.0%	51.3%	38.3%	0.0%	2.2%	0.0%
Killip II		32.7%	42.9%	16.7%	0.0%	25.0%	34.0%	84.6%	65.2%	0.0%
Killip III		8.1%	0.0%	0.0%	0.0%	6.3%	8.5%	15.4%	17.4%	0.0%
Killip IV		12.4%	7.1%	33.3%	0.0%	16.3%	12.8%	0.0%	10.9%	0.0%
不明		3.2%	7.1%	16.7%	0.0%	1.3%	6.4%	0.0%	4.3%	100.0%
計		100.0%	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	100.0%
肺炎	(人)	3,253	133	50	149	198	108	185	256	121
軽症		27.6%	43.6%	34.0%	67.8%	28.3%	40.7%	26.5%	4.3%	38.8%
中等症		8.8%	16.5%	12.0%	8.1%	7.1%	12.0%	15.1%	2.7%	24.0%
重症		33.8%	39.8%	54.0%	24.2%	64.1%	47.2%	58.4%	31.3%	31.4%
不明		29.9%	0.0%	0.0%	0.0%	0.5%	0.0%	0.0%	61.7%	5.8%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
喘息	(人)	1,141	13	4	23	63	2	11	84	7
軽症間欠型		26.6%	23.1%	50.0%	26.1%	12.7%	100.0%	54.5%	2.4%	0.0%
軽症持続型		36.6%	23.1%	25.0%	21.7%	22.2%	0.0%	36.4%	3.6%	0.0%
中等症持続型		22.6%	53.8%	0.0%	47.8%	61.9%	0.0%	9.1%	47.6%	0.0%
重症持続型		11.9%	0.0%	0.0%	4.3%	3.2%	0.0%	0.0%	46.4%	0.0%
不明		2.2%	0.0%	25.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
脳梗塞	(人)	1,735	118	403	5	206	1	71	113	35
JCS 0-1		70.5%	67.8%	61.0%	80.0%	69.4%	100.0%	78.9%	72.6%	77.1%
JCS 2-3		18.2%	18.6%	31.3%	0.0%	20.9%	0.0%	7.0%	10.6%	5.7%
JCS 10-100		8.8%	9.3%	6.9%	20.0%	8.3%	0.0%	11.3%	10.6%	14.3%

【重症度別集計】表

	(単位:%)	平均	病院コード							
			1009	1012	1013	1015	1018	1019	1025	1026
胃の悪性新生物	(人)	969	-	68	1	8	-	-	-	-
ステージ0		1.2%	-	1.5%	0.0%	0.0%	-	-	-	-
ステージⅠ		26.7%	-	29.4%	0.0%	50.0%	-	-	-	-
ステージⅡ		10.6%	-	10.3%	0.0%	0.0%	-	-	-	-
ステージⅢ		13.9%	-	16.2%	0.0%	12.5%	-	-	-	-
ステージⅣ		36.0%	-	27.9%	100.0%	12.5%	-	-	-	-
不明		11.5%	-	14.7%	0.0%	25.0%	-	-	-	-
計		100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
結腸の悪性新生物	(人)	1,187	-	43	0	18	-	-	-	-
ステージ0		6.3%	-	20.9%	0.0%	0.0%	-	-	-	-
ステージⅠ		10.6%	-	16.3%	0.0%	16.7%	-	-	-	-
ステージⅡ		17.6%	-	7.0%	0.0%	22.2%	-	-	-	-
ステージⅢ		22.0%	-	30.2%	0.0%	11.1%	-	-	-	-
ステージⅣ		30.0%	-	11.6%	0.0%	11.1%	-	-	-	-
不明		13.5%	-	14.0%	0.0%	38.9%	-	-	-	-
計		100.0%	-	100.0%	0.0%	100.0%	-	-	-	-
直腸の悪性新生物	(人)	557	-	17	0	14	-	-	-	-
ステージ0		4.8%	-	5.9%	0.0%	0.0%	-	-	-	-
ステージⅠ		12.7%	-	11.8%	0.0%	0.0%	-	-	-	-
ステージⅡ		16.3%	-	17.6%	0.0%	7.1%	-	-	-	-
ステージⅢ		32.1%	-	35.3%	0.0%	21.4%	-	-	-	-
ステージⅣ		25.0%	-	23.5%	0.0%	7.1%	-	-	-	-
不明		9.0%	-	5.9%	0.0%	64.3%	-	-	-	-
計		100.0%	-	100.0%	0.0%	100.0%	-	-	-	-
気管支および肺の悪性新生物	(人)	1,312	-	18	344	4	-	-	-	-
ステージ0		0.5%	-	0.0%	0.0%	0.0%	-	-	-	-
ステージⅠ		16.5%	-	5.6%	13.4%	0.0%	-	-	-	-
ステージⅡ		7.9%	-	0.0%	15.1%	0.0%	-	-	-	-
ステージⅢ		19.4%	-	5.6%	25.0%	0.0%	-	-	-	-
ステージⅣ		38.8%	-	61.1%	42.7%	75.0%	-	-	-	-
不明		16.9%	-	27.8%	3.8%	25.0%	-	-	-	-
計		100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
急性心筋梗塞	(人)	370	-	44	1	1	-	-	-	-
KillipⅠ		43.5%	-	61.4%	100.0%	0.0%	-	-	-	-
KillipⅡ		32.7%	-	13.6%	0.0%	0.0%	-	-	-	-
KillipⅢ		8.1%	-	6.8%	0.0%	100.0%	-	-	-	-
KillipⅣ		12.4%	-	18.2%	0.0%	0.0%	-	-	-	-
不明		3.2%	-	0.0%	0.0%	0.0%	-	-	-	-
計		100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
肺炎	(人)	3,253	-	212	133	246	-	-	-	-
軽症		27.6%	-	44.3%	54.1%	30.9%	-	-	-	-
中等症		8.8%	-	9.0%	21.8%	2.0%	-	-	-	-
重症		33.8%	-	46.7%	24.1%	27.2%	-	-	-	-
不明		29.9%	-	0.0%	0.0%	39.8%	-	-	-	-
計		100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
喘息	(人)	1,141	-	50	37	27	-	-	-	-
軽症間欠型		26.6%	-	32.0%	32.4%	88.9%	-	-	-	-
軽症持続型		36.6%	-	28.0%	29.7%	3.7%	-	-	-	-
中等症持続型		22.6%	-	38.0%	21.6%	3.7%	-	-	-	-
重症持続型		11.9%	-	2.0%	13.5%	0.0%	-	-	-	-
不明		2.2%	-	0.0%	2.7%	3.7%	-	-	-	-
計		100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
脳梗塞	(人)	1,735	-	97	0	62	-	-	-	-
JCS 0-1		70.5%	-	74.2%	0.0%	75.8%	-	-	-	-
JCS 2-3		18.2%	-	16.5%	0.0%	9.7%	-	-	-	-
JCS 10-100		8.8%	-	8.2%	0.0%	9.7%	-	-	-	-

【重症度別集計】表

	(単位:%)	平均	病院コード							
			1039	1041	1046	1047	1049	1058	1069	1129
胃の悪性新生物	(人)	969	87	-	3	-	133	-	70	-
ステージ0		1.2%	0.0%	-	0.0%	-	1.5%	-	4.3%	-
ステージI		26.7%	37.9%	-	33.3%	-	11.3%	-	20.0%	-
ステージII		10.6%	10.3%	-	0.0%	-	12.0%	-	11.4%	-
ステージIII		13.9%	9.2%	-	0.0%	-	8.3%	-	14.3%	-
ステージIV		36.0%	41.4%	-	0.0%	-	47.4%	-	47.1%	-
不明		11.5%	1.1%	-	66.7%	-	19.5%	-	2.9%	-
計		100.0%	100.0%	-	100.0%	-	100.0%	-	100.0%	-
結腸の悪性新生物	(人)	1,187	130	-	4	-	99	-	70	-
ステージ0		6.3%	6.2%	-	25.0%	-	4.0%	-	8.6%	-
ステージI		10.6%	20.0%	-	0.0%	-	6.1%	-	7.1%	-
ステージII		17.6%	20.0%	-	0.0%	-	16.2%	-	20.0%	-
ステージIII		22.0%	14.6%	-	0.0%	-	3.0%	-	14.3%	-
ステージIV		30.0%	25.4%	-	0.0%	-	41.4%	-	50.0%	-
不明		13.5%	13.8%	-	75.0%	-	29.3%	-	0.0%	-
計		100.0%	100.0%	-	100.0%	-	100.0%	-	100.0%	-
直腸の悪性新生物	(人)	557	71	-	0	-	40	-	28	-
ステージ0		4.8%	1.4%	-	0.0%	-	0.0%	-	14.3%	-
ステージI		12.7%	9.9%	-	0.0%	-	25.0%	-	3.6%	-
ステージII		16.3%	15.5%	-	0.0%	-	20.0%	-	10.7%	-
ステージIII		32.1%	36.6%	-	0.0%	-	15.0%	-	39.3%	-
ステージIV		25.0%	32.4%	-	0.0%	-	10.0%	-	28.6%	-
不明		9.0%	4.2%	-	0.0%	-	30.0%	-	3.6%	-
計		100.0%	100.0%	-	0.0%	-	100.0%	-	100.0%	-
気管支および肺の悪性新生物	(人)	1,312	138	-	4	-	212	-	11	-
ステージ0		0.5%	0.7%	-	25.0%	-	0.0%	-	0.0%	-
ステージI		16.5%	9.4%	-	0.0%	-	20.3%	-	9.1%	-
ステージII		7.9%	2.9%	-	0.0%	-	3.8%	-	0.0%	-
ステージIII		19.4%	27.5%	-	0.0%	-	13.2%	-	27.3%	-
ステージIV		38.8%	43.5%	-	0.0%	-	43.4%	-	63.6%	-
不明		16.9%	15.9%	-	75.0%	-	19.3%	-	0.0%	-
計		100.0%	100.0%	-	100.0%	-	100.0%	-	100.0%	-
急性心筋梗塞	(人)	370	59	-	4	-	30	-	0	-
Killip I		43.5%	66.1%	-	0.0%	-	60.0%	-	0.0%	-
Killip II		32.7%	20.3%	-	0.0%	-	30.0%	-	0.0%	-
Killip III		8.1%	5.1%	-	0.0%	-	6.7%	-	0.0%	-
Killip IV		12.4%	8.5%	-	75.0%	-	3.3%	-	0.0%	-
不明		3.2%	0.0%	-	25.0%	-	0.0%	-	0.0%	-
計		100.0%	100.0%	-	100.0%	-	100.0%	-	0.0%	-
肺炎	(人)	3,253	812	-	65	-	262	-	80	-
軽症		27.6%	9.2%	-	36.9%	-	40.8%	-	12.5%	-
中等症		8.8%	1.0%	-	13.8%	-	16.8%	-	30.0%	-
重症		33.8%	7.8%	-	49.2%	-	42.4%	-	57.5%	-
不明		29.9%	82.0%	-	0.0%	-	0.0%	-	0.0%	-
計		100.0%	100.0%	-	100.0%	-	100.0%	-	100.0%	-
喘息	(人)	1,141	703	-	5	-	54	-	13	-
軽症間欠型		26.6%	22.0%	-	80.0%	-	85.2%	-	7.7%	-
軽症持続型		36.6%	48.6%	-	0.0%	-	11.1%	-	7.7%	-
中等症持続型		22.6%	17.9%	-	0.0%	-	1.9%	-	23.1%	-
重症持続型		11.9%	11.4%	-	0.0%	-	1.9%	-	53.8%	-
不明		2.2%	0.0%	-	20.0%	-	0.0%	-	7.7%	-
計		100.0%	100.0%	-	100.0%	-	100.0%	-	100.0%	-
脳梗塞	(人)	1,735	218	-	54	-	126	-	7	-
JCS 0-1		70.5%	73.9%	-	88.9%	-	76.2%	-	28.6%	-
JCS 2-3		18.2%	14.7%	-	5.6%	-	6.3%	-	28.6%	-
JCS 10-100		8.8%	7.8%	-	1.9%	-	12.7%	-	28.6%	-

【重症度別集計】表

	(単位:%)	平均	病院コード							
			9009	9010	9011	9012	9013	9014	9015	9016
胃の悪性新生物	(人)	969	-	55	106	-	-	-	-	-
ステージ0		1.2%	-	0.0%	0.0%	-	-	-	-	-
ステージⅠ		26.7%	-	34.5%	27.4%	-	-	-	-	-
ステージⅡ		10.6%	-	5.5%	14.2%	-	-	-	-	-
ステージⅢ		13.9%	-	5.5%	18.9%	-	-	-	-	-
ステージⅣ		36.0%	-	30.9%	24.5%	-	-	-	-	-
不明		11.5%	-	23.6%	15.1%	-	-	-	-	-
計		100.0%	-	100.0%	100.0%	-	-	-	-	-
結腸の悪性新生物	(人)	1,187	-	53	87	-	-	-	-	-
ステージ0		6.3%	-	5.7%	19.5%	-	-	-	-	-
ステージⅠ		10.6%	-	9.4%	21.8%	-	-	-	-	-
ステージⅡ		17.6%	-	17.0%	11.5%	-	-	-	-	-
ステージⅢ		22.0%	-	18.9%	16.1%	-	-	-	-	-
ステージⅣ		30.0%	-	17.0%	18.4%	-	-	-	-	-
不明		13.5%	-	32.1%	12.6%	-	-	-	-	-
計		100.0%	-	100.0%	100.0%	-	-	-	-	-
直腸の悪性新生物	(人)	557	-	20	68	-	-	-	-	-
ステージ0		4.8%	-	0.0%	8.8%	-	-	-	-	-
ステージⅠ		12.7%	-	15.0%	19.1%	-	-	-	-	-
ステージⅡ		16.3%	-	10.0%	10.3%	-	-	-	-	-
ステージⅢ		32.1%	-	25.0%	27.9%	-	-	-	-	-
ステージⅣ		25.0%	-	30.0%	22.1%	-	-	-	-	-
不明		9.0%	-	20.0%	11.8%	-	-	-	-	-
計		100.0%	-	100.0%	100.0%	-	-	-	-	-
気管支および肺の悪性新生物	(人)	1,312	-	85	26	-	-	-	-	-
ステージ0		0.5%	-	0.0%	3.8%	-	-	-	-	-
ステージⅠ		16.5%	-	8.2%	3.8%	-	-	-	-	-
ステージⅡ		7.9%	-	3.5%	0.0%	-	-	-	-	-
ステージⅢ		19.4%	-	12.9%	23.1%	-	-	-	-	-
ステージⅣ		38.8%	-	37.6%	57.7%	-	-	-	-	-
不明		16.9%	-	37.6%	11.5%	-	-	-	-	-
計		100.0%	-	100.0%	100.0%	-	-	-	-	-
急性心筋梗塞	(人)	370	-	15	9	-	-	-	-	-
KillipⅠ		43.5%	-	46.7%	11.1%	-	-	-	-	-
KillipⅡ		32.7%	-	53.3%	22.2%	-	-	-	-	-
KillipⅢ		8.1%	-	0.0%	22.2%	-	-	-	-	-
KillipⅣ		12.4%	-	0.0%	22.2%	-	-	-	-	-
不明		3.2%	-	0.0%	22.2%	-	-	-	-	-
計		100.0%	-	100.0%	100.0%	-	-	-	-	-
肺炎	(人)	3,253	-	53	190	-	-	-	-	-
軽症		27.6%	-	13.2%	25.8%	-	-	-	-	-
中等症		8.8%	-	3.8%	7.4%	-	-	-	-	-
重症		33.8%	-	66.0%	48.9%	-	-	-	-	-
不明		29.9%	-	17.0%	17.9%	-	-	-	-	-
計		100.0%	-	100.0%	100.0%	-	-	-	-	-
喘息	(人)	1,141	-	10	35	-	-	-	-	-
軽症間欠型		26.6%	-	0.0%	48.6%	-	-	-	-	-
軽症持続型		36.6%	-	0.0%	37.1%	-	-	-	-	-
中等症持続型		22.6%	-	0.0%	5.7%	-	-	-	-	-
重症持続型		11.9%	-	0.0%	0.0%	-	-	-	-	-
不明		2.2%	-	100.0%	8.6%	-	-	-	-	-
計		100.0%	-	100.0%	100.0%	-	-	-	-	-
脳梗塞	(人)	1,735	-	50	169	-	-	-	-	-
JCS 0-1		70.5%	-	60.0%	76.3%	-	-	-	-	-
JCS 2-3		18.2%	-	32.0%	13.0%	-	-	-	-	-
JCS 10-100		8.8%	-	6.0%	10.1%	-	-	-	-	-

【重症度別集計】表

	(単位:%)	平均	病院コード				
			9017	9018	9019	9020	9021
胃の悪性新生物	(人)	969	-	-	-	-	-
ステージ0		1.2%	-	-	-	-	-
ステージI		26.7%	-	-	-	-	-
ステージII		10.6%	-	-	-	-	-
ステージIII		13.9%	-	-	-	-	-
ステージIV		36.0%	-	-	-	-	-
不明		11.5%	-	-	-	-	-
計		100.0%	-	-	-	-	-
結腸の悪性新生物	(人)	1,187	-	-	-	-	-
ステージ0		6.3%	-	-	-	-	-
ステージI		10.6%	-	-	-	-	-
ステージII		17.6%	-	-	-	-	-
ステージIII		22.0%	-	-	-	-	-
ステージIV		30.0%	-	-	-	-	-
不明		13.5%	-	-	-	-	-
計		100.0%	-	-	-	-	-
直腸の悪性新生物	(人)	557	-	-	-	-	-
ステージ0		4.8%	-	-	-	-	-
ステージI		12.7%	-	-	-	-	-
ステージII		16.3%	-	-	-	-	-
ステージIII		32.1%	-	-	-	-	-
ステージIV		25.0%	-	-	-	-	-
不明		9.0%	-	-	-	-	-
計		100.0%	-	-	-	-	-
気管支および肺の悪性新生物	(人)	1,312	-	-	-	-	-
ステージ0		0.5%	-	-	-	-	-
ステージI		16.5%	-	-	-	-	-
ステージII		7.9%	-	-	-	-	-
ステージIII		19.4%	-	-	-	-	-
ステージIV		38.8%	-	-	-	-	-
不明		16.9%	-	-	-	-	-
計		100.0%	-	-	-	-	-
急性心筋梗塞	(人)	370	-	-	-	-	-
Killip I		43.5%	-	-	-	-	-
Killip II		32.7%	-	-	-	-	-
Killip III		8.1%	-	-	-	-	-
Killip IV		12.4%	-	-	-	-	-
不明		3.2%	-	-	-	-	-
計		100.0%	-	-	-	-	-
肺炎	(人)	3,253	-	-	-	-	-
軽症		27.6%	-	-	-	-	-
中等症		8.8%	-	-	-	-	-
重症		33.8%	-	-	-	-	-
不明		29.9%	-	-	-	-	-
計		100.0%	-	-	-	-	-
喘息	(人)	1,141	-	-	-	-	-
軽症間欠型		26.6%	-	-	-	-	-
軽症持続型		36.6%	-	-	-	-	-
中等症持続型		22.6%	-	-	-	-	-
重症持続型		11.9%	-	-	-	-	-
不明		2.2%	-	-	-	-	-
計		100.0%	-	-	-	-	-
脳梗塞	(人)	1,735	-	-	-	-	-
JCS 0-1		70.5%	-	-	-	-	-
JCS 2-3		18.2%	-	-	-	-	-
JCS 10-100		8.8%	-	-	-	-	-

【重症度別集計】表

		平均	病院コード							
			0003	0202	0203	0235	1001	1002	1003	1004
脳梗塞	JCS 200-300	2.5%	4.2%	0.7%	0.0%	1.5%	0.0%	2.8%	6.2%	2.9%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
脳出血	(人)	667	34	161	1	120	4	20	28	22
	JCS 0-1	45.3%	67.6%	39.8%	100.0%	34.2%	75.0%	30.0%	78.6%	59.1%
	JCS 2-3	17.2%	14.7%	24.8%	0.0%	19.2%	0.0%	5.0%	3.6%	4.5%
	JCS 10-100	18.7%	8.8%	19.9%	0.0%	25.8%	0.0%	35.0%	3.6%	18.2%
	JCS 200-300	18.7%	8.8%	15.5%	0.0%	20.8%	25.0%	30.0%	14.3%	18.2%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
糖尿病	(人)	855	72	16	8	82	79	26	34	38
	軽症	62.1%	75.0%	87.5%	37.5%	41.5%	69.6%	61.5%	55.9%	13.2%
	中等症	14.0%	13.9%	6.3%	50.0%	25.6%	20.3%	19.2%	11.8%	0.0%
	重症	10.9%	11.1%	6.3%	12.5%	32.9%	10.1%	19.2%	32.4%	0.0%
	不明	13.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	86.8%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
大腿骨頸部骨折	(人)	648	100	17	1	96	0	3	28	31
	重症度Ⅰ	41.7%	46.0%	41.2%	0.0%	55.2%	0.0%	33.3%	57.1%	0.0%
	重症度Ⅱ	16.5%	16.0%	0.0%	0.0%	12.5%	0.0%	0.0%	3.6%	0.0%
	重症度Ⅲ	24.7%	37.0%	23.5%	0.0%	26.0%	0.0%	33.3%	39.3%	0.0%
	重症度Ⅳ	5.6%	1.0%	35.3%	0.0%	6.3%	0.0%	0.0%	0.0%	0.0%
	不明	11.6%	0.0%	0.0%	100.0%	0.0%	0.0%	33.3%	0.0%	100.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%
胃潰瘍	(人)	252	12	0	2	33	15	7	36	7
	ショックなし(収縮期血圧90mmHg以上)	83.3%	100.0%	0.0%	100.0%	84.8%	93.3%	100.0%	86.1%	57.1%
	ショックあり(収縮期血圧90mmHg未満)	10.3%	0.0%	0.0%	0.0%	15.2%	6.7%	0.0%	13.9%	14.3%
	不明	6.3%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	28.6%
	計	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
急性腸炎	(人)	896	100	2	29	54	30	57	64	13
	血便なし	88.6%	92.0%	100.0%	96.6%	81.5%	76.7%	93.0%	95.3%	15.4%
	血便あり	8.0%	8.0%	0.0%	3.4%	18.5%	23.3%	7.0%	4.7%	0.0%
	不明	3.3%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	84.6%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
急性虫垂炎	(人)	703	58	0	3	91	42	20	56	4
	白血球15000/mm ³ 未満	68.4%	74.1%	0.0%	100.0%	73.6%	71.4%	75.0%	75.0%	75.0%
	白血球15000/mm ³ 以上	28.2%	25.9%	0.0%	0.0%	26.4%	28.6%	25.0%	25.0%	25.0%
	不明	3.4%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
胆石症	(人)	1,413	137	0	9	251	102	45	193	9
	黄疸なし(T Bill 5mg/dl 未満)	92.0%	94.2%	0.0%	100.0%	97.2%	97.1%	97.8%	95.9%	88.9%
	黄疸あり(T Bill 5mg/dl 以上)	5.7%	5.8%	0.0%	0.0%	2.8%	2.9%	2.2%	4.1%	11.1%
	不明	2.3%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
前立腺肥大症	(人)	189	88	0	0	0	7	0	32	0
	前立腺容積20ml未満	25.4%	39.8%	0.0%	0.0%	0.0%	42.9%	0.0%	3.1%	0.0%
	前立腺容積50ml未満	25.9%	17.0%	0.0%	0.0%	0.0%	28.6%	0.0%	46.9%	0.0%
	前立腺容積50ml以上	16.4%	4.5%	0.0%	0.0%	0.0%	14.3%	0.0%	46.9%	0.0%
	不明	32.3%	38.6%	0.0%	0.0%	0.0%	14.3%	0.0%	3.1%	0.0%
	計	100.0%	100.0%	0.0%	0.0%	0.0%	100.0%	0.0%	100.0%	0.0%
白内障	(人)	1,194	163	0	2	2	0	0	0	0
	1眼(処置対象のもの)	58.7%	69.9%	0.0%	100.0%	100.0%	0.0%	0.0%	0.0%	0.0%
	2眼(処置対象のもの)	28.9%	30.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	不明	12.4%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	0.0%	100.0%	100.0%	0.0%	0.0%	0.0%	0.0%
痔核	(人)	140	6	0	2	6	64	5	11	0
	第一度(出血のみ)	27.1%	50.0%	0.0%	0.0%	33.3%	23.4%	20.0%	36.4%	0.0%
	第二度(還納可能)	17.9%	0.0%	0.0%	0.0%	33.3%	4.7%	80.0%	36.4%	0.0%
	第三度(脱肛)	52.1%	50.0%	0.0%	0.0%	33.3%	71.9%	0.0%	27.3%	0.0%
	不明	2.9%	0.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	100.0%	0.0%
子宮筋腫	(人)	476	57	0	0	9	0	0	2	0
	貧血なし(Hb10g/dl以上)	71.2%	42.1%	0.0%	0.0%	55.6%	0.0%	0.0%	100.0%	0.0%
	貧血あり(Hb10g/dl未満)	21.6%	57.9%	0.0%	0.0%	44.4%	0.0%	0.0%	0.0%	0.0%
	不明	7.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	0.0%	0.0%	100.0%	0.0%	0.0%	100.0%	0.0%
狭心症	(人)	2,070	384	44	3	284	327	150	120	4
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
腎結石及び尿管結石	(人)	632	238	0	2	13	78	3	58	2
	水腎・水尿管なし	49.7%	48.3%	0.0%	50.0%	38.5%	39.7%	66.7%	86.2%	50.0%
	水腎・水尿管あり	44.9%	51.7%	0.0%	50.0%	61.5%	59.0%	33.3%	8.6%	0.0%

【重症度別集計】表

		平均	病院コード							
			1009	1012	1013	1015	1018	1019	1025	1026
脳梗塞	JCS 200-300	2.5%	-	1.0%	0.0%	4.8%	-	-	-	-
	不明	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
	計	100.0%	-	100.0%	0.0%	100.0%	-	-	-	-
脳出血	(人)	667	-	49	0	13	-	-	-	-
	JCS 0-1	45.3%	-	36.7%	0.0%	38.5%	-	-	-	-
	JCS 2-3	17.2%	-	20.4%	0.0%	30.8%	-	-	-	-
	JCS 10-100	18.7%	-	10.2%	0.0%	15.4%	-	-	-	-
	JCS 200-300	18.7%	-	32.7%	0.0%	15.4%	-	-	-	-
	不明	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
	計	100.0%	-	100.0%	0.0%	100.0%	-	-	-	-
糖尿病	(人)	855	-	26	2	51	-	-	-	-
	軽症	62.1%	-	61.5%	0.0%	47.1%	-	-	-	-
	中等症	14.0%	-	26.9%	50.0%	21.6%	-	-	-	-
	重症	10.9%	-	11.5%	50.0%	31.4%	-	-	-	-
	不明	13.0%	-	0.0%	0.0%	0.0%	-	-	-	-
	計	100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
大腿骨頸部骨折	(人)	648	-	46	0	79	-	-	-	-
	重症度Ⅰ	41.7%	-	43.5%	0.0%	19.0%	-	-	-	-
	重症度Ⅱ	16.5%	-	30.4%	0.0%	16.5%	-	-	-	-
	重症度Ⅲ	24.7%	-	17.4%	0.0%	50.6%	-	-	-	-
	重症度Ⅳ	5.6%	-	8.7%	0.0%	13.9%	-	-	-	-
	不明	11.6%	-	0.0%	0.0%	0.0%	-	-	-	-
胃潰瘍	(人)	252	-	18	0	12	-	-	-	-
	ショックなし(収縮期血圧90mmHg以上)	83.3%	-	77.8%	0.0%	91.7%	-	-	-	-
	ショックあり(収縮期血圧90mmHg未満)	10.3%	-	22.2%	0.0%	8.3%	-	-	-	-
	不明	6.3%	-	0.0%	0.0%	0.0%	-	-	-	-
急性腸炎	(人)	896	-	59	2	71	-	-	-	-
	血便なし	88.6%	-	93.2%	100.0%	90.1%	-	-	-	-
	血便あり	8.0%	-	6.8%	0.0%	9.9%	-	-	-	-
	不明	3.3%	-	0.0%	0.0%	0.0%	-	-	-	-
急性虫垂炎	(人)	703	-	47	0	21	-	-	-	-
	白血球15000/mm3未満	68.4%	-	70.2%	0.0%	76.2%	-	-	-	-
	白血球15000/mm3以上	28.2%	-	29.8%	0.0%	23.8%	-	-	-	-
	不明	3.4%	-	0.0%	0.0%	0.0%	-	-	-	-
胆石症	(人)	1,413	-	58	0	41	-	-	-	-
	黄疸なし(T Bill 5mg/dl 未満)	92.0%	-	89.7%	0.0%	87.8%	-	-	-	-
	黄疸あり(T Bill 5mg/dl 以上)	5.7%	-	10.3%	0.0%	12.2%	-	-	-	-
	不明	2.3%	-	0.0%	0.0%	0.0%	-	-	-	-
前立腺肥大症	(人)	189	-	0	0	2	-	-	-	-
	前立腺容積20ml未満	25.4%	-	0.0%	0.0%	0.0%	-	-	-	-
	前立腺容積50ml未満	25.9%	-	0.0%	0.0%	50.0%	-	-	-	-
	前立腺容積50ml以上	16.4%	-	0.0%	0.0%	0.0%	-	-	-	-
	不明	32.3%	-	0.0%	0.0%	50.0%	-	-	-	-
白内障	(人)	1,194	-	149	0	121	-	-	-	-
	1眼(処置対象のもの)	58.7%	-	48.3%	0.0%	71.1%	-	-	-	-
	2眼(処置対象のもの)	28.9%	-	51.7%	0.0%	28.9%	-	-	-	-
	不明	12.4%	-	0.0%	0.0%	0.0%	-	-	-	-
痔核	(人)	140	-	9	0	0	-	-	-	-
	第一度(出血のみ)	27.1%	-	44.4%	0.0%	0.0%	-	-	-	-
	第二度(還納可能)	17.9%	-	0.0%	0.0%	0.0%	-	-	-	-
	第三度(脱肛)	52.1%	-	55.6%	0.0%	0.0%	-	-	-	-
	不明	2.9%	-	0.0%	0.0%	0.0%	-	-	-	-
子宮筋腫	(人)	476	-	43	0	0	-	-	-	-
	貧血なし(Hb10g/dl以上)	71.2%	-	74.4%	0.0%	0.0%	-	-	-	-
	貧血あり(Hb10g/dl未満)	21.6%	-	25.6%	0.0%	0.0%	-	-	-	-
	不明	7.1%	-	0.0%	0.0%	0.0%	-	-	-	-
狭心症	(人)	2,070	-	112	0	2	-	-	-	-
	重症度の指標なし	100.0%	-	100.0%	0.0%	100.0%	-	-	-	-
	計	100.0%	-	100.0%	0.0%	100.0%	-	-	-	-
腎結石及び尿管結石	(人)	632	-	2	0	5	-	-	-	-
	水腎・水尿管なし	49.7%	-	100.0%	0.0%	40.0%	-	-	-	-
	水腎・水尿管あり	44.9%	-	0.0%	0.0%	60.0%	-	-	-	-

【重症度別集計】表

		平均	病院コード							
			1039	1041	1046	1047	1049	1058	1069	1129
脳梗塞	JCS 200-300	2.5%	3.7%	-	3.7%	-	4.8%	-	14.3%	-
	不明	N/A	0.0%	-	0.0%	-	0.0%	-	0.0%	-
	計	100.0%	100.0%	-	100.0%	-	100.0%	-	100.0%	-
脳出血	(人)	667	82	-	17	-	39	-	1	-
	JCS 0-1	45.3%	42.7%	-	82.4%	-	51.3%	-	0.0%	-
	JCS 2-3	17.2%	17.1%	-	5.9%	-	7.7%	-	0.0%	-
	JCS 10-100	18.7%	22.0%	-	11.8%	-	12.8%	-	0.0%	-
	JCS 200-300	18.7%	18.3%	-	0.0%	-	28.2%	-	100.0%	-
	不明	N/A	0.0%	-	0.0%	-	0.0%	-	0.0%	-
計	100.0%	100.0%	-	100.0%	-	100.0%	-	100.0%	-	
糖尿病	(人)	855	154	-	9	-	109	-	45	-
	軽症	62.1%	88.3%	-	66.7%	-	84.4%	-	84.4%	-
	中等症	14.0%	9.7%	-	22.2%	-	11.0%	-	8.9%	-
	重症	10.9%	1.3%	-	11.1%	-	4.6%	-	4.4%	-
	不明	13.0%	0.6%	-	0.0%	-	0.0%	-	2.2%	-
計	100.0%	100.0%	-	100.0%	-	100.0%	-	100.0%	-	
大腿骨頸部骨折	(人)	648	79	-	12	-	37	-	0	-
	重症度Ⅰ	41.7%	51.9%	-	33.3%	-	62.2%	-	0.0%	-
	重症度Ⅱ	16.5%	20.3%	-	50.0%	-	13.5%	-	0.0%	-
	重症度Ⅲ	24.7%	22.8%	-	8.3%	-	18.9%	-	0.0%	-
	重症度Ⅳ	5.6%	5.1%	-	8.3%	-	5.4%	-	0.0%	-
	不明	11.6%	0.0%	-	0.0%	-	0.0%	-	0.0%	-
計	100.0%	100.0%	-	100.0%	-	100.0%	-	0.0%	-	
胃潰瘍	(人)	252	11	-	2	-	33	-	21	-
	ショックなし(収縮期血圧90mmHg以上)	83.3%	100.0%	-	100.0%	-	90.9%	-	71.4%	-
	ショックあり(収縮期血圧90mmHg未満)	10.3%	0.0%	-	0.0%	-	9.1%	-	28.6%	-
	不明	6.3%	0.0%	-	0.0%	-	0.0%	-	0.0%	-
計	100.0%	100.0%	-	100.0%	-	100.0%	-	100.0%	-	
急性腸炎	(人)	896	162	-	8	-	103	-	27	-
	血便なし	88.6%	94.4%	-	100.0%	-	88.3%	-	88.9%	-
	血便あり	8.0%	5.6%	-	0.0%	-	11.7%	-	11.1%	-
	不明	3.3%	0.0%	-	0.0%	-	0.0%	-	0.0%	-
計	100.0%	100.0%	-	100.0%	-	100.0%	-	100.0%	-	
急性虫垂炎	(人)	703	139	-	2	-	78	-	32	-
	白血球15000/mm3未満	68.4%	63.3%	-	100.0%	-	79.5%	-	78.1%	-
	白血球15000/mm3以上	28.2%	36.0%	-	0.0%	-	20.5%	-	21.9%	-
	不明	3.4%	0.7%	-	0.0%	-	0.0%	-	0.0%	-
計	100.0%	100.0%	-	100.0%	-	100.0%	-	100.0%	-	
胆石症	(人)	1,413	232	-	11	-	124	-	66	-
	黄疸なし(T Bill 5mg/dl 未満)	92.0%	93.1%	-	72.7%	-	94.4%	-	87.9%	-
	黄疸あり(T Bill 5mg/dl 以上)	5.7%	6.9%	-	27.3%	-	5.6%	-	12.1%	-
	不明	2.3%	0.0%	-	0.0%	-	0.0%	-	0.0%	-
計	100.0%	100.0%	-	100.0%	-	100.0%	-	100.0%	-	
前立腺肥大症	(人)	189	18	-	0	-	8	-	0	-
	前立腺容積20ml未満	25.4%	0.0%	-	0.0%	-	37.5%	-	0.0%	-
	前立腺容積50ml未満	25.9%	27.8%	-	0.0%	-	25.0%	-	0.0%	-
	前立腺容積50ml以上	16.4%	50.0%	-	0.0%	-	25.0%	-	0.0%	-
	不明	32.3%	22.2%	-	0.0%	-	12.5%	-	0.0%	-
計	100.0%	100.0%	-	0.0%	-	100.0%	-	0.0%	-	
白内障	(人)	1,194	229	-	0	-	186	-	0	-
	1眼(処置対象のもの)	58.7%	69.0%	-	0.0%	-	88.7%	-	0.0%	-
	2眼(処置対象のもの)	28.9%	30.6%	-	0.0%	-	11.3%	-	0.0%	-
	不明	12.4%	0.4%	-	0.0%	-	0.0%	-	0.0%	-
計	100.0%	100.0%	-	0.0%	-	100.0%	-	0.0%	-	
痔核	(人)	140	2	-	0	-	32	-	0	-
	第一度(出血のみ)	27.1%	50.0%	-	0.0%	-	21.9%	-	0.0%	-
	第二度(還納可能)	17.9%	0.0%	-	0.0%	-	34.4%	-	0.0%	-
	第三度(脱肛)	52.1%	0.0%	-	0.0%	-	43.8%	-	0.0%	-
	不明	2.9%	50.0%	-	0.0%	-	0.0%	-	0.0%	-
計	100.0%	100.0%	-	0.0%	-	100.0%	-	0.0%	-	
子宮筋腫	(人)	476	37	-	0	-	239	-	0	-
	貧血なし(Hb10g/dl以上)	71.2%	70.3%	-	0.0%	-	85.8%	-	0.0%	-
	貧血あり(Hb10g/dl未満)	21.6%	27.0%	-	0.0%	-	14.2%	-	0.0%	-
	不明	7.1%	2.7%	-	0.0%	-	0.0%	-	0.0%	-
計	100.0%	100.0%	-	0.0%	-	100.0%	-	0.0%	-	
狭心症	(人)	2,070	210	-	1	-	178	-	0	-
	重症度の指標なし	100.0%	100.0%	-	100.0%	-	100.0%	-	0.0%	-
	計	100.0%	100.0%	-	100.0%	-	100.0%	-	0.0%	-
腎結石及び尿管結石	(人)	632	49	-	1	-	114	-	2	-
	水腎・水尿管なし	49.7%	16.3%	-	100.0%	-	55.3%	-	100.0%	-
	水腎・水尿管あり	44.9%	77.6%	-	0.0%	-	44.7%	-	0.0%	-

【重症度別集計】表

		平均	病院コード							
			9009	9010	9011	9012	9013	9014	9015	9016
脳梗塞	JCS 200-300	2.5%	-	2.0%	0.6%	-	-	-	-	-
	不明	N/A	-	0.0%	0.0%	-	-	-	-	-
	計	100.0%	-	100.0%	100.0%	-	-	-	-	-
脳出血	(人)	667	-	20	56	-	-	-	-	-
	JCS 0-1	45.3%	-	35.0%	53.6%	-	-	-	-	-
	JCS 2-3	17.2%	-	30.0%	10.7%	-	-	-	-	-
	JCS 10-100	18.7%	-	15.0%	21.4%	-	-	-	-	-
	JCS 200-300	18.7%	-	20.0%	14.3%	-	-	-	-	-
	不明	N/A	-	0.0%	0.0%	-	-	-	-	-
	計	100.0%	-	100.0%	100.0%	-	-	-	-	-
糖尿病	(人)	855	-	78	26	-	-	-	-	-
	軽症	62.1%	-	2.6%	65.4%	-	-	-	-	-
	中等症	14.0%	-	0.0%	26.9%	-	-	-	-	-
	重症	10.9%	-	0.0%	7.7%	-	-	-	-	-
	不明	13.0%	-	97.4%	0.0%	-	-	-	-	-
	計	100.0%	-	100.0%	100.0%	-	-	-	-	-
大腿骨頸部骨折	(人)	648	-	42	77	-	-	-	-	-
	重症度Ⅰ	41.7%	-	0.0%	57.1%	-	-	-	-	-
	重症度Ⅱ	16.5%	-	0.0%	31.2%	-	-	-	-	-
	重症度Ⅲ	24.7%	-	0.0%	10.4%	-	-	-	-	-
	重症度Ⅳ	5.6%	-	0.0%	1.3%	-	-	-	-	-
	不明	11.6%	-	100.0%	0.0%	-	-	-	-	-
計	100.0%	-	100.0%	100.0%	-	-	-	-	-	
胃潰瘍	(人)	252	-	11	32	-	-	-	-	-
	ショックなし(収縮期血圧90mmHg以上)	83.3%	-	0.0%	90.6%	-	-	-	-	-
	ショックあり(収縮期血圧90mmHg未満)	10.3%	-	0.0%	0.0%	-	-	-	-	-
	不明	6.3%	-	100.0%	9.4%	-	-	-	-	-
計	100.0%	-	100.0%	100.0%	-	-	-	-	-	
急性腸炎	(人)	896	-	19	96	-	-	-	-	-
	血便なし	88.6%	-	10.5%	93.8%	-	-	-	-	-
	血便あり	8.0%	-	0.0%	4.2%	-	-	-	-	-
	不明	3.3%	-	89.5%	2.1%	-	-	-	-	-
計	100.0%	-	100.0%	100.0%	-	-	-	-	-	
急性虫垂炎	(人)	703	-	23	87	-	-	-	-	-
	白血球15000/mm3未満	68.4%	-	0.0%	59.8%	-	-	-	-	-
	白血球15000/mm3以上	28.2%	-	0.0%	40.2%	-	-	-	-	-
	不明	3.4%	-	100.0%	0.0%	-	-	-	-	-
計	100.0%	-	100.0%	100.0%	-	-	-	-	-	
胆石症	(人)	1,413	-	33	102	-	-	-	-	-
	黄疸なし(T Bill 5mg/dl 未満)	92.0%	-	3.0%	92.2%	-	-	-	-	-
	黄疸あり(T Bill 5mg/dl 以上)	5.7%	-	0.0%	7.8%	-	-	-	-	-
	不明	2.3%	-	97.0%	0.0%	-	-	-	-	-
計	100.0%	-	100.0%	100.0%	-	-	-	-	-	
前立腺肥大症	(人)	189	-	6	28	-	-	-	-	-
	前立腺容積20ml未満	25.4%	-	0.0%	21.4%	-	-	-	-	-
	前立腺容積50ml未満	25.9%	-	0.0%	32.1%	-	-	-	-	-
	前立腺容積50ml以上	16.4%	-	0.0%	0.0%	-	-	-	-	-
	不明	32.3%	-	100.0%	46.4%	-	-	-	-	-
計	100.0%	-	100.0%	100.0%	-	-	-	-	-	
白内障	(人)	1,194	-	149	193	-	-	-	-	-
	1眼(処置対象のもの)	58.7%	-	1.3%	51.8%	-	-	-	-	-
	2眼(処置対象のもの)	28.9%	-	0.0%	48.2%	-	-	-	-	-
	不明	12.4%	-	98.7%	0.0%	-	-	-	-	-
計	100.0%	-	100.0%	100.0%	-	-	-	-	-	
痔核	(人)	140	-	1	2	-	-	-	-	-
	第一度(出血のみ)	27.1%	-	0.0%	50.0%	-	-	-	-	-
	第二度(還納可能)	17.9%	-	0.0%	50.0%	-	-	-	-	-
	第三度(脱肛)	52.1%	-	0.0%	0.0%	-	-	-	-	-
	不明	2.9%	-	100.0%	0.0%	-	-	-	-	-
計	100.0%	-	100.0%	100.0%	-	-	-	-	-	
子宮筋腫	(人)	476	-	33	56	-	-	-	-	-
	貧血なし(Hb10g/dl以上)	71.2%	-	0.0%	80.4%	-	-	-	-	-
	貧血あり(Hb10g/dl未満)	21.6%	-	0.0%	19.6%	-	-	-	-	-
	不明	7.1%	-	100.0%	0.0%	-	-	-	-	-
計	100.0%	-	100.0%	100.0%	-	-	-	-	-	
狭心症	(人)	2,070	-	245	6	-	-	-	-	-
	重症度の指標なし	100.0%	-	100.0%	100.0%	-	-	-	-	-
計	100.0%	-	100.0%	100.0%	-	-	-	-	-	
腎結石及び尿管結石	(人)	632	-	26	39	-	-	-	-	-
	水腎・水尿管なし	49.7%	-	0.0%	79.5%	-	-	-	-	-
	水腎・水尿管あり	44.9%	-	0.0%	20.5%	-	-	-	-	-

【重症度別集計】表

		平均	病院コード				
			9017	9018	9019	9020	9021
脳梗塞	JCS 200-300	2.5%	-	-	-	-	-
	不明	N/A	-	-	-	-	-
	計	100.0%	-	-	-	-	-
脳出血	(人)	667	-	-	-	-	-
	JCS 0-1	45.3%	-	-	-	-	-
	JCS 2-3	17.2%	-	-	-	-	-
	JCS 10-100	18.7%	-	-	-	-	-
	JCS 200-300	18.7%	-	-	-	-	-
	不明	N/A	-	-	-	-	-
	計	100.0%	-	-	-	-	-
糖尿病	(人)	855	-	-	-	-	-
	軽症	62.1%	-	-	-	-	-
	中等症	14.0%	-	-	-	-	-
	重症	10.9%	-	-	-	-	-
	不明	13.0%	-	-	-	-	-
	計	100.0%	-	-	-	-	-
大腿骨頸部骨折	(人)	648	-	-	-	-	-
	重症度Ⅰ	41.7%	-	-	-	-	-
	重症度Ⅱ	16.5%	-	-	-	-	-
	重症度Ⅲ	24.7%	-	-	-	-	-
	重症度Ⅳ	5.6%	-	-	-	-	-
	不明	11.6%	-	-	-	-	-
計	100.0%	-	-	-	-	-	
胃潰瘍	(人)	252	-	-	-	-	-
	ショックなし（収縮期血圧90mmHg以上）	83.3%	-	-	-	-	-
	ショックあり（収縮期血圧90mmHg未満）	10.3%	-	-	-	-	-
	不明	6.3%	-	-	-	-	-
計	100.0%	-	-	-	-	-	
急性腸炎	(人)	896	-	-	-	-	-
	血便なし	88.6%	-	-	-	-	-
	血便あり	8.0%	-	-	-	-	-
	不明	3.3%	-	-	-	-	-
計	100.0%	-	-	-	-	-	
急性虫垂炎	(人)	703	-	-	-	-	-
	白血球15000/mm3未満	68.4%	-	-	-	-	-
	白血球15000/mm3以上	28.2%	-	-	-	-	-
	不明	3.4%	-	-	-	-	-
計	100.0%	-	-	-	-	-	
胆石症	(人)	1,413	-	-	-	-	-
	黄疸なし(T Bill 5mg/dl 未満)	92.0%	-	-	-	-	-
	黄疸あり(T Bill 5mg/dl 以上)	5.7%	-	-	-	-	-
	不明	2.3%	-	-	-	-	-
計	100.0%	-	-	-	-	-	
前立腺肥大症	(人)	189	-	-	-	-	-
	前立腺容積20ml未満	25.4%	-	-	-	-	-
	前立腺容積50ml未満	25.9%	-	-	-	-	-
	前立腺容積50ml以上	16.4%	-	-	-	-	-
	不明	32.3%	-	-	-	-	-
計	100.0%	-	-	-	-	-	
白内障	(人)	1,194	-	-	-	-	-
	1眼(処置対象のもの)	58.7%	-	-	-	-	-
	2眼(処置対象のもの)	28.9%	-	-	-	-	-
	不明	12.4%	-	-	-	-	-
計	100.0%	-	-	-	-	-	
痔核	(人)	140	-	-	-	-	-
	第一度(出血のみ)	27.1%	-	-	-	-	-
	第二度(還納可能)	17.9%	-	-	-	-	-
	第三度(脱肛)	52.1%	-	-	-	-	-
	不明	2.9%	-	-	-	-	-
計	100.0%	-	-	-	-	-	
子宮筋腫	(人)	476	-	-	-	-	-
	貧血なし(Hb10g/dl以上)	71.2%	-	-	-	-	-
	貧血あり(Hb10g/dl未満)	21.6%	-	-	-	-	-
	不明	7.1%	-	-	-	-	-
計	100.0%	-	-	-	-	-	
狭心症	(人)	2,070	-	-	-	-	-
	重症度の指標なし	100.0%	-	-	-	-	-
	計	100.0%	-	-	-	-	-
腎結石及び尿管結石	(人)	632	-	-	-	-	-
	水腎・水尿管なし	49.7%	-	-	-	-	-
	水腎・水尿管あり	44.9%	-	-	-	-	-

【重症度別集計】表

		平均	病院コード				
			9017	9018	9019	9020	9021
腎結石及び尿管結石	不明	5.4%	-	-	-	-	-
	計	100.0%	-	-	-	-	-
乳房の悪性新生物	(人)	576	-	-	-	-	-
	ステージ0	4.9%	-	-	-	-	
	ステージI	33.7%	-	-	-	-	
	ステージII	32.1%	-	-	-	-	
	ステージIII	10.4%	-	-	-	-	
	ステージIV	10.1%	-	-	-	-	
	不明	8.9%	-	-	-	-	
計	100.0%	-	-	-	-		
膝関節症	(人)	386	-	-	-	-	
	骨移植なし	90.7%	-	-	-	-	
	骨移植あり	1.0%	-	-	-	-	
	不明	8.3%	-	-	-	-	
計	100.0%	-	-	-	-		
鼠径ヘルニア	(人)	912	-	-	-	-	
	重症度の指標なし	100.0%	-	-	-	-	
計	100.0%	-	-	-	-		
全疾患合計		100.0%	-	-	-	-	
集計病院数		17					