

【重症度別集計】表

絞込条件

対象年月設定: 201404 ~ 201503
 病院の経営主体選択: 全て
 【詳細】病床数: 下限なし ~ 上限なし
 【詳細】手術の有無: 全て
 【詳細】ステージ集約選択: 有り
 【詳細】年代: 全て
 【詳細】性別: 全て
 【詳細】入院経路: 全て
 並べ替え条件: 病院コード・昇順

■ 退院患者数

	(単位:%)	平均	病院コード							
			0003	0202	0203	0235	1001	1002	1003	1004
胃の悪性新生物	(人)	696	73	1	2	130	61	44	190	6
ステージ0		2.3%	1.4%	0.0%	50.0%	1.5%	9.8%	0.0%	0.0%	0.0%
ステージI		28.3%	28.8%	0.0%	0.0%	27.7%	19.7%	18.2%	29.5%	0.0%
ステージII		14.1%	5.5%	0.0%	0.0%	7.7%	19.7%	13.6%	17.4%	0.0%
ステージIII		11.6%	13.7%	0.0%	0.0%	15.4%	14.8%	18.2%	11.6%	0.0%
ステージIV		37.2%	38.4%	0.0%	0.0%	44.6%	34.4%	50.0%	34.7%	0.0%
不明		6.5%	12.3%	100.0%	50.0%	3.1%	1.6%	0.0%	6.8%	100.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
結腸の悪性新生物	(人)	866	89	0	13	119	258	20	157	5
ステージ0		3.7%	4.5%	0.0%	0.0%	4.2%	0.8%	0.0%	1.9%	0.0%
ステージI		8.5%	15.7%	0.0%	0.0%	10.1%	3.5%	5.0%	8.9%	0.0%
ステージII		22.7%	23.6%	0.0%	7.7%	21.8%	26.0%	35.0%	21.7%	0.0%
ステージIII		22.3%	23.6%	0.0%	7.7%	17.6%	26.4%	20.0%	20.4%	0.0%
ステージIV		34.6%	15.7%	0.0%	0.0%	44.5%	39.9%	35.0%	41.4%	20.0%
不明		8.1%	16.9%	0.0%	84.6%	1.7%	3.5%	5.0%	5.7%	80.0%
計		100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
直腸の悪性新生物	(人)	453	20	0	1	90	96	8	71	3
ステージ0		3.1%	5.0%	0.0%	0.0%	3.3%	2.1%	12.5%	5.6%	0.0%
ステージI		10.6%	15.0%	0.0%	0.0%	13.3%	3.1%	0.0%	9.9%	0.0%
ステージII		16.6%	10.0%	0.0%	0.0%	22.2%	11.5%	12.5%	38.0%	0.0%
ステージIII		32.7%	50.0%	0.0%	0.0%	23.3%	46.9%	25.0%	22.5%	0.0%
ステージIV		33.8%	5.0%	0.0%	0.0%	37.8%	36.5%	37.5%	23.9%	0.0%
不明		3.3%	15.0%	0.0%	100.0%	0.0%	0.0%	12.5%	0.0%	100.0%
計		100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
気管支および肺の悪性新生物	(人)	783	28	1	10	127	38	27	257	15
ステージ0		0.4%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.8%	0.0%
ステージI		15.1%	0.0%	0.0%	0.0%	26.8%	18.4%	33.3%	10.5%	0.0%
ステージII		8.2%	0.0%	0.0%	0.0%	3.9%	18.4%	7.4%	5.8%	0.0%
ステージIII		21.7%	14.3%	0.0%	0.0%	15.0%	7.9%	14.8%	30.4%	0.0%
ステージIV		38.2%	64.3%	0.0%	20.0%	37.0%	50.0%	18.5%	35.0%	0.0%
不明		16.5%	21.4%	100.0%	80.0%	17.3%	5.3%	25.9%	17.5%	100.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
急性心筋梗塞	(人)	266	15	1	0	63	39	7	54	0
Killip I		41.7%	66.7%	0.0%	0.0%	47.6%	38.5%	0.0%	31.5%	0.0%
Killip II		27.8%	26.7%	0.0%	0.0%	28.6%	41.0%	42.9%	33.3%	0.0%
Killip III		11.7%	6.7%	0.0%	0.0%	6.3%	5.1%	42.9%	13.0%	0.0%
Killip IV		16.9%	0.0%	100.0%	0.0%	14.3%	15.4%	14.3%	16.7%	0.0%
不明		1.9%	0.0%	0.0%	0.0%	3.2%	0.0%	0.0%	5.6%	0.0%
計		100.0%	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	0.0%
肺炎	(人)	2,889	176	33	165	235	96	199	226	104
軽症		23.6%	40.9%	27.3%	56.4%	20.4%	44.8%	27.1%	10.2%	37.5%
中等症		7.9%	19.9%	18.2%	10.9%	7.2%	14.6%	15.1%	4.0%	26.0%
重症		32.9%	38.6%	54.5%	32.7%	72.3%	40.6%	57.8%	37.2%	25.0%
不明		35.5%	0.6%	0.0%	0.0%	0.0%	0.0%	0.0%	48.7%	11.5%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
喘息	(人)	916	12	1	28	73	0	12	82	2
軽症間欠型		30.2%	50.0%	100.0%	28.6%	12.3%	0.0%	41.7%	6.1%	0.0%
軽症持続型		36.2%	33.3%	0.0%	25.0%	38.4%	0.0%	58.3%	0.0%	0.0%
中等症持続型		20.1%	16.7%	0.0%	46.4%	46.6%	0.0%	0.0%	59.8%	0.0%
重症持続型		12.2%	0.0%	0.0%	0.0%	1.4%	0.0%	0.0%	32.9%	0.0%
不明		1.2%	0.0%	0.0%	0.0%	1.4%	0.0%	0.0%	1.2%	100.0%
計		100.0%	100.0%	100.0%	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%
脳梗塞	(人)	1,365	111	453	5	190	6	74	95	34
JCS 0-1		67.4%	71.2%	54.7%	40.0%	68.9%	83.3%	66.2%	81.1%	73.5%
JCS 2-3		20.8%	17.1%	37.3%	0.0%	18.9%	0.0%	8.1%	5.3%	0.0%
JCS 10-100		9.2%	9.9%	6.8%	40.0%	9.5%	0.0%	14.9%	10.5%	23.5%

【重症度別集計】表

	(単位:%)	平均	病院コード							
			1009	1012	1013	1015	1018	1019	1025	1039
胃の悪性新生物	(人)	696	-	49	0	13	-	-	-	78
ステージ0		2.3%	-	2.0%	0.0%	15.4%	-	-	-	0.0%
ステージⅠ		28.3%	-	36.7%	0.0%	7.7%	-	-	-	46.2%
ステージⅡ		14.1%	-	22.4%	0.0%	7.7%	-	-	-	9.0%
ステージⅢ		11.6%	-	4.1%	0.0%	0.0%	-	-	-	10.3%
ステージⅣ		37.2%	-	30.6%	0.0%	46.2%	-	-	-	32.1%
不明		6.5%	-	4.1%	0.0%	23.1%	-	-	-	2.6%
計		100.0%	-	100.0%	0.0%	100.0%	-	-	-	100.0%
結腸の悪性新生物	(人)	866	-	50	0	15	-	-	-	94
ステージ0		3.7%	-	12.0%	0.0%	0.0%	-	-	-	6.4%
ステージⅠ		8.5%	-	18.0%	0.0%	6.7%	-	-	-	10.6%
ステージⅡ		22.7%	-	24.0%	0.0%	13.3%	-	-	-	22.3%
ステージⅢ		22.3%	-	24.0%	0.0%	13.3%	-	-	-	22.3%
ステージⅣ		34.6%	-	18.0%	0.0%	40.0%	-	-	-	25.5%
不明		8.1%	-	4.0%	0.0%	26.7%	-	-	-	12.8%
計		100.0%	-	100.0%	0.0%	100.0%	-	-	-	100.0%
直腸の悪性新生物	(人)	453	-	21	0	16	-	-	-	66
ステージ0		3.1%	-	9.5%	0.0%	0.0%	-	-	-	0.0%
ステージⅠ		10.6%	-	19.0%	0.0%	6.3%	-	-	-	19.7%
ステージⅡ		16.6%	-	9.5%	0.0%	50.0%	-	-	-	6.1%
ステージⅢ		32.7%	-	0.0%	0.0%	18.8%	-	-	-	50.0%
ステージⅣ		33.8%	-	57.1%	0.0%	6.3%	-	-	-	22.7%
不明		3.3%	-	4.8%	0.0%	18.8%	-	-	-	1.5%
計		100.0%	-	100.0%	0.0%	100.0%	-	-	-	100.0%
気管支および肺の悪性新生物	(人)	783	-	15	157	5	-	-	-	95
ステージ0		0.4%	-	0.0%	0.6%	0.0%	-	-	-	0.0%
ステージⅠ		15.1%	-	6.7%	19.1%	0.0%	-	-	-	9.5%
ステージⅡ		8.2%	-	6.7%	19.7%	0.0%	-	-	-	3.2%
ステージⅢ		21.7%	-	0.0%	31.2%	0.0%	-	-	-	13.7%
ステージⅣ		38.2%	-	66.7%	29.3%	40.0%	-	-	-	60.0%
不明		16.5%	-	20.0%	0.0%	60.0%	-	-	-	13.7%
計		100.0%	-	100.0%	100.0%	100.0%	-	-	-	100.0%
急性心筋梗塞	(人)	266	-	42	0	3	-	-	-	39
KillipⅠ		41.7%	-	42.9%	0.0%	33.3%	-	-	-	51.3%
KillipⅡ		27.8%	-	16.7%	0.0%	33.3%	-	-	-	15.4%
KillipⅢ		11.7%	-	23.8%	0.0%	0.0%	-	-	-	10.3%
KillipⅣ		16.9%	-	16.7%	0.0%	33.3%	-	-	-	23.1%
不明		1.9%	-	0.0%	0.0%	0.0%	-	-	-	0.0%
計		100.0%	-	100.0%	0.0%	100.0%	-	-	-	100.0%
肺炎	(人)	2,889	-	236	73	311	-	-	-	849
軽症		23.6%	-	54.2%	1.4%	23.8%	-	-	-	7.2%
中等症		7.9%	-	10.2%	0.0%	4.2%	-	-	-	0.4%
重症		32.9%	-	35.6%	98.6%	17.7%	-	-	-	6.0%
不明		35.5%	-	0.0%	0.0%	54.3%	-	-	-	86.5%
計		100.0%	-	100.0%	100.0%	100.0%	-	-	-	100.0%
喘息	(人)	916	-	50	15	39	-	-	-	578
軽症間欠型		30.2%	-	54.0%	20.0%	82.1%	-	-	-	30.6%
軽症持続型		36.2%	-	20.0%	20.0%	12.8%	-	-	-	45.0%
中等症持続型		20.1%	-	18.0%	40.0%	2.6%	-	-	-	10.7%
重症持続型		12.2%	-	8.0%	20.0%	2.6%	-	-	-	12.8%
不明		1.2%	-	0.0%	0.0%	0.0%	-	-	-	0.9%
計		100.0%	-	100.0%	100.0%	100.0%	-	-	-	100.0%
脳梗塞	(人)	1,365	-	100	0	53	-	-	-	182
JCS 0-1		67.4%	-	75.0%	0.0%	56.6%	-	-	-	84.1%
JCS 2-3		20.8%	-	15.0%	0.0%	26.4%	-	-	-	8.2%
JCS 10-100		9.2%	-	7.0%	0.0%	11.3%	-	-	-	6.6%

【重症度別集計】表

	(単位:%)	平均	病院コード							
			1041	1046	1047	1049	1058	1069	1129	1133
胃の悪性新生物	(人)	696	-	4	-	-	-	45	-	-
ステージ0		2.3%	-	0.0%	-	-	-	6.7%	-	-
ステージI		28.3%	-	0.0%	-	-	-	20.0%	-	-
ステージII		14.1%	-	0.0%	-	-	-	31.1%	-	-
ステージIII		11.6%	-	25.0%	-	-	-	2.2%	-	-
ステージIV		37.2%	-	25.0%	-	-	-	37.8%	-	-
不明		6.5%	-	50.0%	-	-	-	2.2%	-	-
計		100.0%	-	100.0%	-	-	-	100.0%	-	-
結腸の悪性新生物	(人)	866	-	2	-	-	-	44	-	-
ステージ0		3.7%	-	0.0%	-	-	-	13.6%	-	-
ステージI		8.5%	-	0.0%	-	-	-	9.1%	-	-
ステージII		22.7%	-	0.0%	-	-	-	13.6%	-	-
ステージIII		22.3%	-	50.0%	-	-	-	22.7%	-	-
ステージIV		34.6%	-	0.0%	-	-	-	40.9%	-	-
不明		8.1%	-	50.0%	-	-	-	0.0%	-	-
計		100.0%	-	100.0%	-	-	-	100.0%	-	-
直腸の悪性新生物	(人)	453	-	1	-	-	-	60	-	-
ステージ0		3.1%	-	0.0%	-	-	-	1.7%	-	-
ステージI		10.6%	-	0.0%	-	-	-	8.3%	-	-
ステージII		16.6%	-	0.0%	-	-	-	0.0%	-	-
ステージIII		32.7%	-	0.0%	-	-	-	30.0%	-	-
ステージIV		33.8%	-	100.0%	-	-	-	56.7%	-	-
不明		3.3%	-	0.0%	-	-	-	3.3%	-	-
計		100.0%	-	100.0%	-	-	-	100.0%	-	-
気管支および肺の悪性新生物	(人)	783	-	4	-	-	-	4	-	-
ステージ0		0.4%	-	0.0%	-	-	-	0.0%	-	-
ステージI		15.1%	-	0.0%	-	-	-	25.0%	-	-
ステージII		8.2%	-	0.0%	-	-	-	0.0%	-	-
ステージIII		21.7%	-	0.0%	-	-	-	0.0%	-	-
ステージIV		38.2%	-	25.0%	-	-	-	50.0%	-	-
不明		16.5%	-	75.0%	-	-	-	25.0%	-	-
計		100.0%	-	100.0%	-	-	-	100.0%	-	-
急性心筋梗塞	(人)	266	-	3	-	-	-	0	-	-
Killip I		41.7%	-	0.0%	-	-	-	0.0%	-	-
Killip II		27.8%	-	33.3%	-	-	-	0.0%	-	-
Killip III		11.7%	-	0.0%	-	-	-	0.0%	-	-
Killip IV		16.9%	-	66.7%	-	-	-	0.0%	-	-
不明		1.9%	-	0.0%	-	-	-	0.0%	-	-
計		100.0%	-	100.0%	-	-	-	0.0%	-	-
肺炎	(人)	2,889	-	73	-	-	-	113	-	-
軽症		23.6%	-	31.5%	-	-	-	13.3%	-	-
中等症		7.9%	-	9.6%	-	-	-	23.0%	-	-
重症		32.9%	-	58.9%	-	-	-	63.7%	-	-
不明		35.5%	-	0.0%	-	-	-	0.0%	-	-
計		100.0%	-	100.0%	-	-	-	100.0%	-	-
喘息	(人)	916	-	2	-	-	-	22	-	-
軽症間欠型		30.2%	-	100.0%	-	-	-	9.1%	-	-
軽症持続型		36.2%	-	0.0%	-	-	-	36.4%	-	-
中等症持続型		20.1%	-	0.0%	-	-	-	36.4%	-	-
重症持続型		12.2%	-	0.0%	-	-	-	9.1%	-	-
不明		1.2%	-	0.0%	-	-	-	9.1%	-	-
計		100.0%	-	100.0%	-	-	-	100.0%	-	-
脳梗塞	(人)	1,365	-	56	-	-	-	6	-	-
JCS 0-1		67.4%	-	80.4%	-	-	-	16.7%	-	-
JCS 2-3		20.8%	-	7.1%	-	-	-	16.7%	-	-
JCS 10-100		9.2%	-	12.5%	-	-	-	33.3%	-	-

【重症度別集計】表

	(単位:%)	平均	病院コード		
			9019	9020	9021
胃の悪性新生物	(人)	696	-	-	-
ステージ0		2.3%	-	-	-
ステージI		28.3%	-	-	-
ステージII		14.1%	-	-	-
ステージIII		11.6%	-	-	-
ステージIV		37.2%	-	-	-
不明		6.5%	-	-	-
計		100.0%	-	-	-
結腸の悪性新生物	(人)	866	-	-	-
ステージ0		3.7%	-	-	-
ステージI		8.5%	-	-	-
ステージII		22.7%	-	-	-
ステージIII		22.3%	-	-	-
ステージIV		34.6%	-	-	-
不明		8.1%	-	-	-
計		100.0%	-	-	-
直腸の悪性新生物	(人)	453	-	-	-
ステージ0		3.1%	-	-	-
ステージI		10.6%	-	-	-
ステージII		16.6%	-	-	-
ステージIII		32.7%	-	-	-
ステージIV		33.8%	-	-	-
不明		3.3%	-	-	-
計		100.0%	-	-	-
気管支および肺の悪性新生物	(人)	783	-	-	-
ステージ0		0.4%	-	-	-
ステージI		15.1%	-	-	-
ステージII		8.2%	-	-	-
ステージIII		21.7%	-	-	-
ステージIV		38.2%	-	-	-
不明		16.5%	-	-	-
計		100.0%	-	-	-
急性心筋梗塞	(人)	266	-	-	-
Killip I		41.7%	-	-	-
Killip II		27.8%	-	-	-
Killip III		11.7%	-	-	-
Killip IV		16.9%	-	-	-
不明		1.9%	-	-	-
計		100.0%	-	-	-
肺炎	(人)	2,889	-	-	-
軽症		23.6%	-	-	-
中等症		7.9%	-	-	-
重症		32.9%	-	-	-
不明		35.5%	-	-	-
計		100.0%	-	-	-
喘息	(人)	916	-	-	-
軽症間欠型		30.2%	-	-	-
軽症持続型		36.2%	-	-	-
中等症持続型		20.1%	-	-	-
重症持続型		12.2%	-	-	-
不明		1.2%	-	-	-
計		100.0%	-	-	-
脳梗塞	(人)	1,365	-	-	-
JCS 0-1		67.4%	-	-	-
JCS 2-3		20.8%	-	-	-
JCS 10-100		9.2%	-	-	-

【重症度別集計】表

(単位:%)		平均	病院コード							
			0003	0202	0203	0235	1001	1002	1003	1004
脳梗塞	JCS 200-300	2.6%	1.8%	1.1%	20.0%	2.6%	16.7%	10.8%	3.2%	2.9%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
脳出血	(人)	586	34	169	2	135	2	24	36	7
	JCS 0-1	46.1%	47.1%	37.9%	0.0%	45.2%	50.0%	25.0%	72.2%	85.7%
	JCS 2-3	19.3%	26.5%	26.6%	0.0%	19.3%	50.0%	12.5%	8.3%	0.0%
	JCS 10-100	19.1%	14.7%	18.9%	0.0%	22.2%	0.0%	33.3%	0.0%	0.0%
	JCS 200-300	15.5%	11.8%	16.6%	100.0%	13.3%	0.0%	29.2%	19.4%	14.3%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
糖尿病	(人)	585	78	12	9	60	66	15	39	30
	軽症	65.3%	74.4%	50.0%	66.7%	66.7%	77.3%	40.0%	48.7%	0.0%
	中等症	17.9%	12.8%	50.0%	11.1%	18.3%	15.2%	53.3%	23.1%	0.0%
	重症	11.1%	11.5%	0.0%	22.2%	15.0%	7.6%	6.7%	28.2%	0.0%
	不明	5.6%	1.3%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
大腿骨頸部骨折	(人)	511	140	0	1	114	3	2	36	21
	重症度Ⅰ	41.5%	55.0%	0.0%	0.0%	50.0%	33.3%	0.0%	44.4%	0.0%
	重症度Ⅱ	19.8%	11.4%	0.0%	0.0%	21.1%	66.7%	0.0%	22.2%	0.0%
	重症度Ⅲ	27.4%	29.3%	0.0%	0.0%	24.6%	0.0%	0.0%	27.8%	0.0%
	重症度Ⅳ	5.9%	3.6%	0.0%	0.0%	4.4%	0.0%	0.0%	5.6%	0.0%
	不明	5.5%	0.7%	0.0%	100.0%	0.0%	0.0%	100.0%	0.0%	100.0%
計	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
胃潰瘍	(人)	202	35	0	6	31	9	8	44	5
	ショックなし(収縮期血圧90mmHg以上)	87.6%	88.6%	0.0%	100.0%	90.3%	88.9%	100.0%	86.4%	80.0%
	ショックあり(収縮期血圧90mmHg未満)	11.9%	11.4%	0.0%	0.0%	9.7%	11.1%	0.0%	13.6%	0.0%
	不明	0.5%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	20.0%
計	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
急性腸炎	(人)	701	92	2	15	51	18	51	60	15
	血便なし	92.3%	95.7%	100.0%	86.7%	90.2%	83.3%	86.3%	93.3%	53.3%
	血便あり	6.6%	4.3%	0.0%	13.3%	9.8%	16.7%	13.7%	6.7%	0.0%
	不明	1.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	46.7%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
急性虫垂炎	(人)	522	78	0	2	109	30	28	66	1
	白血球15000/mm3未満	67.4%	67.9%	0.0%	50.0%	72.5%	66.7%	64.3%	71.2%	100.0%
	白血球15000/mm3以上	32.2%	32.1%	0.0%	0.0%	27.5%	33.3%	35.7%	28.8%	0.0%
	不明	0.4%	0.0%	0.0%	50.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
胆石症	(人)	1,108	120	1	17	221	110	46	225	11
	黄疸なし(T Bill 5mg/dl 未満)	93.1%	90.8%	100.0%	94.1%	97.3%	87.3%	95.7%	93.8%	100.0%
	黄疸あり(T Bill 5mg/dl 以上)	6.9%	9.2%	0.0%	5.9%	2.7%	12.7%	4.3%	6.2%	0.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
前立腺肥大症	(人)	95	33	0	0	0	10	2	36	0
	前立腺容積20ml未満	20.0%	48.5%	0.0%	0.0%	0.0%	10.0%	0.0%	2.8%	0.0%
	前立腺容積50ml未満	40.0%	39.4%	0.0%	0.0%	0.0%	50.0%	0.0%	38.9%	0.0%
	前立腺容積50ml以上	30.5%	9.1%	0.0%	0.0%	0.0%	10.0%	0.0%	58.3%	0.0%
	不明	9.5%	3.0%	0.0%	0.0%	0.0%	30.0%	100.0%	0.0%	0.0%
計	100.0%	100.0%	0.0%	0.0%	0.0%	100.0%	100.0%	100.0%	0.0%	
白内障	(人)	679	274	0	2	0	0	0	0	0
	1眼(処置対象のもの)	91.2%	81.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	2眼(処置対象のもの)	8.8%	19.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	0.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	
痔核	(人)	74	3	0	2	5	42	5	11	0
	第一度(出血のみ)	24.3%	100.0%	0.0%	100.0%	60.0%	9.5%	0.0%	45.5%	0.0%
	第二度(還納可能)	12.2%	0.0%	0.0%	0.0%	20.0%	11.9%	40.0%	9.1%	0.0%
	第三度(脱肛)	62.2%	0.0%	0.0%	0.0%	20.0%	76.2%	60.0%	45.5%	0.0%
	不明	1.4%	0.0%	0.0%	0.0%	0.0%	2.4%	0.0%	0.0%	0.0%
計	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	100.0%	0.0%	
子宮筋腫	(人)	142	51	0	0	0	0	0	4	0
	貧血なし(Hb10g/dl以上)	65.5%	70.6%	0.0%	0.0%	0.0%	0.0%	0.0%	50.0%	0.0%
	貧血あり(Hb10g/dl未満)	33.8%	29.4%	0.0%	0.0%	0.0%	0.0%	0.0%	50.0%	0.0%
	不明	0.7%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
計	100.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	
狭心症	(人)	1,963	264	29	2	356	178	170	535	7
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	
腎結石及び尿管結石	(人)	487	194	0	1	10	88	0	82	0
	水腎・水尿管なし	35.7%	26.3%	0.0%	0.0%	70.0%	44.3%	0.0%	45.1%	0.0%
	水腎・水尿管あり	57.5%	73.7%	0.0%	100.0%	30.0%	53.4%	0.0%	41.5%	0.0%

【重症度別集計】表

		平均	病院コード							
			1009	1012	1013	1015	1018	1019	1025	1039
脳梗塞	JCS 200-300	2.6%	-	3.0%	0.0%	5.7%	-	-	-	1.1%
	不明	N/A	-	0.0%	0.0%	0.0%	-	-	-	0.0%
	計	100.0%	-	100.0%	0.0%	100.0%	-	-	-	100.0%
脳出血	(人)	586	-	60	0	8	-	-	-	91
	JCS 0-1	46.1%	-	46.7%	0.0%	25.0%	-	-	-	50.5%
	JCS 2-3	19.3%	-	11.7%	0.0%	50.0%	-	-	-	15.4%
	JCS 10-100	19.1%	-	21.7%	0.0%	25.0%	-	-	-	20.9%
	JCS 200-300	15.5%	-	20.0%	0.0%	0.0%	-	-	-	13.2%
	不明	N/A	-	0.0%	0.0%	0.0%	-	-	-	0.0%
	計	100.0%	-	100.0%	0.0%	100.0%	-	-	-	100.0%
糖尿病	(人)	585	-	24	1	64	-	-	-	135
	軽症	65.3%	-	66.7%	0.0%	40.6%	-	-	-	88.9%
	中等症	17.9%	-	25.0%	0.0%	35.9%	-	-	-	8.9%
	重症	11.1%	-	8.3%	100.0%	23.4%	-	-	-	1.5%
	不明	5.6%	-	0.0%	0.0%	0.0%	-	-	-	0.7%
	計	100.0%	-	100.0%	100.0%	100.0%	-	-	-	100.0%
大腿骨頸部骨折	(人)	511	-	29	0	77	-	-	-	62
	重症度Ⅰ	41.5%	-	31.0%	0.0%	28.6%	-	-	-	43.5%
	重症度Ⅱ	19.8%	-	20.7%	0.0%	19.5%	-	-	-	25.8%
	重症度Ⅲ	27.4%	-	37.9%	0.0%	31.2%	-	-	-	30.6%
	重症度Ⅳ	5.9%	-	10.3%	0.0%	19.5%	-	-	-	0.0%
	不明	5.5%	-	0.0%	0.0%	1.3%	-	-	-	0.0%
計	100.0%	-	100.0%	0.0%	100.0%	-	-	-	100.0%	
胃潰瘍	(人)	202	-	16	0	13	-	-	-	11
	ショックなし(収縮期血圧90mmHg以上)	87.6%	-	81.3%	0.0%	92.3%	-	-	-	100.0%
	ショックあり(収縮期血圧90mmHg未満)	11.9%	-	18.8%	0.0%	7.7%	-	-	-	0.0%
	不明	0.5%	-	0.0%	0.0%	0.0%	-	-	-	0.0%
計	100.0%	-	100.0%	0.0%	100.0%	-	-	-	100.0%	
急性腸炎	(人)	701	-	81	0	63	-	-	-	170
	血便なし	92.3%	-	98.8%	0.0%	87.3%	-	-	-	94.7%
	血便あり	6.6%	-	1.2%	0.0%	12.7%	-	-	-	5.3%
	不明	1.1%	-	0.0%	0.0%	0.0%	-	-	-	0.0%
計	100.0%	-	100.0%	0.0%	100.0%	-	-	-	100.0%	
急性虫垂炎	(人)	522	-	33	0	25	-	-	-	119
	白血球15000/mm3未満	67.4%	-	48.5%	0.0%	68.0%	-	-	-	66.4%
	白血球15000/mm3以上	32.2%	-	51.5%	0.0%	32.0%	-	-	-	32.8%
	不明	0.4%	-	0.0%	0.0%	0.0%	-	-	-	0.8%
計	100.0%	-	100.0%	0.0%	100.0%	-	-	-	100.0%	
胆石症	(人)	1,108	-	76	0	42	-	-	-	167
	黄疸なし(T Bill 5mg/dl 未満)	93.1%	-	94.7%	0.0%	88.1%	-	-	-	93.4%
	黄疸あり(T Bill 5mg/dl 以上)	6.9%	-	5.3%	0.0%	11.9%	-	-	-	6.6%
	不明	N/A	-	0.0%	0.0%	0.0%	-	-	-	0.0%
計	100.0%	-	100.0%	0.0%	100.0%	-	-	-	100.0%	
前立腺肥大症	(人)	95	-	0	0	1	-	-	-	13
	前立腺容積20ml未満	20.0%	-	0.0%	0.0%	0.0%	-	-	-	7.7%
	前立腺容積50ml未満	40.0%	-	0.0%	0.0%	0.0%	-	-	-	46.2%
	前立腺容積50ml以上	30.5%	-	0.0%	0.0%	0.0%	-	-	-	30.8%
	不明	9.5%	-	0.0%	0.0%	100.0%	-	-	-	15.4%
計	100.0%	-	0.0%	0.0%	100.0%	-	-	-	100.0%	
白内障	(人)	679	-	196	0	193	-	-	-	14
	1眼(処置対象のもの)	91.2%	-	97.4%	0.0%	100.0%	-	-	-	78.6%
	2眼(処置対象のもの)	8.8%	-	2.6%	0.0%	0.0%	-	-	-	21.4%
	不明	N/A	-	0.0%	0.0%	0.0%	-	-	-	0.0%
計	100.0%	-	100.0%	0.0%	100.0%	-	-	-	100.0%	
痔核	(人)	74	-	3	0	3	-	-	-	0
	第一度(出血のみ)	24.3%	-	33.3%	0.0%	0.0%	-	-	-	0.0%
	第二度(還納可能)	12.2%	-	0.0%	0.0%	0.0%	-	-	-	0.0%
	第三度(脱肛)	62.2%	-	66.7%	0.0%	100.0%	-	-	-	0.0%
	不明	1.4%	-	0.0%	0.0%	0.0%	-	-	-	0.0%
計	100.0%	-	100.0%	0.0%	100.0%	-	-	-	0.0%	
子宮筋腫	(人)	142	-	40	0	0	-	-	-	47
	貧血なし(Hb10g/dl以上)	65.5%	-	72.5%	0.0%	0.0%	-	-	-	55.3%
	貧血あり(Hb10g/dl未満)	33.8%	-	27.5%	0.0%	0.0%	-	-	-	42.6%
	不明	0.7%	-	0.0%	0.0%	0.0%	-	-	-	2.1%
計	100.0%	-	100.0%	0.0%	0.0%	-	-	-	100.0%	
狭心症	(人)	1,963	-	118	0	4	-	-	-	300
	重症度の指標なし	100.0%	-	100.0%	0.0%	100.0%	-	-	-	100.0%
	計	100.0%	-	100.0%	0.0%	100.0%	-	-	-	100.0%
腎結石及び尿管結石	(人)	487	-	7	0	3	-	-	-	101
	水腎・水尿管なし	35.7%	-	57.1%	0.0%	66.7%	-	-	-	33.7%
	水腎・水尿管あり	57.5%	-	42.9%	0.0%	33.3%	-	-	-	46.5%

【重症度別集計】表

		平均	病院コード							
			1041	1046	1047	1049	1058	1069	1129	1133
脳梗塞	JCS 200-300	2.6%	-	0.0%	-	-	-	33.3%	-	-
	不明	N/A	-	0.0%	-	-	-	0.0%	-	-
	計	100.0%	-	100.0%	-	-	-	100.0%	-	-
脳出血	(人)	586	-	16	-	-	-	2	-	-
	JCS 0-1	46.1%	-	87.5%	-	-	-	0.0%	-	-
	JCS 2-3	19.3%	-	6.3%	-	-	-	0.0%	-	-
	JCS 10-100	19.1%	-	6.3%	-	-	-	100.0%	-	-
	JCS 200-300	15.5%	-	0.0%	-	-	-	0.0%	-	-
	不明	N/A	-	0.0%	-	-	-	0.0%	-	-
	計	100.0%	-	100.0%	-	-	-	100.0%	-	-
糖尿病	(人)	585	-	21	-	-	-	31	-	-
	軽症	65.3%	-	66.7%	-	-	-	64.5%	-	-
	中等症	17.9%	-	14.3%	-	-	-	19.4%	-	-
	重症	11.1%	-	19.0%	-	-	-	12.9%	-	-
	不明	5.6%	-	0.0%	-	-	-	3.2%	-	-
	計	100.0%	-	100.0%	-	-	-	100.0%	-	-
大腿骨頸部骨折	(人)	511	-	25	-	-	-	1	-	-
	重症度Ⅰ	41.5%	-	12.0%	-	-	-	0.0%	-	-
	重症度Ⅱ	19.8%	-	56.0%	-	-	-	0.0%	-	-
	重症度Ⅲ	27.4%	-	28.0%	-	-	-	0.0%	-	-
	重症度Ⅳ	5.9%	-	0.0%	-	-	-	0.0%	-	-
	不明	5.5%	-	4.0%	-	-	-	100.0%	-	-
胃潰瘍	(人)	202	-	0	-	-	-	24	-	-
	ショックなし（収縮期血圧90mmHg以上）	87.6%	-	0.0%	-	-	-	75.0%	-	-
	ショックあり（収縮期血圧90mmHg未満）	11.9%	-	0.0%	-	-	-	25.0%	-	-
	不明	0.5%	-	0.0%	-	-	-	0.0%	-	-
	計	100.0%	-	0.0%	-	-	-	100.0%	-	-
急性腸炎	(人)	701	-	12	-	-	-	71	-	-
	血便なし	92.3%	-	83.3%	-	-	-	97.2%	-	-
	血便あり	6.6%	-	8.3%	-	-	-	2.8%	-	-
	不明	1.1%	-	8.3%	-	-	-	0.0%	-	-
急性虫垂炎	(人)	522	-	0	-	-	-	31	-	-
	白血球15000/mm3未満	67.4%	-	0.0%	-	-	-	67.7%	-	-
	白血球15000/mm3以上	32.2%	-	0.0%	-	-	-	32.3%	-	-
	不明	0.4%	-	0.0%	-	-	-	0.0%	-	-
	計	100.0%	-	0.0%	-	-	-	100.0%	-	-
胆石症	(人)	1,108	-	1	-	-	-	71	-	-
	黄疸なし(T Bill 5mg/dl 未満)	93.1%	-	100.0%	-	-	-	87.3%	-	-
	黄疸あり(T Bill 5mg/dl 以上)	6.9%	-	0.0%	-	-	-	12.7%	-	-
	不明	N/A	-	0.0%	-	-	-	0.0%	-	-
前立腺肥大症	(人)	95	-	0	-	-	-	0	-	-
	前立腺容積20ml未満	20.0%	-	0.0%	-	-	-	0.0%	-	-
	前立腺容積50ml未満	40.0%	-	0.0%	-	-	-	0.0%	-	-
	前立腺容積50ml以上	30.5%	-	0.0%	-	-	-	0.0%	-	-
	不明	9.5%	-	0.0%	-	-	-	0.0%	-	-
白内障	(人)	679	-	0	-	-	-	0	-	-
	1眼(処置対象のもの)	91.2%	-	0.0%	-	-	-	0.0%	-	-
	2眼(処置対象のもの)	8.8%	-	0.0%	-	-	-	0.0%	-	-
	不明	N/A	-	0.0%	-	-	-	0.0%	-	-
痔核	(人)	74	-	0	-	-	-	0	-	-
	第一度(出血のみ)	24.3%	-	0.0%	-	-	-	0.0%	-	-
	第二度(還納可能)	12.2%	-	0.0%	-	-	-	0.0%	-	-
	第三度(脱肛)	62.2%	-	0.0%	-	-	-	0.0%	-	-
	不明	1.4%	-	0.0%	-	-	-	0.0%	-	-
子宮筋腫	(人)	142	-	0	-	-	-	0	-	-
	貧血なし(Hb10g/dl以上)	65.5%	-	0.0%	-	-	-	0.0%	-	-
	貧血あり(Hb10g/dl未満)	33.8%	-	0.0%	-	-	-	0.0%	-	-
	不明	0.7%	-	0.0%	-	-	-	0.0%	-	-
狭心症	(人)	1,963	-	0	-	-	-	0	-	-
	重症度の指標なし	100.0%	-	0.0%	-	-	-	0.0%	-	-
	計	100.0%	-	0.0%	-	-	-	0.0%	-	-
腎結石及び尿管結石	(人)	487	-	0	-	-	-	1	-	-
	水腎・水尿管なし	35.7%	-	0.0%	-	-	-	0.0%	-	-
	水腎・水尿管あり	57.5%	-	0.0%	-	-	-	100.0%	-	-

【重症度別集計】表

		平均	病院コード						
			1167	1201	1202	9001	9004	9005	9007
脳梗塞	JCS 200-300	2.6%	-	-	-	-	-	-	-
	不明	N/A	-	-	-	-	-	-	-
	計	100.0%	-	-	-	-	-	-	-
脳出血	(人)	586	-	-	-	-	-	-	-
	JCS 0-1	46.1%	-	-	-	-	-	-	-
	JCS 2-3	19.3%	-	-	-	-	-	-	-
	JCS 10-100	19.1%	-	-	-	-	-	-	-
	JCS 200-300	15.5%	-	-	-	-	-	-	-
	不明	N/A	-	-	-	-	-	-	-
計	100.0%	-	-	-	-	-	-	-	
糖尿病	(人)	585	-	-	-	-	-	-	-
	軽症	65.3%	-	-	-	-	-	-	-
	中等症	17.9%	-	-	-	-	-	-	-
	重症	11.1%	-	-	-	-	-	-	-
	不明	5.6%	-	-	-	-	-	-	-
計	100.0%	-	-	-	-	-	-	-	
大腿骨頸部骨折	(人)	511	-	-	-	-	-	-	-
	重症度Ⅰ	41.5%	-	-	-	-	-	-	-
	重症度Ⅱ	19.8%	-	-	-	-	-	-	-
	重症度Ⅲ	27.4%	-	-	-	-	-	-	-
	重症度Ⅳ	5.9%	-	-	-	-	-	-	-
	不明	5.5%	-	-	-	-	-	-	-
計	100.0%	-	-	-	-	-	-	-	
胃潰瘍	(人)	202	-	-	-	-	-	-	-
	ショックなし(収縮期血圧90mmHg以上)	87.6%	-	-	-	-	-	-	-
	ショックあり(収縮期血圧90mmHg未満)	11.9%	-	-	-	-	-	-	-
	不明	0.5%	-	-	-	-	-	-	-
計	100.0%	-	-	-	-	-	-	-	
急性腸炎	(人)	701	-	-	-	-	-	-	-
	血便なし	92.3%	-	-	-	-	-	-	-
	血便あり	6.6%	-	-	-	-	-	-	-
	不明	1.1%	-	-	-	-	-	-	-
計	100.0%	-	-	-	-	-	-	-	
急性虫垂炎	(人)	522	-	-	-	-	-	-	-
	白血球15000/mm3未満	67.4%	-	-	-	-	-	-	-
	白血球15000/mm3以上	32.2%	-	-	-	-	-	-	-
	不明	0.4%	-	-	-	-	-	-	-
計	100.0%	-	-	-	-	-	-	-	
胆石症	(人)	1,108	-	-	-	-	-	-	-
	黄疸なし(T Bill 5mg/dl 未満)	93.1%	-	-	-	-	-	-	-
	黄疸あり(T Bill 5mg/dl 以上)	6.9%	-	-	-	-	-	-	-
	不明	N/A	-	-	-	-	-	-	-
計	100.0%	-	-	-	-	-	-	-	
前立腺肥大症	(人)	95	-	-	-	-	-	-	-
	前立腺容積20ml未満	20.0%	-	-	-	-	-	-	-
	前立腺容積50ml未満	40.0%	-	-	-	-	-	-	-
	前立腺容積50ml以上	30.5%	-	-	-	-	-	-	-
	不明	9.5%	-	-	-	-	-	-	-
計	100.0%	-	-	-	-	-	-	-	
白内障	(人)	679	-	-	-	-	-	-	-
	1眼(処置対象のもの)	91.2%	-	-	-	-	-	-	-
	2眼(処置対象のもの)	8.8%	-	-	-	-	-	-	-
	不明	N/A	-	-	-	-	-	-	-
計	100.0%	-	-	-	-	-	-	-	
痔核	(人)	74	-	-	-	-	-	-	-
	第一度(出血のみ)	24.3%	-	-	-	-	-	-	-
	第二度(還納可能)	12.2%	-	-	-	-	-	-	-
	第三度(脱肛)	62.2%	-	-	-	-	-	-	-
	不明	1.4%	-	-	-	-	-	-	-
計	100.0%	-	-	-	-	-	-	-	
子宮筋腫	(人)	142	-	-	-	-	-	-	-
	貧血なし(Hb10g/dl以上)	65.5%	-	-	-	-	-	-	-
	貧血あり(Hb10g/dl未満)	33.8%	-	-	-	-	-	-	-
	不明	0.7%	-	-	-	-	-	-	-
計	100.0%	-	-	-	-	-	-	-	
狭心症	(人)	1,963	-	-	-	-	-	-	-
	重症度の指標なし	100.0%	-	-	-	-	-	-	-
	計	100.0%	-	-	-	-	-	-	-
腎結石及び尿管結石	(人)	487	-	-	-	-	-	-	-
	水腎・水尿管なし	35.7%	-	-	-	-	-	-	-
	水腎・水尿管あり	57.5%	-	-	-	-	-	-	-

【重症度別集計】表

		平均	病院コード		
			9019	9020	9021
脳梗塞	JCS 200-300	2.6%	-	-	-
	不明	N/A	-	-	-
	計	100.0%	-	-	-
脳出血	(人)	586	-	-	-
	JCS 0-1	46.1%	-	-	-
	JCS 2-3	19.3%	-	-	-
	JCS 10-100	19.1%	-	-	-
	JCS 200-300	15.5%	-	-	-
	不明	N/A	-	-	-
	計	100.0%	-	-	-
糖尿病	(人)	585	-	-	-
	軽症	65.3%	-	-	-
	中等症	17.9%	-	-	-
	重症	11.1%	-	-	-
	不明	5.6%	-	-	-
	計	100.0%	-	-	-
大腿骨頸部骨折	(人)	511	-	-	-
	重症度Ⅰ	41.5%	-	-	-
	重症度Ⅱ	19.8%	-	-	-
	重症度Ⅲ	27.4%	-	-	-
	重症度Ⅳ	5.9%	-	-	-
	不明	5.5%	-	-	-
	計	100.0%	-	-	-
胃潰瘍	(人)	202	-	-	-
	ショックなし（収縮期血圧90mmHg以上）	87.6%	-	-	-
	ショックあり（収縮期血圧90mmHg未満）	11.9%	-	-	-
	不明	0.5%	-	-	-
	計	100.0%	-	-	-
急性腸炎	(人)	701	-	-	-
	血便なし	92.3%	-	-	-
	血便あり	6.6%	-	-	-
	不明	1.1%	-	-	-
	計	100.0%	-	-	-
急性虫垂炎	(人)	522	-	-	-
	白血球15000/mm3未満	67.4%	-	-	-
	白血球15000/mm3以上	32.2%	-	-	-
	不明	0.4%	-	-	-
	計	100.0%	-	-	-
胆石症	(人)	1,108	-	-	-
	黄疸なし(T Bill 5mg/dl 未満)	93.1%	-	-	-
	黄疸あり(T Bill 5mg/dl 以上)	6.9%	-	-	-
	不明	N/A	-	-	-
	計	100.0%	-	-	-
前立腺肥大症	(人)	95	-	-	-
	前立腺容積20ml未満	20.0%	-	-	-
	前立腺容積50ml未満	40.0%	-	-	-
	前立腺容積50ml以上	30.5%	-	-	-
	不明	9.5%	-	-	-
	計	100.0%	-	-	-
白内障	(人)	679	-	-	-
	1眼(処置対象のもの)	91.2%	-	-	-
	2眼(処置対象のもの)	8.8%	-	-	-
	不明	N/A	-	-	-
	計	100.0%	-	-	-
痔核	(人)	74	-	-	-
	第一度(出血のみ)	24.3%	-	-	-
	第二度(還納可能)	12.2%	-	-	-
	第三度(脱肛)	62.2%	-	-	-
	不明	1.4%	-	-	-
	計	100.0%	-	-	-
子宮筋腫	(人)	142	-	-	-
	貧血なし(Hb10g/dl以上)	65.5%	-	-	-
	貧血あり(Hb10g/dl未満)	33.8%	-	-	-
	不明	0.7%	-	-	-
	計	100.0%	-	-	-
狭心症	(人)	1,963	-	-	-
	重症度の指標なし	100.0%	-	-	-
	計	100.0%	-	-	-
腎結石及び尿管結石	(人)	487	-	-	-
	水腎・水尿管なし	35.7%	-	-	-
	水腎・水尿管あり	57.5%	-	-	-

【重症度別集計】表

		平均	病院コード		
			9019	9020	9021
腎結石及び尿管結石	不明	6.8%	-	-	-
	計	100.0%	-	-	-
乳房の悪性新生物	(人)	493	-	-	-
	ステージ0	6.5%	-	-	-
	ステージI	34.7%	-	-	-
	ステージII	36.1%	-	-	-
	ステージIII	8.9%	-	-	-
	ステージIV	8.3%	-	-	-
	不明	5.5%	-	-	-
計	100.0%	-	-	-	
膝関節症	(人)	313	-	-	-
	骨移植なし	100.0%	-	-	-
	骨移植あり	N/A	-	-	-
	不明	N/A	-	-	-
計	100.0%	-	-	-	
鼠径ヘルニア	(人)	727	-	-	-
	重症度の指標なし	100.0%	-	-	-
計	100.0%	-	-	-	
全疾患合計		100.0%	-	-	-
集計病院数		14			