

【重症度別集計】表

絞込条件

対象年月設定: 201604 ~ 201703
 病院の経営主体選択: 全て
 【詳細】病床数: 下限なし ~ 上限なし
 【詳細】手術の有無: 全て
 【詳細】ステージ集約選択: 有り
 【詳細】年代: 全て
 【詳細】性別: 全て
 【詳細】入院経路: 全て
 並べ替え条件: 病院コード・昇順

■ 退院患者数

	(単位:%)	平均	病院コード							
			0003	0202	0203	0235	1001	1002	1003	1004
胃の悪性新生物	(人)	578	65	41	4	62	110	30	183	-
ステージ0		3.6%	3.1%	0.0%	0.0%	19.4%	1.8%	0.0%	0.0%	-
ステージI		30.8%	26.2%	22.0%	0.0%	24.2%	32.7%	70.0%	37.7%	-
ステージII		11.6%	6.2%	17.1%	0.0%	9.7%	17.3%	3.3%	12.0%	-
ステージIII		14.5%	27.7%	12.2%	0.0%	11.3%	9.1%	0.0%	17.5%	-
ステージIV		29.2%	29.2%	41.5%	0.0%	29.0%	33.6%	23.3%	29.5%	-
不明		10.2%	7.7%	7.3%	100.0%	6.5%	5.5%	3.3%	3.3%	-
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
結腸の悪性新生物	(人)	946	99	31	4	180	300	42	114	-
ステージ0		2.3%	7.1%	3.2%	0.0%	1.1%	1.7%	0.0%	0.9%	-
ステージI		9.7%	18.2%	19.4%	0.0%	10.6%	6.3%	14.3%	11.4%	-
ステージII		16.8%	21.2%	12.9%	0.0%	21.1%	12.7%	28.6%	23.7%	-
ステージIII		33.4%	21.2%	16.1%	0.0%	29.4%	48.3%	23.8%	22.8%	-
ステージIV		31.6%	21.2%	35.5%	0.0%	35.6%	29.0%	23.8%	37.7%	-
不明		6.1%	11.1%	12.9%	100.0%	2.2%	2.0%	9.5%	3.5%	-
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
直腸の悪性新生物	(人)	523	38	18	7	110	129	12	86	-
ステージ0		3.8%	10.5%	0.0%	0.0%	5.5%	0.8%	0.0%	2.3%	-
ステージI		9.0%	13.2%	0.0%	0.0%	9.1%	0.8%	25.0%	24.4%	-
ステージII		16.3%	18.4%	27.8%	0.0%	23.6%	17.1%	25.0%	17.4%	-
ステージIII		22.8%	13.2%	61.1%	14.3%	26.4%	24.8%	33.3%	17.4%	-
ステージIV		43.6%	42.1%	0.0%	0.0%	33.6%	56.6%	16.7%	36.0%	-
不明		4.6%	2.6%	11.1%	85.7%	1.8%	0.0%	0.0%	2.3%	-
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
気管支および肺の悪性新生物	(人)	518	21	5	7	152	34	45	227	-
ステージ0		N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
ステージI		18.7%	0.0%	0.0%	0.0%	30.9%	26.5%	11.1%	15.9%	-
ステージII		4.1%	0.0%	20.0%	0.0%	7.2%	0.0%	8.9%	1.8%	-
ステージIII		21.6%	0.0%	60.0%	0.0%	8.6%	2.9%	13.3%	37.0%	-
ステージIV		35.7%	85.7%	20.0%	14.3%	36.8%	52.9%	22.2%	32.2%	-
不明		19.9%	14.3%	0.0%	85.7%	16.4%	17.6%	44.4%	13.2%	-
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
急性心筋梗塞	(人)	289	14	18	0	86	66	21	61	-
Killip I		53.6%	57.1%	22.2%	0.0%	65.1%	62.1%	0.0%	63.9%	-
Killip II		22.1%	7.1%	38.9%	0.0%	14.0%	16.7%	90.5%	18.0%	-
Killip III		8.0%	7.1%	22.2%	0.0%	3.5%	9.1%	0.0%	8.2%	-
Killip IV		14.5%	21.4%	16.7%	0.0%	15.1%	12.1%	4.8%	8.2%	-
不明		1.7%	7.1%	0.0%	0.0%	2.3%	0.0%	4.8%	1.6%	-
計		100.0%	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	-
肺炎	(人)	1,858	186	105	181	292	113	212	149	-
軽症		30.1%	31.7%	19.0%	49.2%	35.6%	36.3%	36.8%	26.8%	-
中等症		9.1%	12.4%	16.2%	14.4%	3.1%	10.6%	10.4%	6.0%	-
重症		44.0%	55.4%	64.8%	36.5%	61.3%	53.1%	52.8%	32.2%	-
不明		16.8%	0.5%	0.0%	0.0%	0.0%	0.0%	0.0%	34.9%	-
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
喘息	(人)	274	20	5	42	58	5	18	54	-
軽症間欠型		32.1%	25.0%	80.0%	23.8%	13.8%	80.0%	38.9%	13.0%	-
軽症持続型		24.1%	25.0%	0.0%	31.0%	51.7%	20.0%	22.2%	0.0%	-
中等症持続型		31.0%	40.0%	20.0%	40.5%	32.8%	0.0%	33.3%	44.4%	-
重症持続型		12.4%	5.0%	0.0%	4.8%	1.7%	0.0%	5.6%	42.6%	-
不明		0.4%	5.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
脳梗塞	(人)	1,184	82	566	15	284	8	61	53	-
JCS 0-1		62.3%	76.8%	52.1%	60.0%	73.2%	100.0%	68.9%	66.0%	-
JCS 2-3		27.4%	18.3%	38.3%	13.3%	18.7%	0.0%	18.0%	11.3%	-
JCS 10-100		8.0%	3.7%	7.8%	26.7%	6.0%	0.0%	11.5%	13.2%	-

【重症度別集計】表

	(単位:%)	平均	病院コード							
			1012	1013	1015	1018	1019	1025	1039	1046
胃の悪性新生物	(人)	578	30	-	18	-	-	-	-	4
ステージ0		3.6%	0.0%	-	11.1%	-	-	-	-	0.0%
ステージI		30.8%	23.3%	-	16.7%	-	-	-	-	0.0%
ステージII		11.6%	6.7%	-	5.6%	-	-	-	-	0.0%
ステージIII		14.5%	13.3%	-	0.0%	-	-	-	-	0.0%
ステージIV		29.2%	13.3%	-	38.9%	-	-	-	-	0.0%
不明		10.2%	43.3%	-	27.8%	-	-	-	-	100.0%
計		100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
結腸の悪性新生物	(人)	946	36	-	19	-	-	-	-	2
ステージ0		2.3%	5.6%	-	0.0%	-	-	-	-	0.0%
ステージI		9.7%	19.4%	-	0.0%	-	-	-	-	0.0%
ステージII		16.8%	5.6%	-	21.1%	-	-	-	-	0.0%
ステージIII		33.4%	19.4%	-	10.5%	-	-	-	-	0.0%
ステージIV		31.6%	25.0%	-	15.8%	-	-	-	-	0.0%
不明		6.1%	25.0%	-	52.6%	-	-	-	-	100.0%
計		100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
直腸の悪性新生物	(人)	523	7	-	12	-	-	-	-	0
ステージ0		3.8%	14.3%	-	0.0%	-	-	-	-	0.0%
ステージI		9.0%	0.0%	-	0.0%	-	-	-	-	0.0%
ステージII		16.3%	28.6%	-	16.7%	-	-	-	-	0.0%
ステージIII		22.8%	42.9%	-	8.3%	-	-	-	-	0.0%
ステージIV		43.6%	0.0%	-	16.7%	-	-	-	-	0.0%
不明		4.6%	14.3%	-	58.3%	-	-	-	-	0.0%
計		100.0%	100.0%	-	100.0%	-	-	-	-	0.0%
気管支および肺の悪性新生物	(人)	518	10	-	9	-	-	-	-	6
ステージ0		N/A	0.0%	-	0.0%	-	-	-	-	0.0%
ステージI		18.7%	0.0%	-	0.0%	-	-	-	-	0.0%
ステージII		4.1%	10.0%	-	0.0%	-	-	-	-	0.0%
ステージIII		21.6%	40.0%	-	0.0%	-	-	-	-	16.7%
ステージIV		35.7%	10.0%	-	44.4%	-	-	-	-	16.7%
不明		19.9%	40.0%	-	55.6%	-	-	-	-	66.7%
計		100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
急性心筋梗塞	(人)	289	11	-	4	-	-	-	-	6
Killip I		53.6%	54.5%	-	25.0%	-	-	-	-	0.0%
Killip II		22.1%	9.1%	-	25.0%	-	-	-	-	16.7%
Killip III		8.0%	9.1%	-	50.0%	-	-	-	-	16.7%
Killip IV		14.5%	27.3%	-	0.0%	-	-	-	-	66.7%
不明		1.7%	0.0%	-	0.0%	-	-	-	-	0.0%
計		100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
肺炎	(人)	1,858	137	-	355	-	-	-	-	50
軽症		30.1%	46.7%	-	11.5%	-	-	-	-	24.0%
中等症		9.1%	10.9%	-	2.5%	-	-	-	-	24.0%
重症		44.0%	42.3%	-	13.0%	-	-	-	-	52.0%
不明		16.8%	0.0%	-	73.0%	-	-	-	-	0.0%
計		100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
喘息	(人)	274	24	-	38	-	-	-	-	0
軽症間欠型		32.1%	75.0%	-	65.8%	-	-	-	-	0.0%
軽症持続型		24.1%	4.2%	-	21.1%	-	-	-	-	0.0%
中等症持続型		31.0%	12.5%	-	2.6%	-	-	-	-	0.0%
重症持続型		12.4%	8.3%	-	10.5%	-	-	-	-	0.0%
不明		0.4%	0.0%	-	0.0%	-	-	-	-	0.0%
計		100.0%	100.0%	-	100.0%	-	-	-	-	0.0%
脳梗塞	(人)	1,184	48	-	31	-	-	-	-	34
JCS 0-1		62.3%	68.8%	-	51.6%	-	-	-	-	79.4%
JCS 2-3		27.4%	25.0%	-	19.4%	-	-	-	-	8.8%
JCS 10-100		8.0%	4.2%	-	25.8%	-	-	-	-	8.8%

【重症度別集計】表

	(単位:%)	平均	病院コード							
			1047	1069	1129	1167	1201	1202	1501	1502
胃の悪性新生物	(人)	578	-	31	-	-	-	-	-	-
ステージ0		3.6%	-	9.7%	-	-	-	-	-	-
ステージI		30.8%	-	3.2%	-	-	-	-	-	-
ステージII		11.6%	-	16.1%	-	-	-	-	-	-
ステージIII		14.5%	-	25.8%	-	-	-	-	-	-
ステージIV		29.2%	-	19.4%	-	-	-	-	-	-
不明		10.2%	-	25.8%	-	-	-	-	-	-
計		100.0%	-	100.0%	-	-	-	-	-	-
結腸の悪性新生物	(人)	946	-	119	-	-	-	-	-	-
ステージ0		2.3%	-	3.4%	-	-	-	-	-	-
ステージI		9.7%	-	3.4%	-	-	-	-	-	-
ステージII		16.8%	-	10.9%	-	-	-	-	-	-
ステージIII		33.4%	-	39.5%	-	-	-	-	-	-
ステージIV		31.6%	-	42.9%	-	-	-	-	-	-
不明		6.1%	-	0.0%	-	-	-	-	-	-
計		100.0%	-	100.0%	-	-	-	-	-	-
直腸の悪性新生物	(人)	523	-	104	-	-	-	-	-	-
ステージ0		3.8%	-	5.8%	-	-	-	-	-	-
ステージI		9.0%	-	6.7%	-	-	-	-	-	-
ステージII		16.3%	-	2.9%	-	-	-	-	-	-
ステージIII		22.8%	-	17.3%	-	-	-	-	-	-
ステージIV		43.6%	-	64.4%	-	-	-	-	-	-
不明		4.6%	-	2.9%	-	-	-	-	-	-
計		100.0%	-	100.0%	-	-	-	-	-	-
気管支および肺の悪性新生物	(人)	518	-	2	-	-	-	-	-	-
ステージ0		N/A	-	0.0%	-	-	-	-	-	-
ステージI		18.7%	-	0.0%	-	-	-	-	-	-
ステージII		4.1%	-	0.0%	-	-	-	-	-	-
ステージIII		21.6%	-	0.0%	-	-	-	-	-	-
ステージIV		35.7%	-	100.0%	-	-	-	-	-	-
不明		19.9%	-	0.0%	-	-	-	-	-	-
計		100.0%	-	100.0%	-	-	-	-	-	-
急性心筋梗塞	(人)	289	-	2	-	-	-	-	-	-
Killip I		53.6%	-	0.0%	-	-	-	-	-	-
Killip II		22.1%	-	0.0%	-	-	-	-	-	-
Killip III		8.0%	-	0.0%	-	-	-	-	-	-
Killip IV		14.5%	-	100.0%	-	-	-	-	-	-
不明		1.7%	-	0.0%	-	-	-	-	-	-
計		100.0%	-	100.0%	-	-	-	-	-	-
肺炎	(人)	1,858	-	78	-	-	-	-	-	-
軽症		30.1%	-	15.4%	-	-	-	-	-	-
中等症		9.1%	-	19.2%	-	-	-	-	-	-
重症		44.0%	-	65.4%	-	-	-	-	-	-
不明		16.8%	-	0.0%	-	-	-	-	-	-
計		100.0%	-	100.0%	-	-	-	-	-	-
喘息	(人)	274	-	10	-	-	-	-	-	-
軽症間欠型		32.1%	-	0.0%	-	-	-	-	-	-
軽症持続型		24.1%	-	40.0%	-	-	-	-	-	-
中等症持続型		31.0%	-	60.0%	-	-	-	-	-	-
重症持続型		12.4%	-	0.0%	-	-	-	-	-	-
不明		0.4%	-	0.0%	-	-	-	-	-	-
計		100.0%	-	100.0%	-	-	-	-	-	-
脳梗塞	(人)	1,184	-	2	-	-	-	-	-	-
JCS 0-1		62.3%	-	100.0%	-	-	-	-	-	-
JCS 2-3		27.4%	-	0.0%	-	-	-	-	-	-
JCS 10-100		8.0%	-	0.0%	-	-	-	-	-	-

【重症度別集計】表

		平均	病院コード							
			0003	0202	0203	0235	1001	1002	1003	1004
脳梗塞	JCS 200-300	2.2%	1.2%	1.8%	0.0%	2.1%	0.0%	1.6%	9.4%	-
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
脳出血	(人)	480	21	216	3	153	1	21	12	-
	JCS 0-1	40.8%	33.3%	34.7%	33.3%	50.3%	0.0%	28.6%	75.0%	-
	JCS 2-3	21.5%	23.8%	24.1%	0.0%	17.6%	0.0%	23.8%	8.3%	-
	JCS 10-100	20.8%	38.1%	20.4%	33.3%	17.6%	0.0%	33.3%	8.3%	-
	JCS 200-300	16.9%	4.8%	20.8%	33.3%	14.4%	100.0%	14.3%	8.3%	-
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-	
糖尿病	(人)	348	59	29	7	64	64	29	26	-
	軽症	53.4%	55.9%	86.2%	57.1%	53.1%	42.2%	27.6%	84.6%	-
	中等症	31.6%	35.6%	13.8%	42.9%	20.3%	43.8%	51.7%	3.8%	-
	重症	14.9%	8.5%	0.0%	0.0%	26.6%	14.1%	20.7%	11.5%	-
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-	
大腿骨頸部骨折	(人)	393	122	0	1	121	1	2	24	-
	重症度Ⅰ	46.3%	47.5%	0.0%	0.0%	62.0%	100.0%	0.0%	37.5%	-
	重症度Ⅱ	26.5%	43.4%	0.0%	0.0%	19.8%	0.0%	0.0%	12.5%	-
	重症度Ⅲ	20.4%	6.6%	0.0%	0.0%	10.7%	0.0%	0.0%	50.0%	-
	重症度Ⅳ	6.4%	2.5%	0.0%	100.0%	7.4%	0.0%	0.0%	0.0%	-
	不明	0.5%	0.0%	0.0%	0.0%	0.0%	0.0%	100.0%	0.0%	-
計	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-	
胃潰瘍	(人)	138	18	12	1	24	15	4	25	-
	ショックなし(収縮期血圧90mmHg以上)	89.9%	88.9%	100.0%	100.0%	91.7%	86.7%	100.0%	88.0%	-
	ショックあり(収縮期血圧90mmHg未満)	10.1%	11.1%	0.0%	0.0%	8.3%	13.3%	0.0%	12.0%	-
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-	
急性腸炎	(人)	559	113	43	23	46	60	44	59	-
	血便なし	90.5%	91.2%	81.4%	100.0%	89.1%	93.3%	84.1%	91.5%	-
	血便あり	9.3%	8.8%	18.6%	0.0%	10.9%	6.7%	15.9%	8.5%	-
	不明	0.2%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-	
急性虫垂炎	(人)	377	71	11	0	81	46	23	75	-
	白血球15000/mm3未満	73.5%	74.6%	90.9%	0.0%	72.8%	78.3%	56.5%	68.0%	-
	白血球15000/mm3以上	26.3%	25.4%	9.1%	0.0%	27.2%	21.7%	43.5%	32.0%	-
	不明	0.3%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
計	100.0%	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	-	
胆石症	(人)	955	138	42	9	200	130	68	198	-
	黄疸なし(T Bill 5mg/dl 未満)	91.6%	85.5%	83.3%	88.9%	96.0%	90.8%	98.5%	89.9%	-
	黄疸あり(T Bill 5mg/dl 以上)	8.4%	14.5%	16.7%	11.1%	4.0%	9.2%	1.5%	10.1%	-
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-	
前立腺肥大症	(人)	54	22	0	0	0	3	1	28	-
	前立腺容積20ml未満	1.9%	4.5%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
	前立腺容積50ml未満	31.5%	27.3%	0.0%	0.0%	0.0%	0.0%	0.0%	39.3%	-
	前立腺容積50ml以上	50.0%	45.5%	0.0%	0.0%	0.0%	33.3%	0.0%	57.1%	-
	不明	16.7%	22.7%	0.0%	0.0%	0.0%	66.7%	100.0%	3.6%	-
計	100.0%	100.0%	0.0%	0.0%	0.0%	100.0%	100.0%	100.0%	-	
白内障	(人)	588	278	0	0	0	0	0	0	-
	1眼(処置対象のもの)	99.8%	99.6%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
	2眼(処置対象のもの)	0.2%	0.4%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
計	100.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-	
痔核	(人)	109	4	1	5	17	60	3	3	-
	第一度(出血のみ)	25.7%	50.0%	100.0%	80.0%	23.5%	18.3%	33.3%	33.3%	-
	第二度(還納可能)	21.1%	0.0%	0.0%	0.0%	17.6%	25.0%	66.7%	0.0%	-
	第三度(脱肛)	50.5%	50.0%	0.0%	0.0%	52.9%	56.7%	0.0%	66.7%	-
	不明	2.8%	0.0%	0.0%	20.0%	5.9%	0.0%	0.0%	0.0%	-
計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-	
子宮筋腫	(人)	115	83	17	0	0	0	0	2	-
	貧血なし(Hb10g/dl以上)	81.7%	85.5%	52.9%	0.0%	0.0%	0.0%	0.0%	100.0%	-
	貧血あり(Hb10g/dl未満)	18.3%	14.5%	47.1%	0.0%	0.0%	0.0%	0.0%	0.0%	-
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
計	100.0%	100.0%	100.0%	0.0%	0.0%	0.0%	0.0%	100.0%	-	
狭心症	(人)	1,649	177	124	1	335	198	155	598	-
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
腎結石及び尿管結石	(人)	352	155	1	3	11	81	4	94	-
	水腎・水尿管なし	51.1%	56.1%	100.0%	66.7%	72.7%	39.5%	100.0%	46.8%	-
	水腎・水尿管あり	46.9%	43.9%	0.0%	33.3%	27.3%	53.1%	0.0%	52.1%	-

【重症度別集計】表

		平均	病院コード							
			1012	1013	1015	1018	1019	1025	1039	1046
脳梗塞	JCS 200-300	2.2%	2.1%	-	3.2%	-	-	-	-	2.9%
	不明	N/A	0.0%	-	0.0%	-	-	-	-	0.0%
	計	100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
脳出血	(人)	480	21	-	9	-	-	-	-	21
	JCS 0-1	40.8%	23.8%	-	55.6%	-	-	-	-	52.4%
	JCS 2-3	21.5%	23.8%	-	44.4%	-	-	-	-	19.0%
	JCS 10-100	20.8%	33.3%	-	0.0%	-	-	-	-	14.3%
	JCS 200-300	16.9%	19.0%	-	0.0%	-	-	-	-	14.3%
	不明	N/A	0.0%	-	0.0%	-	-	-	-	0.0%
	計	100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
糖尿病	(人)	348	5	-	34	-	-	-	-	9
	軽症	53.4%	60.0%	-	44.1%	-	-	-	-	44.4%
	中等症	31.6%	20.0%	-	35.3%	-	-	-	-	55.6%
	重症	14.9%	20.0%	-	20.6%	-	-	-	-	0.0%
	不明	N/A	0.0%	-	0.0%	-	-	-	-	0.0%
	計	100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
大腿骨頸部骨折	(人)	393	20	-	95	-	-	-	-	7
	重症度Ⅰ	46.3%	65.0%	-	25.3%	-	-	-	-	28.6%
	重症度Ⅱ	26.5%	20.0%	-	16.8%	-	-	-	-	57.1%
	重症度Ⅲ	20.4%	15.0%	-	45.3%	-	-	-	-	14.3%
	重症度Ⅳ	6.4%	0.0%	-	12.6%	-	-	-	-	0.0%
	不明	0.5%	0.0%	-	0.0%	-	-	-	-	0.0%
	計	100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
胃潰瘍	(人)	138	8	-	10	-	-	-	-	1
	ショックなし(収縮期血圧90mmHg以上)	89.9%	100.0%	-	100.0%	-	-	-	-	100.0%
	ショックあり(収縮期血圧90mmHg未満)	10.1%	0.0%	-	0.0%	-	-	-	-	0.0%
	不明	N/A	0.0%	-	0.0%	-	-	-	-	0.0%
	計	100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
急性腸炎	(人)	559	32	-	74	-	-	-	-	3
	血便なし	90.5%	100.0%	-	89.2%	-	-	-	-	100.0%
	血便あり	9.3%	0.0%	-	10.8%	-	-	-	-	0.0%
	不明	0.2%	0.0%	-	0.0%	-	-	-	-	0.0%
	計	100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
急性虫垂炎	(人)	377	16	-	16	-	-	-	-	1
	白血球15000/mm3未満	73.5%	81.3%	-	62.5%	-	-	-	-	100.0%
	白血球15000/mm3以上	26.3%	18.8%	-	37.5%	-	-	-	-	0.0%
	不明	0.3%	0.0%	-	0.0%	-	-	-	-	0.0%
	計	100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
胆石症	(人)	955	54	-	30	-	-	-	-	1
	黄疸なし(T Bill 5mg/dl 未満)	91.6%	98.1%	-	90.0%	-	-	-	-	100.0%
	黄疸あり(T Bill 5mg/dl 以上)	8.4%	1.9%	-	10.0%	-	-	-	-	0.0%
	不明	N/A	0.0%	-	0.0%	-	-	-	-	0.0%
	計	100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
前立腺肥大症	(人)	54	0	-	0	-	-	-	-	0
	前立腺容積20ml未満	1.9%	0.0%	-	0.0%	-	-	-	-	0.0%
	前立腺容積50ml未満	31.5%	0.0%	-	0.0%	-	-	-	-	0.0%
	前立腺容積50ml以上	50.0%	0.0%	-	0.0%	-	-	-	-	0.0%
	不明	16.7%	0.0%	-	0.0%	-	-	-	-	0.0%
	計	100.0%	0.0%	-	0.0%	-	-	-	-	0.0%
白内障	(人)	588	109	-	201	-	-	-	-	0
	1眼(処置対象のもの)	99.8%	100.0%	-	100.0%	-	-	-	-	0.0%
	2眼(処置対象のもの)	0.2%	0.0%	-	0.0%	-	-	-	-	0.0%
	不明	N/A	0.0%	-	0.0%	-	-	-	-	0.0%
	計	100.0%	100.0%	-	100.0%	-	-	-	-	0.0%
痔核	(人)	109	7	-	4	-	-	-	-	0
	第一度(出血のみ)	25.7%	0.0%	-	75.0%	-	-	-	-	0.0%
	第二度(還納可能)	21.1%	28.6%	-	25.0%	-	-	-	-	0.0%
	第三度(脱肛)	50.5%	71.4%	-	0.0%	-	-	-	-	0.0%
	不明	2.8%	0.0%	-	0.0%	-	-	-	-	0.0%
	計	100.0%	100.0%	-	100.0%	-	-	-	-	0.0%
子宮筋腫	(人)	115	13	-	0	-	-	-	-	0
	貧血なし(Hb10g/dl以上)	81.7%	92.3%	-	0.0%	-	-	-	-	0.0%
	貧血あり(Hb10g/dl未満)	18.3%	7.7%	-	0.0%	-	-	-	-	0.0%
	不明	N/A	0.0%	-	0.0%	-	-	-	-	0.0%
	計	100.0%	100.0%	-	0.0%	-	-	-	-	0.0%
狭心症	(人)	1,649	59	-	1	-	-	-	-	0
	重症度の指標なし	100.0%	100.0%	-	100.0%	-	-	-	-	0.0%
	計	100.0%	100.0%	-	100.0%	-	-	-	-	0.0%
腎結石及び尿管結石	(人)	352	1	-	1	-	-	-	-	0
	水腎・水尿管なし	51.1%	100.0%	-	0.0%	-	-	-	-	0.0%
	水腎・水尿管あり	46.9%	0.0%	-	100.0%	-	-	-	-	0.0%

【重症度別集計】表

		平均	病院コード							
			1047	1069	1129	1167	1201	1202	1501	1502
脳梗塞	JCS 200-300	2.2%	-	0.0%	-	-	-	-	-	-
	不明	N/A	-	0.0%	-	-	-	-	-	-
	計	100.0%	-	100.0%	-	-	-	-	-	-
脳出血	(人)	480	-	2	-	-	-	-	-	-
	JCS 0-1	40.8%	-	0.0%	-	-	-	-	-	-
	JCS 2-3	21.5%	-	0.0%	-	-	-	-	-	-
	JCS 10-100	20.8%	-	100.0%	-	-	-	-	-	-
	JCS 200-300	16.9%	-	0.0%	-	-	-	-	-	-
	不明	N/A	-	0.0%	-	-	-	-	-	-
	計	100.0%	-	100.0%	-	-	-	-	-	-
糖尿病	(人)	348	-	22	-	-	-	-	-	-
	軽症	53.4%	-	50.0%	-	-	-	-	-	-
	中等症	31.6%	-	31.8%	-	-	-	-	-	-
	重症	14.9%	-	18.2%	-	-	-	-	-	-
	計	100.0%	-	100.0%	-	-	-	-	-	-
大腿骨頸部骨折	(人)	393	-	0	-	-	-	-	-	-
	重症度Ⅰ	46.3%	-	0.0%	-	-	-	-	-	-
	重症度Ⅱ	26.5%	-	0.0%	-	-	-	-	-	-
	重症度Ⅲ	20.4%	-	0.0%	-	-	-	-	-	-
	重症度Ⅳ	6.4%	-	0.0%	-	-	-	-	-	-
	不明	0.5%	-	0.0%	-	-	-	-	-	-
胃潰瘍	(人)	138	-	20	-	-	-	-	-	-
	ショックなし(収縮期血圧90mmHg以上)	89.9%	-	75.0%	-	-	-	-	-	-
	ショックあり(収縮期血圧90mmHg未満)	10.1%	-	25.0%	-	-	-	-	-	-
	計	100.0%	-	100.0%	-	-	-	-	-	-
急性腸炎	(人)	559	-	62	-	-	-	-	-	-
	血便なし	90.5%	-	90.3%	-	-	-	-	-	-
	血便あり	9.3%	-	8.1%	-	-	-	-	-	-
	計	100.0%	-	100.0%	-	-	-	-	-	-
急性虫垂炎	(人)	377	-	37	-	-	-	-	-	-
	白血球15000/mm3未満	73.5%	-	83.8%	-	-	-	-	-	-
	白血球15000/mm3以上	26.3%	-	13.5%	-	-	-	-	-	-
	計	100.0%	-	100.0%	-	-	-	-	-	-
胆石症	(人)	955	-	85	-	-	-	-	-	-
	黄疸なし(T Bill 5mg/dl 未満)	91.6%	-	91.8%	-	-	-	-	-	-
	黄疸あり(T Bill 5mg/dl 以上)	8.4%	-	8.2%	-	-	-	-	-	-
	計	100.0%	-	100.0%	-	-	-	-	-	-
前立腺肥大症	(人)	54	-	0	-	-	-	-	-	-
	前立腺容積20ml未満	1.9%	-	0.0%	-	-	-	-	-	-
	前立腺容積50ml未満	31.5%	-	0.0%	-	-	-	-	-	-
	前立腺容積50ml以上	50.0%	-	0.0%	-	-	-	-	-	-
	計	100.0%	-	0.0%	-	-	-	-	-	-
白内障	(人)	588	-	0	-	-	-	-	-	-
	1眼(処置対象のもの)	99.8%	-	0.0%	-	-	-	-	-	-
	2眼(処置対象のもの)	0.2%	-	0.0%	-	-	-	-	-	-
	計	100.0%	-	0.0%	-	-	-	-	-	-
痔核	(人)	109	-	5	-	-	-	-	-	-
	第一度(出血のみ)	25.7%	-	20.0%	-	-	-	-	-	-
	第二度(還納可能)	21.1%	-	0.0%	-	-	-	-	-	-
	第三度(脱肛)	50.5%	-	60.0%	-	-	-	-	-	-
	計	100.0%	-	100.0%	-	-	-	-	-	-
子宮筋腫	(人)	115	-	0	-	-	-	-	-	-
	貧血なし(Hb10g/dl以上)	81.7%	-	0.0%	-	-	-	-	-	-
	貧血あり(Hb10g/dl未満)	18.3%	-	0.0%	-	-	-	-	-	-
	計	100.0%	-	0.0%	-	-	-	-	-	-
狭心症	(人)	1,649	-	1	-	-	-	-	-	-
	重症度の指標なし	100.0%	-	100.0%	-	-	-	-	-	-
	計	100.0%	-	100.0%	-	-	-	-	-	-
腎結石及び尿管結石	(人)	352	-	1	-	-	-	-	-	-
	水腎・水尿管なし	51.1%	-	100.0%	-	-	-	-	-	-
	水腎・水尿管あり	46.9%	-	0.0%	-	-	-	-	-	-

