

【重症度別集計】表

絞込条件

対象年月設定: 201704 ~ 201803
 病院の経営主体選択: 全て
 【詳細】病床数: 下限なし ~ 上限なし
 【詳細】手術の有無: 全て
 【詳細】ステージ集約選択: 有り
 【詳細】年代: 全て
 【詳細】性別: 全て
 【詳細】入院経路: 全て
 並べ替え条件: 病院コード・昇順

■ 退院患者数

	(単位:%)	平均	病院コード							
			0003	0202	0203	0235	1001	1002	1003	1004
胃の悪性新生物	(人)	491	69	57	9	30	84	24	173	-
ステージ0		0.2%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
ステージI		32.6%	24.6%	36.8%	0.0%	30.0%	38.1%	12.5%	37.0%	-
ステージII		11.2%	4.3%	7.0%	0.0%	10.0%	16.7%	4.2%	14.5%	-
ステージIII		13.0%	20.3%	10.5%	0.0%	16.7%	9.5%	16.7%	13.3%	-
ステージIV		36.3%	43.5%	40.4%	0.0%	43.3%	33.3%	58.3%	34.1%	-
不明		6.7%	7.2%	5.3%	100.0%	0.0%	2.4%	8.3%	1.2%	-
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
結腸の悪性新生物	(人)	755	86	47	5	119	255	56	137	-
ステージ0		1.6%	0.0%	2.1%	0.0%	1.7%	0.4%	5.4%	2.2%	-
ステージI		7.2%	11.6%	8.5%	20.0%	6.7%	3.9%	5.4%	10.2%	-
ステージII		19.3%	27.9%	23.4%	0.0%	14.3%	18.8%	14.3%	25.5%	-
ステージIII		34.6%	15.1%	40.4%	0.0%	37.0%	48.6%	23.2%	29.9%	-
ステージIV		28.9%	32.6%	19.1%	0.0%	37.0%	26.7%	26.8%	29.9%	-
不明		8.5%	12.8%	6.4%	80.0%	3.4%	1.6%	25.0%	2.2%	-
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
直腸の悪性新生物	(人)	424	21	21	3	100	150	20	91	-
ステージ0		2.4%	0.0%	9.5%	0.0%	2.0%	0.7%	0.0%	5.5%	-
ステージI		12.5%	19.0%	9.5%	0.0%	23.0%	2.0%	25.0%	14.3%	-
ステージII		15.3%	19.0%	0.0%	33.3%	14.0%	12.7%	20.0%	24.2%	-
ステージIII		26.2%	33.3%	57.1%	33.3%	24.0%	21.3%	25.0%	27.5%	-
ステージIV		39.4%	14.3%	23.8%	0.0%	36.0%	62.7%	20.0%	25.3%	-
不明		4.2%	14.3%	0.0%	33.3%	1.0%	0.7%	10.0%	3.3%	-
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
気管支および肺の悪性新生物	(人)	515	12	2	3	184	74	36	188	-
ステージ0		1.2%	0.0%	0.0%	0.0%	1.6%	2.7%	0.0%	0.5%	-
ステージI		18.6%	0.0%	0.0%	0.0%	28.8%	14.9%	27.8%	11.7%	-
ステージII		8.0%	0.0%	100.0%	0.0%	7.6%	5.4%	13.9%	8.5%	-
ステージIII		18.8%	8.3%	0.0%	0.0%	19.6%	8.1%	2.8%	27.7%	-
ステージIV		38.1%	75.0%	0.0%	0.0%	32.1%	56.8%	19.4%	39.4%	-
不明		15.3%	16.7%	0.0%	100.0%	10.3%	12.2%	36.1%	12.2%	-
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
急性心筋梗塞	(人)	306	16	18	4	81	64	19	69	-
Killip I		52.0%	62.5%	50.0%	50.0%	66.7%	18.8%	5.3%	69.6%	-
Killip II		28.8%	6.3%	22.2%	0.0%	16.0%	65.6%	73.7%	11.6%	-
Killip III		6.2%	0.0%	11.1%	0.0%	4.9%	6.3%	15.8%	4.3%	-
Killip IV		10.1%	31.3%	11.1%	0.0%	9.9%	7.8%	5.3%	10.1%	-
不明		2.9%	0.0%	5.6%	50.0%	2.5%	1.6%	0.0%	4.3%	-
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
肺炎	(人)	1,687	175	139	119	299	134	218	144	-
軽症		29.9%	25.7%	24.5%	50.4%	31.8%	35.8%	30.7%	19.4%	-
中等症		9.3%	12.0%	15.1%	16.8%	4.3%	9.0%	10.1%	6.3%	-
重症		48.0%	62.3%	60.4%	32.8%	63.9%	55.2%	59.2%	52.8%	-
不明		12.8%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	21.5%	-
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
喘息	(人)	257	9	9	19	95	4	16	48	-
軽症間欠型		30.7%	33.3%	55.6%	21.1%	23.2%	50.0%	0.0%	4.2%	-
軽症持続型		22.6%	33.3%	33.3%	26.3%	38.9%	0.0%	18.8%	0.0%	-
中等症持続型		31.5%	33.3%	0.0%	31.6%	30.5%	50.0%	81.3%	52.1%	-
重症持続型		12.5%	0.0%	11.1%	21.1%	2.1%	0.0%	0.0%	43.8%	-
不明		2.7%	0.0%	0.0%	0.0%	5.3%	0.0%	0.0%	0.0%	-
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
脳梗塞	(人)	1,218	45	624	6	266	4	68	65	-
JCS 0-1		65.6%	68.9%	62.8%	33.3%	63.2%	100.0%	80.9%	76.9%	-
JCS 2-3		24.1%	20.0%	28.2%	16.7%	25.6%	0.0%	5.9%	13.8%	-
JCS 10-100		8.5%	8.9%	7.7%	33.3%	9.8%	0.0%	8.8%	6.2%	-

【重症度別集計】表

	(単位:%)	平均	病院コード							
			1012	1013	1015	1018	1019	1025	1039	1046
胃の悪性新生物	(人)	491	31	-	13	-	-	-	-	1
ステージ0		0.2%	0.0%	-	7.7%	-	-	-	-	0.0%
ステージI		32.6%	32.3%	-	23.1%	-	-	-	-	100.0%
ステージII		11.2%	9.7%	-	15.4%	-	-	-	-	0.0%
ステージIII		13.0%	6.5%	-	15.4%	-	-	-	-	0.0%
ステージIV		36.3%	22.6%	-	30.8%	-	-	-	-	0.0%
不明		6.7%	29.0%	-	7.7%	-	-	-	-	0.0%
計		100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
結腸の悪性新生物	(人)	755	34	-	15	-	-	-	-	1
ステージ0		1.6%	5.9%	-	0.0%	-	-	-	-	0.0%
ステージI		7.2%	11.8%	-	0.0%	-	-	-	-	0.0%
ステージII		19.3%	8.8%	-	0.0%	-	-	-	-	0.0%
ステージIII		34.6%	20.6%	-	0.0%	-	-	-	-	0.0%
ステージIV		28.9%	38.2%	-	0.0%	-	-	-	-	0.0%
不明		8.5%	14.7%	-	100.0%	-	-	-	-	100.0%
計		100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
直腸の悪性新生物	(人)	424	6	-	11	-	-	-	-	1
ステージ0		2.4%	0.0%	-	0.0%	-	-	-	-	0.0%
ステージI		12.5%	50.0%	-	0.0%	-	-	-	-	0.0%
ステージII		15.3%	16.7%	-	0.0%	-	-	-	-	0.0%
ステージIII		26.2%	33.3%	-	27.3%	-	-	-	-	0.0%
ステージIV		39.4%	0.0%	-	18.2%	-	-	-	-	0.0%
不明		4.2%	0.0%	-	54.5%	-	-	-	-	100.0%
計		100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
気管支および肺の悪性新生物	(人)	515	15	-	1	-	-	-	-	0
ステージ0		1.2%	0.0%	-	0.0%	-	-	-	-	0.0%
ステージI		18.6%	0.0%	-	0.0%	-	-	-	-	0.0%
ステージII		8.0%	0.0%	-	0.0%	-	-	-	-	0.0%
ステージIII		18.8%	6.7%	-	0.0%	-	-	-	-	0.0%
ステージIV		38.1%	26.7%	-	100.0%	-	-	-	-	0.0%
不明		15.3%	66.7%	-	0.0%	-	-	-	-	0.0%
計		100.0%	100.0%	-	100.0%	-	-	-	-	0.0%
急性心筋梗塞	(人)	306	29	-	2	-	-	-	-	4
Killip I		52.0%	75.9%	-	0.0%	-	-	-	-	25.0%
Killip II		28.8%	13.8%	-	50.0%	-	-	-	-	25.0%
Killip III		6.2%	6.9%	-	50.0%	-	-	-	-	0.0%
Killip IV		10.1%	3.4%	-	0.0%	-	-	-	-	50.0%
不明		2.9%	0.0%	-	0.0%	-	-	-	-	0.0%
計		100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
肺炎	(人)	1,687	107	-	304	-	-	-	-	48
軽症		29.9%	62.6%	-	14.1%	-	-	-	-	37.5%
中等症		9.3%	9.3%	-	6.9%	-	-	-	-	16.7%
重症		48.0%	28.0%	-	18.1%	-	-	-	-	45.8%
不明		12.8%	0.0%	-	60.9%	-	-	-	-	0.0%
計		100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
喘息	(人)	257	25	-	30	-	-	-	-	2
軽症間欠型		30.7%	72.0%	-	76.7%	-	-	-	-	0.0%
軽症持続型		22.6%	16.0%	-	10.0%	-	-	-	-	0.0%
中等症持続型		31.5%	8.0%	-	3.3%	-	-	-	-	0.0%
重症持続型		12.5%	4.0%	-	10.0%	-	-	-	-	0.0%
不明		2.7%	0.0%	-	0.0%	-	-	-	-	100.0%
計		100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
脳梗塞	(人)	1,218	59	-	56	-	-	-	-	25
JCS 0-1		65.6%	61.0%	-	73.2%	-	-	-	-	80.0%
JCS 2-3		24.1%	27.1%	-	19.6%	-	-	-	-	0.0%
JCS 10-100		8.5%	6.8%	-	7.1%	-	-	-	-	20.0%

【重症度別集計】表

		平均	病院コード							
			0003	0202	0203	0235	1001	1002	1003	1004
脳梗塞	JCS 200-300	1.8%	2.2%	1.3%	16.7%	1.5%	0.0%	4.4%	3.1%	-
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
脳出血	(人)	432	19	188	1	142	3	14	16	-
	JCS 0-1	37.7%	15.8%	34.0%	0.0%	40.1%	66.7%	14.3%	68.8%	-
	JCS 2-3	20.8%	42.1%	21.3%	0.0%	20.4%	0.0%	21.4%	6.3%	-
	JCS 10-100	24.1%	42.1%	27.7%	100.0%	19.0%	33.3%	28.6%	6.3%	-
	JCS 200-300	17.4%	0.0%	17.0%	0.0%	20.4%	0.0%	35.7%	18.8%	-
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
糖尿病	(人)	345	60	38	5	58	83	29	27	-
	軽症	60.9%	76.7%	47.4%	100.0%	50.0%	75.9%	34.5%	66.7%	-
	中等症	23.8%	15.0%	50.0%	0.0%	22.4%	15.7%	48.3%	11.1%	-
	重症	13.3%	8.3%	2.6%	0.0%	27.6%	8.4%	17.2%	18.5%	-
	不明	2.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	3.7%	-
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
大腿骨頸部骨折	(人)	379	73	1	0	131	2	3	53	-
	重症度Ⅰ	41.7%	42.5%	0.0%	0.0%	48.9%	50.0%	0.0%	49.1%	-
	重症度Ⅱ	27.7%	50.7%	100.0%	0.0%	29.0%	0.0%	0.0%	9.4%	-
	重症度Ⅲ	22.2%	4.1%	0.0%	0.0%	16.0%	0.0%	0.0%	26.4%	-
	重症度Ⅳ	6.9%	2.7%	0.0%	0.0%	5.3%	50.0%	0.0%	15.1%	-
	不明	1.6%	0.0%	0.0%	0.0%	0.8%	0.0%	100.0%	0.0%	-
	計	100.0%	100.0%	100.0%	0.0%	100.0%	100.0%	100.0%	100.0%	-
胃潰瘍	(人)	133	14	16	1	38	8	8	28	-
	ショックなし(収縮期血圧90mmHg以上)	92.5%	100.0%	100.0%	100.0%	89.5%	87.5%	87.5%	89.3%	-
	ショックあり(収縮期血圧90mmHg未満)	7.5%	0.0%	0.0%	0.0%	10.5%	12.5%	12.5%	10.7%	-
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
急性腸炎	(人)	487	103	44	20	63	49	52	53	-
	血便なし	91.2%	93.2%	95.5%	100.0%	81.0%	89.8%	84.6%	92.5%	-
	血便あり	8.8%	6.8%	4.5%	0.0%	19.0%	10.2%	15.4%	7.5%	-
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
急性虫垂炎	(人)	368	86	19	2	89	45	23	80	-
	白血球15000/mm3未満	71.7%	66.3%	68.4%	50.0%	67.4%	75.6%	78.3%	76.3%	-
	白血球15000/mm3以上	28.3%	33.7%	31.6%	50.0%	32.6%	24.4%	21.7%	23.8%	-
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
胆石症	(人)	910	101	56	9	228	175	43	230	-
	黄疸なし(T Bill 5mg/dl 未満)	91.4%	85.1%	91.1%	88.9%	96.5%	81.1%	93.0%	95.2%	-
	黄疸あり(T Bill 5mg/dl 以上)	8.6%	14.9%	8.9%	11.1%	3.5%	18.9%	7.0%	4.8%	-
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
前立腺肥大症	(人)	43	21	0	0	1	3	0	18	-
	前立腺容積20ml未満	11.6%	9.5%	0.0%	0.0%	0.0%	0.0%	0.0%	16.7%	-
	前立腺容積50ml未満	32.6%	23.8%	0.0%	0.0%	0.0%	33.3%	0.0%	44.4%	-
	前立腺容積50ml以上	41.9%	47.6%	0.0%	0.0%	0.0%	66.7%	0.0%	33.3%	-
	不明	14.0%	19.0%	0.0%	0.0%	100.0%	0.0%	0.0%	5.6%	-
	計	100.0%	100.0%	0.0%	0.0%	100.0%	100.0%	0.0%	100.0%	-
白内障	(人)	530	264	0	0	0	0	0	0	-
	1眼(処置対象のもの)	99.8%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
	2眼(処置対象のもの)	0.2%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
	計	100.0%	100.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
痔核	(人)	105	6	2	4	36	43	4	6	-
	第一度(出血のみ)	20.0%	33.3%	50.0%	75.0%	13.9%	18.6%	25.0%	16.7%	-
	第二度(還納可能)	37.1%	33.3%	50.0%	0.0%	25.0%	55.8%	0.0%	33.3%	-
	第三度(脱肛)	41.9%	33.3%	0.0%	0.0%	61.1%	25.6%	75.0%	50.0%	-
	不明	1.0%	0.0%	0.0%	25.0%	0.0%	0.0%	0.0%	0.0%	-
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
子宮筋腫	(人)	118	76	21	0	0	0	0	5	-
	貧血なし(Hb10g/dl以上)	58.5%	51.3%	71.4%	0.0%	0.0%	0.0%	0.0%	60.0%	-
	貧血あり(Hb10g/dl未満)	41.5%	48.7%	28.6%	0.0%	0.0%	0.0%	0.0%	40.0%	-
	不明	N/A	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	-
	計	100.0%	100.0%	100.0%	0.0%	0.0%	0.0%	0.0%	100.0%	-
狭心症	(人)	1,534	164	112	3	342	174	141	544	-
	重症度の指標なし	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
	計	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	-
腎結石及び尿管結石	(人)	306	148	0	1	3	85	3	59	-
	水腎・水尿管なし	47.7%	52.0%	0.0%	0.0%	66.7%	28.2%	66.7%	61.0%	-
	水腎・水尿管あり	52.3%	48.0%	0.0%	100.0%	33.3%	71.8%	33.3%	39.0%	-

【重症度別集計】表

		平均	病院コード							
			1012	1013	1015	1018	1019	1025	1039	1046
脳梗塞	JCS 200-300	1.8%	5.1%	-	0.0%	-	-	-	-	0.0%
	不明	N/A	0.0%	-	0.0%	-	-	-	-	0.0%
	計	100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
脳出血	(人)	432	27	-	8	-	-	-	-	14
	JCS 0-1	37.7%	37.0%	-	62.5%	-	-	-	-	64.3%
	JCS 2-3	20.8%	18.5%	-	25.0%	-	-	-	-	14.3%
	JCS 10-100	24.1%	33.3%	-	0.0%	-	-	-	-	7.1%
	JCS 200-300	17.4%	11.1%	-	12.5%	-	-	-	-	14.3%
	不明	N/A	0.0%	-	0.0%	-	-	-	-	0.0%
	計	100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
糖尿病	(人)	345	10	-	25	-	-	-	-	10
	軽症	60.9%	30.0%	-	56.0%	-	-	-	-	40.0%
	中等症	23.8%	40.0%	-	24.0%	-	-	-	-	10.0%
	重症	13.3%	10.0%	-	20.0%	-	-	-	-	10.0%
	不明	2.0%	20.0%	-	0.0%	-	-	-	-	40.0%
	計	100.0%	100.0%	-	100.0%	-	-	-	-	100.0%
大腿骨頸部骨折	(人)	379	20	-	92	-	-	-	-	4
	重症度Ⅰ	41.7%	60.0%	-	26.1%	-	-	-	-	0.0%
	重症度Ⅱ	27.7%	25.0%	-	18.5%	-	-	-	-	50.0%
	重症度Ⅲ	22.2%	10.0%	-	47.8%	-	-	-	-	0.0%
	重症度Ⅳ	6.9%	5.0%	-	7.6%	-	-	-	-	0.0%
	不明	1.6%	0.0%	-	0.0%	-	-	-	-	50.0%
計	100.0%	100.0%	-	100.0%	-	-	-	-	100.0%	
胃潰瘍	(人)	133	5	-	13	-	-	-	-	2
	ショックなし(収縮期血圧90mmHg以上)	92.5%	100.0%	-	92.3%	-	-	-	-	100.0%
	ショックあり(収縮期血圧90mmHg未満)	7.5%	0.0%	-	7.7%	-	-	-	-	0.0%
	不明	N/A	0.0%	-	0.0%	-	-	-	-	0.0%
計	100.0%	100.0%	-	100.0%	-	-	-	-	100.0%	
急性腸炎	(人)	487	35	-	64	-	-	-	-	4
	血便なし	91.2%	100.0%	-	92.2%	-	-	-	-	100.0%
	血便あり	8.8%	0.0%	-	7.8%	-	-	-	-	0.0%
	不明	N/A	0.0%	-	0.0%	-	-	-	-	0.0%
計	100.0%	100.0%	-	100.0%	-	-	-	-	100.0%	
急性虫垂炎	(人)	368	12	-	11	-	-	-	-	1
	白血球15000/mm3未満	71.7%	83.3%	-	81.8%	-	-	-	-	100.0%
	白血球15000/mm3以上	28.3%	16.7%	-	18.2%	-	-	-	-	0.0%
	不明	N/A	0.0%	-	0.0%	-	-	-	-	0.0%
計	100.0%	100.0%	-	100.0%	-	-	-	-	100.0%	
胆石症	(人)	910	44	-	23	-	-	-	-	1
	黄疸なし(T Bill 5mg/dl 未満)	91.4%	97.7%	-	95.7%	-	-	-	-	100.0%
	黄疸あり(T Bill 5mg/dl 以上)	8.6%	2.3%	-	4.3%	-	-	-	-	0.0%
	不明	N/A	0.0%	-	0.0%	-	-	-	-	0.0%
計	100.0%	100.0%	-	100.0%	-	-	-	-	100.0%	
前立腺肥大症	(人)	43	0	-	0	-	-	-	-	0
	前立腺容積20ml未満	11.6%	0.0%	-	0.0%	-	-	-	-	0.0%
	前立腺容積50ml未満	32.6%	0.0%	-	0.0%	-	-	-	-	0.0%
	前立腺容積50ml以上	41.9%	0.0%	-	0.0%	-	-	-	-	0.0%
	不明	14.0%	0.0%	-	0.0%	-	-	-	-	0.0%
計	100.0%	0.0%	-	0.0%	-	-	-	-	0.0%	
白内障	(人)	530	77	-	189	-	-	-	-	0
	1眼(処置対象のもの)	99.8%	100.0%	-	99.5%	-	-	-	-	0.0%
	2眼(処置対象のもの)	0.2%	0.0%	-	0.5%	-	-	-	-	0.0%
	不明	N/A	0.0%	-	0.0%	-	-	-	-	0.0%
計	100.0%	100.0%	-	100.0%	-	-	-	-	0.0%	
痔核	(人)	105	2	-	2	-	-	-	-	0
	第一度(出血のみ)	20.0%	0.0%	-	0.0%	-	-	-	-	0.0%
	第二度(還納可能)	37.1%	0.0%	-	50.0%	-	-	-	-	0.0%
	第三度(脱肛)	41.9%	100.0%	-	50.0%	-	-	-	-	0.0%
	不明	1.0%	0.0%	-	0.0%	-	-	-	-	0.0%
計	100.0%	100.0%	-	100.0%	-	-	-	-	0.0%	
子宮筋腫	(人)	118	16	-	0	-	-	-	-	0
	貧血なし(Hb10g/dl以上)	58.5%	75.0%	-	0.0%	-	-	-	-	0.0%
	貧血あり(Hb10g/dl未満)	41.5%	25.0%	-	0.0%	-	-	-	-	0.0%
	不明	N/A	0.0%	-	0.0%	-	-	-	-	0.0%
計	100.0%	100.0%	-	0.0%	-	-	-	-	0.0%	
狭心症	(人)	1,534	52	-	2	-	-	-	-	0
	重症度の指標なし	100.0%	100.0%	-	100.0%	-	-	-	-	0.0%
	計	100.0%	100.0%	-	100.0%	-	-	-	-	0.0%
腎結石及び尿管結石	(人)	306	1	-	6	-	-	-	-	0
	水腎・水尿管なし	47.7%	100.0%	-	66.7%	-	-	-	-	0.0%
	水腎・水尿管あり	52.3%	0.0%	-	33.3%	-	-	-	-	0.0%

