

【重症度別集計】表

絞込条件

対象年月設定: 202004 ~ 202103
 病院の経営主体選択: 全て
 【詳細】病床数: 下限なし ~ 上限なし
 【詳細】手術の有無: 全て
 【詳細】ステージ集約選択: 有り
 【詳細】年代: 全て
 【詳細】性別: 全て
 【詳細】入院経路: 全て
 並べ替え条件: 病院コード・昇順

■ 退院患者数

	(単位:%)	平均	病院コード							
			0003	0202	0203	0235	1001	1002	1003	1004
胃の悪性新生物	(人)	-	-	57	-	60	-	-	-	-
ステージ0		2.5%	-	0.0%	0.0%	5.0%	-	-	-	-
ステージI		40.3%	-	42.1%	0.0%	40.0%	-	-	-	-
ステージII		10.9%	-	12.3%	0.0%	10.0%	-	-	-	-
ステージIII		10.1%	-	10.5%	50.0%	8.3%	-	-	-	-
ステージIV		28.6%	-	26.3%	0.0%	31.7%	-	-	-	-
不明		7.6%	-	8.8%	50.0%	5.0%	-	-	-	-
計		100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
結腸の悪性新生物	(人)	-	-	70	-	-	-	-	-	-
ステージ0		15.2%	-	8.6%	0.0%	18.2%	-	-	-	-
ステージI		11.3%	-	7.1%	0.0%	13.2%	-	-	-	-
ステージII		17.3%	-	22.9%	50.0%	14.5%	-	-	-	-
ステージIII		27.3%	-	34.3%	0.0%	24.5%	-	-	-	-
ステージIV		27.3%	-	24.3%	0.0%	28.9%	-	-	-	-
不明		1.7%	-	2.9%	50.0%	0.6%	-	-	-	-
計		100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
直腸の悪性新生物	(人)	-	-	35	-	-	-	-	-	-
ステージ0		5.6%	-	8.6%	0.0%	4.6%	-	-	-	-
ステージI		25.0%	-	28.6%	0.0%	24.1%	-	-	-	-
ステージII		15.3%	-	5.7%	0.0%	19.5%	-	-	-	-
ステージIII		21.0%	-	28.6%	0.0%	18.4%	-	-	-	-
ステージIV		31.5%	-	28.6%	0.0%	33.3%	-	-	-	-
不明		1.6%	-	0.0%	100.0%	0.0%	-	-	-	-
計		100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
気管支および肺の悪性新生物	(人)	-	-	11	-	155	-	-	-	-
ステージ0		2.3%	-	0.0%	0.0%	2.6%	-	-	-	-
ステージI		30.9%	-	36.4%	0.0%	32.3%	-	-	-	-
ステージII		4.6%	-	0.0%	0.0%	5.2%	-	-	-	-
ステージIII		14.9%	-	36.4%	0.0%	14.2%	-	-	-	-
ステージIV		28.0%	-	18.2%	0.0%	30.3%	-	-	-	-
不明		19.4%	-	9.1%	100.0%	15.5%	-	-	-	-
計		100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
急性心筋梗塞	(人)	-	-	28	-	-	-	-	-	-
Killip I		55.4%	-	17.9%	0.0%	69.9%	-	-	-	-
Killip II		7.9%	-	7.1%	0.0%	8.2%	-	-	-	-
Killip III		5.9%	-	3.6%	0.0%	6.8%	-	-	-	-
Killip IV		29.7%	-	71.4%	0.0%	13.7%	-	-	-	-
不明		1.0%	-	0.0%	0.0%	1.4%	-	-	-	-
計		100.0%	-	100.0%	0.0%	100.0%	-	-	-	-
肺炎	(人)	-	-	114	-	-	-	-	-	-
軽症		45.4%	-	32.5%	71.2%	35.5%	-	-	-	-
中等症		10.9%	-	18.4%	12.6%	3.5%	-	-	-	-
重症		43.7%	-	49.1%	16.2%	61.0%	-	-	-	-
不明		N/A	-	0.0%	0.0%	0.0%	-	-	-	-
計		100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
喘息	(人)	-	-	0	-	-	-	-	-	-
軽症間欠型		26.5%	-	0.0%	35.3%	17.6%	-	-	-	-
軽症持続型		26.5%	-	0.0%	23.5%	29.4%	-	-	-	-
中等症持続型		29.4%	-	0.0%	17.6%	41.2%	-	-	-	-
重症持続型		14.7%	-	0.0%	23.5%	5.9%	-	-	-	-
不明		2.9%	-	0.0%	0.0%	5.9%	-	-	-	-
計		100.0%	-	0.0%	100.0%	100.0%	-	-	-	-
脳梗塞	(人)	-	-	667	-	-	-	-	-	-
JCS 0-1		67.8%	-	61.5%	100.0%	84.9%	-	-	-	-
JCS 2-3		24.6%	-	29.4%	0.0%	11.7%	-	-	-	-
JCS 10-100		5.1%	-	6.0%	0.0%	2.5%	-	-	-	-

【重症度別集計】表

	(単位:%)	平均	病院コード	
			9022	9023
胃の悪性新生物	(人)	-	-	-
ステージ0		2.5%	-	-
ステージI		40.3%	-	-
ステージII		10.9%	-	-
ステージIII		10.1%	-	-
ステージIV		28.6%	-	-
不明		7.6%	-	-
計		100.0%	-	-
結腸の悪性新生物	(人)	-	-	-
ステージ0		15.2%	-	-
ステージI		11.3%	-	-
ステージII		17.3%	-	-
ステージIII		27.3%	-	-
ステージIV		27.3%	-	-
不明		1.7%	-	-
計		100.0%	-	-
直腸の悪性新生物	(人)	-	-	-
ステージ0		5.6%	-	-
ステージI		25.0%	-	-
ステージII		15.3%	-	-
ステージIII		21.0%	-	-
ステージIV		31.5%	-	-
不明		1.6%	-	-
計		100.0%	-	-
気管支および肺の悪性新生物	(人)	-	-	-
ステージ0		2.3%	-	-
ステージI		30.9%	-	-
ステージII		4.6%	-	-
ステージIII		14.9%	-	-
ステージIV		28.0%	-	-
不明		19.4%	-	-
計		100.0%	-	-
急性心筋梗塞	(人)	-	-	-
Killip I		55.4%	-	-
Killip II		7.9%	-	-
Killip III		5.9%	-	-
Killip IV		29.7%	-	-
不明		1.0%	-	-
計		100.0%	-	-
肺炎	(人)	-	-	-
軽症		45.4%	-	-
中等症		10.9%	-	-
重症		43.7%	-	-
不明		N/A	-	-
計		100.0%	-	-
喘息	(人)	-	-	-
軽症間欠型		26.5%	-	-
軽症持続型		26.5%	-	-
中等症持続型		29.4%	-	-
重症持続型		14.7%	-	-
不明		2.9%	-	-
計		100.0%	-	-
脳梗塞	(人)	-	-	-
JCS 0-1		67.8%	-	-
JCS 2-3		24.6%	-	-
JCS 10-100		5.1%	-	-

【重症度別集計】表

		平均	病院コード							
			0003	0202	0203	0235	1001	1002	1003	1004
脳梗塞	JCS 200-300	2.5%	-	3.1%	0.0%	0.8%	-	-	-	-
	不明	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
	計	100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
脳出血	(人)	-	-	245	-	-	-	-	-	-
	JCS 0-1	48.3%	-	37.6%	0.0%	66.7%	-	-	-	-
	JCS 2-3	22.9%	-	28.6%	0.0%	13.2%	-	-	-	-
	JCS 10-100	10.8%	-	12.2%	0.0%	8.3%	-	-	-	-
	JCS 200-300	18.0%	-	21.6%	0.0%	11.8%	-	-	-	-
	不明	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
	計	100.0%	-	100.0%	0.0%	100.0%	-	-	-	-
糖尿病	(人)	-	-	19	-	-	-	-	-	-
	軽症	34.3%	-	42.1%	40.0%	30.2%	-	-	-	-
	中等症	37.3%	-	52.6%	60.0%	27.9%	-	-	-	-
	重症	28.4%	-	5.3%	0.0%	41.9%	-	-	-	-
	不明	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
	計	100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
大腿骨頸部骨折	(人)	-	-	54	-	-	-	-	-	-
	重症度Ⅰ	54.7%	-	38.9%	0.0%	62.4%	-	-	-	-
	重症度Ⅱ	24.4%	-	40.7%	0.0%	17.1%	-	-	-	-
	重症度Ⅲ	16.3%	-	13.0%	0.0%	17.9%	-	-	-	-
	重症度Ⅳ	4.1%	-	7.4%	0.0%	2.6%	-	-	-	-
	不明	0.6%	-	0.0%	100.0%	0.0%	-	-	-	-
	計	100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
胃潰瘍	(人)	-	-	14	-	-	-	-	-	-
	ショックなし(収縮期血圧90mmHg以上)	79.1%	-	100.0%	0.0%	69.0%	-	-	-	-
	ショックあり(収縮期血圧90mmHg未満)	20.9%	-	0.0%	0.0%	31.0%	-	-	-	-
	不明	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
	計	100.0%	-	100.0%	0.0%	100.0%	-	-	-	-
急性腸炎	(人)	-	-	31	-	-	-	-	-	-
	血便なし	87.5%	-	87.1%	100.0%	71.4%	-	-	-	-
	血便あり	12.5%	-	12.9%	0.0%	28.6%	-	-	-	-
	不明	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
	計	100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
急性虫垂炎	(人)	-	-	21	-	-	-	-	-	-
	白血球15000/mm3未満	73.9%	-	71.4%	0.0%	100.0%	-	-	-	-
	白血球15000/mm3以上	26.1%	-	28.6%	0.0%	0.0%	-	-	-	-
	不明	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
	計	100.0%	-	100.0%	0.0%	100.0%	-	-	-	-
胆石症	(人)	-	-	69	-	-	-	-	-	-
	黄疸なし(T Bill 5mg/dl 未満)	93.9%	-	84.1%	100.0%	96.7%	-	-	-	-
	黄疸あり(T Bill 5mg/dl 以上)	6.1%	-	15.9%	0.0%	3.3%	-	-	-	-
	不明	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
	計	100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
前立腺肥大症	(人)	-	-	0	-	-	-	-	-	-
	前立腺容積20ml未満	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
	前立腺容積50ml未満	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
	前立腺容積50ml以上	50.0%	-	0.0%	0.0%	50.0%	-	-	-	-
	不明	50.0%	-	0.0%	0.0%	50.0%	-	-	-	-
	計	100.0%	-	0.0%	0.0%	100.0%	-	-	-	-
白内障	(人)	-	-	0	-	-	-	-	-	-
	1眼(処置対象のもの)	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
	2眼(処置対象のもの)	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
	不明	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
	計	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
痔核	(人)	-	-	1	-	-	-	-	-	-
	第一度(出血のみ)	66.7%	-	100.0%	50.0%	0.0%	-	-	-	-
	第二度(還納可能)	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
	第三度(脱肛)	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
	不明	33.3%	-	0.0%	50.0%	0.0%	-	-	-	-
	計	100.0%	-	100.0%	100.0%	0.0%	-	-	-	-
子宮筋腫	(人)	-	-	27	-	-	-	-	-	-
	貧血なし(Hb10g/dl以上)	70.4%	-	70.4%	0.0%	0.0%	-	-	-	-
	貧血あり(Hb10g/dl未満)	29.6%	-	29.6%	0.0%	0.0%	-	-	-	-
	不明	N/A	-	0.0%	0.0%	0.0%	-	-	-	-
	計	100.0%	-	100.0%	0.0%	0.0%	-	-	-	-
狭心症	(人)	-	-	95	-	338	-	-	-	-
	重症度の指標なし	100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
	計	100.0%	-	100.0%	100.0%	100.0%	-	-	-	-
腎結石及び尿管結石	(人)	-	-	2	-	-	-	-	-	-
	水腎・水尿管なし	64.3%	-	100.0%	0.0%	63.6%	-	-	-	-
	水腎・水尿管あり	35.7%	-	0.0%	100.0%	36.4%	-	-	-	-

【重症度別集計】表

		平均	病院コード	
			9022	9023
脳梗塞	JCS 200-300	2.5%	-	-
	不明	N/A	-	-
	計	100.0%	-	-
脳出血	(人)	-	-	-
	JCS 0-1	48.3%	-	-
	JCS 2-3	22.9%	-	-
	JCS 10-100	10.8%	-	-
	JCS 200-300	18.0%	-	-
	不明	N/A	-	-
	計	100.0%	-	-
糖尿病	(人)	-	-	-
	軽症	34.3%	-	-
	中等症	37.3%	-	-
	重症	28.4%	-	-
	不明	N/A	-	-
	計	100.0%	-	-
大腿骨頸部骨折	(人)	-	-	-
	重症度Ⅰ	54.7%	-	-
	重症度Ⅱ	24.4%	-	-
	重症度Ⅲ	16.3%	-	-
	重症度Ⅳ	4.1%	-	-
	不明	0.6%	-	-
	計	100.0%	-	-
胃潰瘍	(人)	-	-	-
	ショックなし(収縮期血圧90mmHg以上)	79.1%	-	-
	ショックあり(収縮期血圧90mmHg未満)	20.9%	-	-
	不明	N/A	-	-
	計	100.0%	-	-
急性腸炎	(人)	-	-	-
	血便なし	87.5%	-	-
	血便あり	12.5%	-	-
	不明	N/A	-	-
	計	100.0%	-	-
急性虫垂炎	(人)	-	-	-
	白血球15000/mm3未満	73.9%	-	-
	白血球15000/mm3以上	26.1%	-	-
	不明	N/A	-	-
	計	100.0%	-	-
胆石症	(人)	-	-	-
	黄疸なし(T Bill 5mg/dl 未満)	93.9%	-	-
	黄疸あり(T Bill 5mg/dl 以上)	6.1%	-	-
	不明	N/A	-	-
	計	100.0%	-	-
前立腺肥大症	(人)	-	-	-
	前立腺容積20ml未満	N/A	-	-
	前立腺容積50ml未満	N/A	-	-
	前立腺容積50ml以上	50.0%	-	-
	不明	50.0%	-	-
	計	100.0%	-	-
白内障	(人)	-	-	-
	1眼(処置対象のもの)	N/A	-	-
	2眼(処置対象のもの)	N/A	-	-
	不明	N/A	-	-
	計	N/A	-	-
痔核	(人)	-	-	-
	第一度(出血のみ)	66.7%	-	-
	第二度(還納可能)	N/A	-	-
	第三度(脱肛)	N/A	-	-
	不明	33.3%	-	-
	計	100.0%	-	-
子宮筋腫	(人)	-	-	-
	貧血なし(Hb10g/dl以上)	70.4%	-	-
	貧血あり(Hb10g/dl未満)	29.6%	-	-
	不明	N/A	-	-
	計	100.0%	-	-
狭心症	(人)	-	-	-
	重症度の指標なし	100.0%	-	-
	計	100.0%	-	-
腎結石及び尿管結石	(人)	-	-	-
	水腎・水尿管なし	64.3%	-	-
	水腎・水尿管あり	35.7%	-	-

【重症度別集計】表

		平均	病院コード	
			9022	9023
腎結石及び尿管結石	不明	N/A	-	-
	計	100.0%	-	-
乳房の悪性新生物	(人)	-	-	-
	ステージ0	13.6%	-	-
	ステージI	38.0%	-	-
	ステージII	34.8%	-	-
	ステージIII	8.2%	-	-
	ステージIV	3.8%	-	-
	不明	1.6%	-	-
計	100.0%	-	-	
膝関節症	(人)	-	-	-
	骨移植なし	100.0%	-	-
	骨移植あり	N/A	-	-
	不明	N/A	-	-
計	100.0%	-	-	
鼠径ヘルニア	(人)	-	-	-
	重症度の指標なし	100.0%	-	-
計	100.0%	-	-	
全疾患合計		100.0%	-	-
集計病院数		3		